

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> :  A61K	A2	(11) International Publication Number: <b>WO 00/38618</b>  (43) International Publication Date: 6 July 2000 (06.07.00)
<p>(21) International Application Number: PCT/US99/30815</p> <p>(22) International Filing Date: 23 December 1999 (23.12.99)</p> <p>(30) Priority Data: 60/113,588 24 December 1998 (24.12.98) US</p> <p>(71) Applicant: DU PONT PHARMACEUTICALS COMPANY [US/US]; Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19807 (US).</p> <p>(72) Inventor: OLSON, Richard, E.; 7 Pelham Road, Wilmington, DE 19803 (US).</p> <p>(74) Agent: LARSEN, Scott, K.; Du Pont Pharmaceuticals Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).</p>	<p>(81) Designated States: AL, AU, BR, CA, CN, CZ, EE, HU, IL, IN, JP, KR, LT, LV, MK, MX, NO, NZ, PL, RO, SG, SI, SK, TR, UA, VN, ZA, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p><b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i></p>	

(54) Title: SUCCINOYLAMINO BENZODIAZEPINES AS INHIBITORS OF A $\beta$  PROTEIN PRODUCTION



(57) Abstract

This invention relates to novel lactams having formula (I) to their pharmaceutical compositions and to their methods of use. These novel compounds inhibit the processing of amyloid precursor protein and, more specifically, inhibit the production of A $\beta$ -peptide, thereby acting to prevent the formation of neurological deposits of amyloid protein. More particularly, the present invention relates to the treatment of neurological disorders related to  $\beta$ -amyloid production such as Alzheimer's disease and Down's Syndrome.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with 0-1 R<sup>11a</sup>;  
phenyl substituted with 0-3 R<sup>11b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>11b</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
5 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>11b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
10 pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>11a</sup>, at each occurrence, is independently selected from H,  
C<sub>1</sub>-C<sub>4</sub> alkyl, OR<sup>14</sup>, F, Cl, =O, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, or phenyl  
15 substituted with 0-3 R<sup>11b</sup>;

R<sup>11b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, methyl, ethyl, propyl, butyl,  
methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub>  
20 haloalkoxy;

Z is H;  
C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-3 R<sup>12a</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>12a</sup>; or  
25 C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>12a</sup>;

R<sup>12a</sup>, at each occurrence, is independently selected from  
H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
30 ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

R<sup>13</sup>, at each occurrence, is independently selected from  
H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
Cl, F, Br, CN, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;

35 R<sup>14</sup> is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;


R<sup>15</sup>, at each occurrence, is independently selected from H, methyl, ethyl, propyl, and butyl;

R<sup>16</sup>, at each occurrence, is independently selected from  
 5 H, OH, methyl, ethyl, propyl, butyl, benzyl, phenethyl, methyl-C(=O)-, ethyl-C(=O)-, methyl-S(=O)<sub>2</sub>-, and ethyl-S(=O)<sub>2</sub>-;

R<sup>18</sup>, at each occurrence, is independently selected from  
 10 H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl; and

R<sup>19</sup>, at each occurrence, is independently selected from  
 15 H, methyl, and ethyl.

8. A compound according to Claim 5 of Formula (Ie):



(Ie)

20 or a pharmaceutically acceptable salt or prodrug thereof wherein:

R<sup>3</sup> is R<sup>4</sup>,

25 R<sup>4</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>4a</sup>,  
 C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>4a</sup>, or  
 C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>4a</sup>;

R<sup>4a</sup>, at each occurrence, is independently selected from  
 30 H, F, CF<sub>3</sub>,  
 C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,  
 phenyl substituted with 0-3 R<sup>4b</sup>, or



- 5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R<sup>4b</sup>; wherein said 5 to 6 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;
- 10 R<sup>4b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;
- 15 R<sup>5</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>5b</sup>; C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>5b</sup>; C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>5b</sup>;
- R<sup>5b</sup>, at each occurrence, is independently selected from:
- 20 H, methyl, ethyl, propyl, butyl, CF<sub>3</sub>, OR<sup>14</sup>, =O; C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-2 R<sup>5c</sup>; phenyl substituted with 0-3 R<sup>5c</sup>; or 5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R<sup>5c</sup>; wherein said 5 to 6 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;
- 25 30
- R<sup>5c</sup>, at each occurrence, is independently selected from H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;
- 35
- R<sup>10</sup> is H, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>; C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with 0-1 R<sup>10a</sup>;

phenyl substituted with 0-4 R<sup>10b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>10b</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
5 substituted with 0-3 R<sup>10b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
10 imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>10a</sup>, at each occurrence, is independently selected from H,  
methyl, ethyl, propyl, butyl, OR<sup>14</sup>, Cl, F, =O, NR<sup>15</sup>R<sup>16</sup>,  
CF<sub>3</sub>, or phenyl substituted with 0-4 R<sup>10b</sup>;

15 R<sup>10b</sup>, at each occurrence, is independently selected from H,  
OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
propoxy, Cl, F, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;

20 Z is H;

C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-3 R<sup>12a</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>12a</sup>; or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>12a</sup>;

25 R<sup>12a</sup>, at each occurrence, is independently selected from  
H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

30 R<sup>13</sup>, at each occurrence, is independently selected from  
H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
Cl, F, Br, CN, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;

R<sup>14</sup> is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;

35

R<sup>15</sup>, at each occurrence, is independently selected from H,  
methyl, ethyl, propyl, and butyl;

R<sup>16</sup>, at each occurrence, is independently selected from  
H, OH, methyl, ethyl, propyl, butyl, benzyl,  
phenethyl, methyl-C(=O)-, ethyl-C(=O)-,  
methyl-S(=O)<sub>2</sub>-, and ethyl-S(=O)<sub>2</sub>-;

5

R<sup>17</sup> is H, methyl, ethyl, propyl, butyl, methoxymethyl,  
ethoxymethyl, methoxyethyl, ethoxyethyl,  
phenyl substituted by 0-3 R<sup>17a</sup>, or  
-CH<sub>2</sub>-phenyl substituted by 0-3 R<sup>17a</sup>;

10


R<sup>17a</sup> is H, methyl, methoxy, -OH, F, Cl, CF<sub>3</sub>, or OCF<sub>3</sub>;

R<sup>18</sup>, at each occurrence, is independently selected from  
H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and  
phenethyl; and

15

R<sup>19</sup>, at each occurrence, is independently selected from  
H, methyl, and ethyl.

20 9. A compound according to Claim 5 of Formula (If):



(If)

or a pharmaceutically acceptable salt or prodrug thereof

25 wherein:

R<sup>3</sup> is R<sup>4</sup>,

R<sup>4</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>4a</sup>,  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>4a</sup>, or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>4a</sup>;

30

R<sup>4a</sup>, at each occurrence, is independently selected from

H, F, CF<sub>3</sub>,  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,  
phenyl substituted with 0-3 R<sup>4b</sup>, or  
5 to 6 membered heterocycle containing 1 to 4  
5 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>4b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
10 pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>4b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
15 S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

R<sup>5</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>5b</sup>;  
20 C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>5b</sup>;

R<sup>5b</sup>, at each occurrence, is independently selected from:  
H, methyl, ethyl, propyl, butyl, CF<sub>3</sub>, OR<sup>14</sup>, =O;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-2 R<sup>5c</sup>;  
25 phenyl substituted with 0-3 R<sup>5c</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>5c</sup>; wherein said 5 to 6  
30 membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

35 R<sup>5c</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

Z is H;

C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-3 R<sup>12a</sup>;

C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>12a</sup>; or

5 C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>12a</sup>;

R<sup>12a</sup>, at each occurrence, is independently selected from

H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,

S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,

10 ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

R<sup>13</sup>, at each occurrence, is independently selected from

H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy,

15 Cl, F, Br, CN, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;

R<sup>14</sup> is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;

R<sup>15</sup>, at each occurrence, is independently selected from H,  
methyl, ethyl, propyl, and butyl;

20

R<sup>16</sup>, at each occurrence, is independently selected from

H, OH, methyl, ethyl, propyl, butyl, benzyl,

phenethyl, methyl-C(=O)-, ethyl-C(=O)-,

methyl-S(=O)<sub>2</sub>-, and ethyl-S(=O)<sub>2</sub>-;

25

R<sup>18</sup>, at each occurrence, is independently selected from

H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and  
phenethyl; and

30 R<sup>19</sup>, at each occurrence, is independently selected from  
H, methyl, and ethyl.

10. A compound, according to one of Claims 6, 7, 8, or 9,  
35 wherein:

R<sup>3</sup> is -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>,

-CH<sub>2</sub>(CH<sub>3</sub>)<sub>2</sub>, -CH(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>C(CH<sub>3</sub>)<sub>3</sub>,

- $-\text{CF}_3$ ,  $-\text{CH}_2\text{CF}_3$ ,  $-\text{CH}_2\text{CH}_2\text{CF}_3$ ,  $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CF}_3$ ,  
 $-\text{CH}=\text{CH}_2$ ,  $-\text{CH}_2\text{CH}=\text{CH}_2$ ,  $-\text{CH}_2\text{C}(\text{CH}_3)=\text{CH}_2$ ,  $-\text{CH}_2\text{CH}=\text{C}(\text{CH}_3)_2$ ,  
 $-\text{CH}_2\text{CH}_2\text{CH}=\text{CH}_2$ ,  $-\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)=\text{CH}_2$ ,  $-\text{CH}_2\text{CH}_2\text{CH}=\text{C}(\text{CH}_3)_2$ ,  
 $\text{cis-CH}_2\text{CH}=\text{CH}(\text{CH}_3)$ ,  $\text{cis-CH}_2\text{CH}_2\text{CH}=\text{CH}(\text{CH}_3)$ ,  
5  $\text{trans-CH}_2\text{CH}=\text{CH}(\text{CH}_3)$ ,  $\text{trans-CH}_2\text{CH}_2\text{CH}=\text{CH}(\text{CH}_3)$ ;  
 $-\text{C}\equiv\text{CH}$ ,  $-\text{CH}_2\text{C}\equiv\text{CH}$ ,  $-\text{CH}_2\text{C}\equiv\text{C}(\text{CH}_3)$ ,  
cyclopropyl- $\text{CH}_2-$ , cyclobutyl- $\text{CH}_2-$ , cyclopentyl- $\text{CH}_2-$ ,  
cyclohexyl- $\text{CH}_2-$ , cyclopropyl- $\text{CH}_2\text{CH}_2-$ ,  
cyclobutyl- $\text{CH}_2\text{CH}_2-$ , cyclopentyl- $\text{CH}_2\text{CH}_2-$ ,  
10 cyclohexyl- $\text{CH}_2\text{CH}_2-$ , phenyl- $\text{CH}_2-$ ,  
(2-F-phenyl) $\text{CH}_2-$ , (3-F-phenyl) $\text{CH}_2-$ , (4-F-phenyl) $\text{CH}_2-$ ,  
(2-Cl-phenyl) $\text{CH}_2-$ , (3-Cl-phenyl) $\text{CH}_2-$ , (4-Cl-phenyl) $\text{CH}_2-$ ,  
(2,3-diF-phenyl) $\text{CH}_2-$ , (2,4-diF-phenyl) $\text{CH}_2-$ ,  
(2,5-diF-phenyl) $\text{CH}_2-$ , (2,6-diF-phenyl) $\text{CH}_2-$ ,  
15 (3,4-diF-phenyl) $\text{CH}_2-$ , (3,5-diF-phenyl) $\text{CH}_2-$ ,  
(2,3-diCl-phenyl) $\text{CH}_2-$ , (2,4-diCl-phenyl) $\text{CH}_2-$ ,  
(2,5-diCl-phenyl) $\text{CH}_2-$ , (2,6-diCl-phenyl) $\text{CH}_2-$ ,  
(3,4-diCl-phenyl) $\text{CH}_2-$ , (3,5-diCl-phenyl) $\text{CH}_2-$ ,  
(3-F-4-Cl-phenyl) $\text{CH}_2-$ , (3-F-5-Cl-phenyl) $\text{CH}_2-$ ,  
20 (3-Cl-4-F-phenyl) $\text{CH}_2-$ , phenyl- $\text{CH}_2\text{CH}_2-$ ,  
(2-F-phenyl) $\text{CH}_2\text{CH}_2-$ , (3-F-phenyl) $\text{CH}_2\text{CH}_2-$ ,  
(4-F-phenyl) $\text{CH}_2\text{CH}_2-$ , (2-Cl-phenyl) $\text{CH}_2\text{CH}_2-$ ,  
(3-Cl-phenyl) $\text{CH}_2\text{CH}_2-$ , (4-Cl-phenyl) $\text{CH}_2\text{CH}_2-$ ,  
(2,3-diF-phenyl) $\text{CH}_2\text{CH}_2-$ , (2,4-diF-phenyl) $\text{CH}_2\text{CH}_2-$ ,  
25 (2,5-diF-phenyl) $\text{CH}_2\text{CH}_2-$ , (2,6-diF-phenyl) $\text{CH}_2\text{CH}_2-$ ,  
(3,4-diF-phenyl) $\text{CH}_2\text{CH}_2-$ , (3,5-diF-phenyl) $\text{CH}_2\text{CH}_2-$ ,  
(2,3-diCl-phenyl) $\text{CH}_2\text{CH}_2-$ , (2,4-diCl-phenyl) $\text{CH}_2\text{CH}_2-$ ,  
(2,5-diCl-phenyl) $\text{CH}_2\text{CH}_2-$ , (2,6-diCl-phenyl) $\text{CH}_2\text{CH}_2-$ ,  
(3,4-diCl-phenyl) $\text{CH}_2\text{CH}_2-$ , (3,5-diCl-phenyl) $\text{CH}_2\text{CH}_2-$ ,  
30 (3-F-4-Cl-phenyl) $\text{CH}_2\text{CH}_2-$ , or (3-F-5-Cl-phenyl) $\text{CH}_2\text{CH}_2-$ ,  
  
 $\text{R}^5$  is  $-\text{CH}_3$ ,  $-\text{CH}_2\text{CH}_3$ ,  $-\text{CH}_2\text{CH}_2\text{CH}_3$ ,  $-\text{CH}(\text{CH}_3)_2$ ,  $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ ,  
 $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$ ,  $-\text{CH}_2\text{CH}(\text{CH}_3)_2$ ,  $-\text{CH}_2\text{C}(\text{CH}_3)_3$ ,  
 $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ ,  $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_3$ ,  $-\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$ ,  
35  $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$ ,  $-\text{CH}(\text{CH}_2\text{CH}_3)_2$ ,  $-\text{CF}_3$ ,  $-\text{CH}_2\text{CF}_3$ ,  $-\text{CH}_2\text{CH}_2\text{CF}_3$ ,  
 $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CF}_3$ ,  $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CF}_3$ ,  $-\text{CH}=\text{CH}_2$ ,  $-\text{CH}_2\text{CH}=\text{CH}_2$ ,  
 $-\text{CH}=\text{CHCH}_3$ ,  $\text{cis-CH}_2\text{CH}=\text{CH}(\text{CH}_3)$ ,  $\text{trans-CH}_2\text{CH}=\text{CH}(\text{CH}_3)$ ,  
 $\text{trans-CH}_2\text{CH}=\text{CH}(\text{C}_6\text{H}_5)$ ,  $-\text{CH}_2\text{CH}=\text{C}(\text{CH}_3)_2$ ,  $\text{cis-CH}_2\text{CH}=\text{CHCH}_2\text{CH}_3$ ,

- trans-CH<sub>2</sub>CH=CHCH<sub>2</sub>CH<sub>3</sub>, cis-CH<sub>2</sub>CH<sub>2</sub>CH=CH(CH<sub>3</sub>),  
 trans-CH<sub>2</sub>CH<sub>2</sub>CH=CH(CH<sub>3</sub>), trans-CH<sub>2</sub>CH=CHCH<sub>2</sub>(C<sub>6</sub>H<sub>5</sub>),  
 -C≡CH, -CH<sub>2</sub>C≡CH, -CH<sub>2</sub>C≡C(CH<sub>3</sub>), -CH<sub>2</sub>C≡C(C<sub>6</sub>H<sub>5</sub>)  
 -CH<sub>2</sub>CH<sub>2</sub>C≡CH, -CH<sub>2</sub>CH<sub>2</sub>C≡C(CH<sub>3</sub>), -CH<sub>2</sub>CH<sub>2</sub>C≡C(C<sub>6</sub>H<sub>5</sub>)  
 5 -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C≡CH, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C≡C(CH<sub>3</sub>), -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C≡C(C<sub>6</sub>H<sub>5</sub>)  
 cyclopropyl-CH<sub>2</sub>-, cyclobutyl-CH<sub>2</sub>-, cyclopentyl-CH<sub>2</sub>-,  
 cyclohexyl-CH<sub>2</sub>-, (2-CH<sub>3</sub>-cyclopropyl)CH<sub>2</sub>-,  
 (3-CH<sub>3</sub>-cyclobutyl)CH<sub>2</sub>-,  
 cyclopropyl-CH<sub>2</sub>CH<sub>2</sub>-, cyclobutyl-CH<sub>2</sub>CH<sub>2</sub>-,  
 10 cyclopentyl-CH<sub>2</sub>CH<sub>2</sub>-, cyclohexyl-CH<sub>2</sub>CH<sub>2</sub>-,  
 (2-CH<sub>3</sub>-cyclopropyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-CH<sub>3</sub>-cyclobutyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 phenyl-CH<sub>2</sub>-, (2-F-phenyl)CH<sub>2</sub>-, (3-F-phenyl)CH<sub>2</sub>-,  
 (4-F-phenyl)CH<sub>2</sub>-, furanyl-CH<sub>2</sub>-, thienyl-CH<sub>2</sub>-,  
 pyridyl-CH<sub>2</sub>-, 1-imidazolyl-CH<sub>2</sub>-, oxazolyl-CH<sub>2</sub>-,  
 15 isoxazolyl-CH<sub>2</sub>-,  
 phenyl-CH<sub>2</sub>CH<sub>2</sub>-, (2-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (4-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, furanyl-CH<sub>2</sub>CH<sub>2</sub>-, thienyl-CH<sub>2</sub>CH<sub>2</sub>-,  
 pyridyl-CH<sub>2</sub>CH<sub>2</sub>-, 1-imidazolyl-CH<sub>2</sub>CH<sub>2</sub>-, oxazolyl-CH<sub>2</sub>CH<sub>2</sub>-,  
 isoxazolyl-CH<sub>2</sub>CH<sub>2</sub>-,

20

Z is methyl, ethyl, i-propyl, n-propyl, n-butyl, i-butyl,  
 s-butyl, t-butyl, or allyl;

- R<sup>10</sup> is H, methyl, ethyl, phenyl, benzyl, phenethyl,  
 25 4-F-phenyl, (4-F-phenyl)CH<sub>2</sub>-, (4-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 4-Cl-phenyl, (4-Cl-phenyl)CH<sub>2</sub>-, (4-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 4-CH<sub>3</sub>-phenyl, (4-CH<sub>3</sub>-phenyl)CH<sub>2</sub>-, (4-CH<sub>3</sub>-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 4-CF<sub>3</sub>-phenyl, (4-CF<sub>3</sub>-phenyl)CH<sub>2</sub>-, or  
 (4-CF<sub>3</sub>-phenyl)CH<sub>2</sub>CH<sub>2</sub>-;

30

- R<sup>11</sup>, at each occurrence, is independently selected from  
 H, =O, methyl, ethyl, phenyl, benzyl, phenethyl,  
 4-F-phenyl, (4-F-phenyl)CH<sub>2</sub>-, (4-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 3-F-phenyl, (3-F-phenyl)CH<sub>2</sub>-, (3-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 35 2-F-phenyl, (2-F-phenyl)CH<sub>2</sub>-, (2-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 4-Cl-phenyl, (4-Cl-phenyl)CH<sub>2</sub>-, (4-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 3-Cl-phenyl, (3-Cl-phenyl)CH<sub>2</sub>-, (3-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 4-CH<sub>3</sub>-phenyl, (4-CH<sub>3</sub>-phenyl)CH<sub>2</sub>-, (4-CH<sub>3</sub>-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,

3-CH<sub>3</sub>-phenyl, (3-CH<sub>3</sub>-phenyl)CH<sub>2</sub>-, (3-CH<sub>3</sub>-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
4-CF<sub>3</sub>-phenyl, (4-CF<sub>3</sub>-phenyl)CH<sub>2</sub>-, (4-CF<sub>3</sub>-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
pyrid-2-yl, pyrid-3-yl, or pyrid-4-yl, and

- 5 R<sup>13</sup>, at each occurrence, is independently selected from  
H, F, Cl, OH, -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -OCH<sub>3</sub>, or -CF<sub>3</sub>.

11. A compound according to Claim 2 selected from:

- 10 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-  
benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-  
butanediamide;

- (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-  
15 benzodiazepin-3-yl]-2-(2-methylpropyl)-3-propyl-  
butanediamide;

- (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-  
1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-  
20 butanediamide;

- (2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-  
1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-  
butanediamide;

- 25 (2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-  
1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-propyl-  
butanediamide;

- 30 (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-  
1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-propyl-  
butanediamide;

- (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-  
35 benzodiazepin-3-yl]-2-methyl-3-allyl-butanediamide;

- (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-  
1,4-benzodiazepin-3-yl]-2-methyl-3-allyl-butanediamide;



(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-methyl-3-propyl-butanediamide;

5 (2R) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-methyl-butanediamide;

(2R,3S) N1-[1,3-dihydro-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

10

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

15

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

20 (2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(2-fluorophenyl)-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

25

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(2-fluorophenyl)-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

30

(2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-(2-fluorophenyl)-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

35

(2S,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-propyl-butanediamide;

5

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(2-fluorophenyl)-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-propyl-butanediamide;

10 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(4-fluorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

15 (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-fluorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

20 (2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-(4-fluorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

25 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(pyrid-2-yl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(N-morpholino)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

30

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(dimethylamino)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

35 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(N-methyl-N-phenylamino)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(N-piperidinyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

- 5 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(N-homopiperidinyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

- 10 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(3-methoxyphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

- 15 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(pyrid-4-yl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

- 20 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-7-methoxy-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(pyrid-3-yl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

- 25 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-(cyclopropylmethyl)-butanediamide;

- 30 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(3-fluorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

- 35 (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(3-fluorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

- (2R, 3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-(3-fluorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;
- 5 (2R, 3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-(3-buten-1-yl)-butanediamide;
- (2R, 3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-
- 10 (cyclopentylethyl)-butanediamide;
- (2R, 3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-
- 15 methylpropyl)-3-(3-buten-1-yl)-butanediamide;
- (2R, 3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-
- 20 methylpropyl)-3-(3-buten-1-yl)-butanediamide;
- (2R, 3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-
- methylpropyl)-3-allyl-butanediamide;
- 25 (2R, 3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-
- methylpropyl)-3-allyl-butanediamide;
- (2R, 3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-(4-
- 30 trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-
- methylpropyl)-3-allyl-butanediamide;
- (2R, 3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-
- 35 trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-
- methylpropyl)-3-n-butyl-butanediamide;

- (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-propyl-butanediamide;
- 5 (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-chlorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-(3-buten-1-yl)-butanediamide;
- (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-chlorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-n-butyl-butanediamide;
- 10 (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-N4-[benzyl]-butanediamide;
- 15 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-methyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;
- 20 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-n-butyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;
- 25 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(2-methylpropyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;
- (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(4-chlorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;
- 30 (2R,3S) N1-[1,3-dihydro-1-ethyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;
- 35

(2R,3S) N1-[1,3-dihydro-1-propyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

5 (2R,3S) N1-[1,3-dihydro-1-(isopropyl)-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;


10 (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3,3-diallyl-butanediamide;

15 (2R,3S) N1-[6,7-dihydro-5-methyl-6-oxo-5H-dibenz[b,d]azepin-7-yl]-2-(2-methylpropyl)-3-allyl-butanediamide; and

(2R,3S) N1-[1,3,4,5-tetrahydro-1,5-dimethyl-2,4-dioxo-2H-1,5-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide.

20

12. A compound, according to Claim 1, of Formula (Ia):



(Ia)

25

or a pharmaceutically acceptable salt or prodrug thereof, wherein:

30 Z is C<sub>1</sub>-C<sub>8</sub> alkyl substituted with 1-3 R<sup>12</sup>;  
 C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 1-3 R<sup>12</sup>;  
 C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 1-3 R<sup>12</sup>;  
 C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
 C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-4 R<sup>12b</sup>; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R<sup>12b</sup>;

5

provided, when R<sup>13</sup> is H,

then Z is C<sub>4</sub>-C<sub>8</sub> alkyl substituted with 1-3 R<sup>12</sup>;

C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 1-3 R<sup>12</sup>; or

C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 1-3 R<sup>12</sup>; and

10

provided, when ring B is a 1,3,4,5-tetrahydro-1-(Z)-5-(R<sup>10</sup>)-6,6,7,7-tetra(R<sup>11</sup>)-2,4-dioxo-2H-1,5-diazepin-3-yl core, and R<sup>13</sup> is H; then

15 R<sup>10</sup> is H, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>,

S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, S(=O)<sub>2</sub>R<sup>17</sup>; or

C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 0-3 R<sup>10a</sup>;

R<sup>10a</sup>, at each occurrence, is independently selected from

20 H, C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>,

NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>.

13. A compound according to Claim 12 of Formula (Ia)

25 wherein:

R<sup>3</sup> is -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-R<sup>4</sup>,

-(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-S-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>,

-(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-O-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>, or

30 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-N(R<sup>7b</sup>)-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>;

n is 0, 1, or 2;

m is 0, 1, or 2;

35

R<sup>3a</sup> is H, OH, methyl, ethyl, propyl, butyl, methoxy,

ethoxy, propoxy, butoxy, allyl, or 3-buten-1-yl;

R<sup>4</sup> is H, OH, OR<sup>14a</sup>,

C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 0-3 R<sup>4a</sup>,

C<sub>2</sub>-C<sub>6</sub> alkenyl substituted with 0-3 R<sup>4a</sup>,

C<sub>2</sub>-C<sub>6</sub> alkynyl substituted with 0-3 R<sup>4a</sup>,

5 C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,

C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>4b</sup>, or

5 to 10 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle

10 is substituted with 0-3 R<sup>4b</sup>;

R<sup>4a</sup>, at each occurrence, is independently selected from is

H, F, Cl, Br, I, CF<sub>3</sub>,

C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,

15 C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>4b</sup>, or

5 to 10 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle

is substituted with 0-3 R<sup>4b</sup>;

20

R<sup>4b</sup>, at each occurrence, is independently selected from H,

OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>,

S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub>

haloalkyl, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy;

25

R<sup>5</sup> is H, OR<sup>14</sup>;

C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 0-3 R<sup>5b</sup>;

C<sub>1</sub>-C<sub>6</sub> alkoxy substituted with 0-3 R<sup>5b</sup>;

C<sub>2</sub>-C<sub>6</sub> alkenyl substituted with 0-3 R<sup>5b</sup>;

30 C<sub>2</sub>-C<sub>6</sub> alkynyl substituted with 0-3 R<sup>5b</sup>;

C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>5c</sup>;

C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>5c</sup>; or

5 to 10 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

35 sulphur, wherein said 5 to 10 membered heterocycle

is substituted with 0-3 R<sup>5c</sup>;

R<sup>5a</sup> is H or C<sub>1</sub>-C<sub>4</sub> alkyl;



- R<sup>5b</sup>, at each occurrence, is independently selected from:  
H, C<sub>1</sub>-C<sub>6</sub> alkyl, CF<sub>3</sub>, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>,  
NR<sup>15</sup>R<sup>16</sup>;
- 5 C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>5c</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>5c</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
10 is substituted with 0-3 R<sup>5c</sup>;
- R<sup>5c</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>,  
S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub>  
15 haloalkyl, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy;
- R<sup>6</sup> is H, methyl, or ethyl;
- R<sup>7</sup>, at each occurrence, is independently selected from  
20 H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, CF<sub>3</sub>, phenyl, and C<sub>1</sub>-C<sub>4</sub>  
alkyl;
- R<sup>7a</sup>, at each occurrence, is independently selected from  
H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, CF<sub>3</sub>, and C<sub>1</sub>-C<sub>4</sub> alkyl;
- 25 R<sup>7b</sup> is independently selected from H, methyl, ethyl,  
propyl, and butyl;
- Ring B is a 7 membered lactam,  
30 wherein the lactam is saturated, partially saturated  
or unsaturated;  
wherein each additional lactam carbon is substituted  
with 0-2 R<sup>11</sup>; and,  
optionally, the lactam contains a heteroatom selected  
35 from -O-, -S-, -S(=O)-, -S(=O)<sub>2</sub>-, -N=, -NH-, and -  
N(R<sup>10</sup>)-;

additionally, two  $R^{11}$  substituents on adjacent atoms may be combined to form a benzo fused radical; wherein said benzo fused radical is substituted with 0-3  $R^{13}$ ;

5 additionally, two  $R^{11}$  substituents on adjacent atoms may be combined to form a 5 to 6 membered heteroaryl fused radical, wherein said 5 to 6 membered heteroaryl fused radical comprises 1 or 2 heteroatoms selected from N, O, and S; wherein said 5 to 6 membered heteroaryl  
10 fused radical is substituted with 0-3  $R^{13}$ ;

additionally, two  $R^{11}$  substituents on the same or adjacent carbon atoms may be combined to form a  $C_3$ - $C_6$  carbocycle substituted with 0-3  $R^{13}$ ;

15

$R^{10}$  is H,  $C(=O)R^{17}$ ,  $C(=O)OR^{17}$ ,  $C(=O)NR^{18}R^{19}$ ,  
 $S(=O)_2NR^{18}R^{19}$ ,  $S(=O)_2R^{17}$ ;

$C_1$ - $C_6$  alkyl optionally substituted with 0-2  $R^{10a}$ ;

$C_6$ - $C_{10}$  aryl substituted with 0-4  $R^{10b}$ ;

20  $C_3$ - $C_{10}$  carbocycle substituted with 0-3  $R^{10b}$ ; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3  $R^{10b}$ ;

25

$R^{10a}$ , at each occurrence, is independently selected from H,  $C_1$ - $C_6$  alkyl,  $OR^{14}$ , Cl, F, Br, I, =O, CN,  $NO_2$ ,  $NR^{15}R^{16}$ ,  $CF_3$ , or phenyl substituted with 0-4  $R^{10b}$ ;

30  $R^{10b}$ , at each occurrence, is independently selected from H, OH,  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_4$  alkoxy, Cl, F, Br, I, CN,  $NO_2$ ,  $NR^{15}R^{16}$ , or  $CF_3$ ;

$R^{11}$ , at each occurrence, is independently selected from  
35 H,  $C_1$ - $C_4$  alkoxy, Cl, F, Br, I, =O, CN,  $NO_2$ ,  $NR^{18}R^{19}$ ,  $C(=O)R^{17}$ ,  $C(=O)OR^{17}$ ,  $C(=O)NR^{18}R^{19}$ ,  $S(=O)_2NR^{18}R^{19}$ ,  $CF_3$ ;  
 $C_1$ - $C_6$  alkyl optionally substituted with 0-3  $R^{11a}$ ;  
 $C_6$ - $C_{10}$  aryl substituted with 0-3  $R^{11b}$ ;

C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>11b</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
5 is substituted with 0-3 R<sup>11b</sup>;

R<sup>11a</sup>, at each occurrence, is independently selected from H,  
C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>,  
CF<sub>3</sub>, or phenyl substituted with 0-3 R<sup>11b</sup>;

10

R<sup>11b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>,  
S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub>  
haloalkyl, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy;

15

Z is C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 1-3 R<sup>12</sup>;

C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 1-3 R<sup>12</sup>;

C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 1-3 R<sup>12</sup>;

C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;

20

C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-4 R<sup>12b</sup>; or

5 to 10 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle

is substituted with 0-3 R<sup>12b</sup>;

25

R<sup>12</sup>, at each occurrence, is independently selected from

C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;

C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-4 R<sup>12b</sup>; or

5 to 10 membered heterocycle containing 1 to 4

30

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle

is substituted with 0-3 R<sup>12b</sup>;

R<sup>12b</sup>, at each occurrence, is independently selected from  
35 H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl,  
SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy,  
C<sub>1</sub>-C<sub>4</sub> haloalkyl, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy;

- $R^{13}$ , at each occurrence, is independently selected from  
H, OH,  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_4$  alkoxy, Cl, F, Br, I, CN,  
 $NO_2$ ,  $NR^{15}R^{16}$ , and  $CF_3$ ;
- 5  $R^{14}$  is H, phenyl, benzyl,  $C_1$ - $C_6$  alkyl, or  $C_2$ - $C_6$  alkoxyalkyl;
- $R^{14a}$  is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;
- 10  $R^{15}$ , at each occurrence, is independently selected from H,  
 $C_1$ - $C_6$  alkyl, benzyl, phenethyl,  $(C_1$ - $C_6$  alkyl)- $C(=O)-$ ,  
and  $(C_1$ - $C_6$  alkyl)- $S(=O)_2-$ ;
- 15  $R^{16}$ , at each occurrence, is independently selected from  
H, OH,  $C_1$ - $C_6$  alkyl, benzyl, phenethyl,  
 $(C_1$ - $C_6$  alkyl)- $C(=O)-$ , and  $(C_1$ - $C_6$  alkyl)- $S(=O)_2-$ ;
- $R^{17}$  is H,  $C_1$ - $C_6$  alkyl,  $C_2$ - $C_6$  alkoxyalkyl,  
aryl substituted by 0-4  $R^{17a}$ , or  
- $CH_2$ -aryl substituted by 0-4  $R^{17a}$ ;
- 20  $R^{17a}$  is H, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
propoxy, butoxy, -OH, F, Cl, Br, I,  $CF_3$ ,  $OCF_3$ ,  $SCH_3$ ,  
 $S(O)CH_3$ ,  $SO_2CH_3$ ,  $-NH_2$ ,  $-N(CH_3)_2$ , or  $C_1$ - $C_4$  haloalkyl;
- 25  $R^{18}$ , at each occurrence, is independently selected from  
H,  $C_1$ - $C_6$  alkyl, phenyl, benzyl, phenethyl,  
 $(C_1$ - $C_6$  alkyl)- $C(=O)-$ , and  $(C_1$ - $C_6$  alkyl)- $S(=O)_2-$ ; and
- 30  $R^{19}$ , at each occurrence, is independently selected from  
H, OH,  $C_1$ - $C_6$  alkyl, phenyl, benzyl, phenethyl,  
 $(C_1$ - $C_6$  alkyl)- $C(=O)-$ , and  $(C_1$ - $C_6$  alkyl)- $S(=O)_2-$ ;
- provided, when  $R^{13}$  is H,  
then Z is  $C_4$ - $C_6$  alkyl substituted with 1-3  $R^{12}$ ;
- 35  $C_2$ - $C_4$  alkenyl substituted with 1-3  $R^{12}$ ; or  
 $C_2$ - $C_4$  alkynyl substituted with 1-3  $R^{12}$ .

WO 00/38618

PCT/US99/30815

14. A compound according to Claim 13 of Formula (Ia)  
wherein:

5 R<sub>3</sub> is  $-(CH_2)_n-R_4$ ,  
n is 0 or 1;

10 R<sub>3a</sub> is H, OH, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, butoxy, allyl, or 3-buten-1-yl;  
R<sub>4</sub> is H, OH, OR<sub>4a</sub>,  
C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-2 R<sub>4a</sub>,  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-2 R<sub>4a</sub>,  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sub>4a</sub>,  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sub>4b</sub>, or  
5-10 membered heterocycle containing  
atoms selected from nitrogen  
wherein said 5 to 6  
with 0-3 R<sub>4b</sub>;

C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>5b</sup>;

R<sup>5a</sup> is H, methyl, ethyl, propyl, or butyl;

- 5 R<sup>5b</sup>, at each occurrence, is independently selected from:  
H, methyl, ethyl, propyl, butyl, CF<sub>3</sub>, OR<sup>14</sup>, Cl, F, Br,  
I, =O;

C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>5c</sup>;

phenyl substituted with 0-3 R<sup>5c</sup>; or

- 10 5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>5c</sup>;

- 15 R<sup>5c</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>,  
S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkoxy, C<sub>1</sub>-C<sub>2</sub>  
haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

- 20 R<sup>6</sup> is H;

R<sup>7</sup>, at each occurrence, is independently selected from H,  
F, CF<sub>3</sub>, methyl, and ethyl;

- 25 Ring B is a 7 membered lactam,  
wherein the lactam is saturated, partially saturated  
or unsaturated;  
wherein each additional lactam carbon is substituted  
with 0-2 R<sup>11</sup>; and,  
30 optionally, the lactam contains a heteroatom selected  
from -N=, -NH-, and -N(R<sup>10</sup>)-;

- additionally, two R<sup>11</sup> substituents on adjacent atoms may be  
combined to form a benzo fused radical; wherein said  
35 benzo fused radical is substituted with 0-2 R<sup>13</sup>;

additionally, two R<sup>11</sup> substituents on adjacent atoms may be  
combined to form a 5 to 6 membered heteroaryl fused

radical, wherein said 5 to 6 membered heteroaryl fused radical comprises 1 or 2 heteroatoms selected from N, O, and S; wherein said 5 to 6 membered heteroaryl fused radical is substituted with 0-2 R<sup>13</sup>;

5

additionally, two R<sup>11</sup> substituents on the same or adjacent carbon atoms may be combined to form a C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-2 R<sup>13</sup>;

- 10 R<sup>10</sup> is H, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>;  
C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with 0-1 R<sup>10a</sup>;  
phenyl substituted with 0-4 R<sup>10b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>10b</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
15 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>10b</sup>;

- R<sup>10a</sup>, at each occurrence, is independently selected from H,  
20 C<sub>1</sub>-C<sub>4</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>,  
CF<sub>3</sub>, or phenyl substituted with 0-4 R<sup>10b</sup>;

- R<sup>10b</sup>, at each occurrence, is independently selected from H,  
OH, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkoxy, Cl, F, Br, I, CN, NO<sub>2</sub>,  
25 NR<sup>15</sup>R<sup>16</sup>, or CF<sub>3</sub>;

- R<sup>11</sup>, at each occurrence, is independently selected from  
H, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, =O, NR<sup>18</sup>R<sup>19</sup>, C(=O)R<sup>17</sup>,  
C(=O)OR<sup>17</sup>, CF<sub>3</sub>;  
30 C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 0-3 R<sup>11a</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>11b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>11b</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
35 sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>11b</sup>;

R<sup>11a</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>4</sub> alkyl, OR<sup>14</sup>, F, =O, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, or phenyl substituted with 0-3 R<sup>11b</sup>;

5 R<sup>11b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

Z is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 1-3 R<sup>12</sup>;  
10 C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 1-3 R<sup>12</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-4 R<sup>12b</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
15 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R<sup>12b</sup>;

R<sup>12</sup>, at each occurrence, is independently selected from  
20 C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-4 R<sup>12b</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
25 is substituted with 0-3 R<sup>12b</sup>;

R<sup>12b</sup>, at each occurrence, is independently selected from  
H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl,  
30 and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

R<sup>13</sup>, at each occurrence, is independently selected from  
H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, CN,  
NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;

35 R<sup>14</sup> is H, phenyl, benzyl, C<sub>1</sub>-C<sub>4</sub> alkyl, or C<sub>2</sub>-C<sub>4</sub> alkoxyalkyl;



R<sup>15</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>4</sub> alkyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>4</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>4</sub> alkyl)-S(=O)<sub>2</sub>-;

5 R<sup>16</sup>, at each occurrence, is independently selected from H, OH, C<sub>1</sub>-C<sub>4</sub> alkyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>4</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>4</sub> alkyl)-S(=O)<sub>2</sub>-;

10 R<sup>17</sup> is H, methyl, ethyl, propyl, butyl, methoxymethyl, ethoxymethyl, methoxyethyl, ethoxyethyl, phenyl substituted by 0-3 R<sup>17a</sup>, or -CH<sub>2</sub>-phenyl substituted by 0-3 R<sup>17a</sup>;

R<sup>17a</sup> is H, methyl, methoxy, -OH, F, Cl, CF<sub>3</sub>, or OCF<sub>3</sub>;

15 R<sup>18</sup>, at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl; and


20 R<sup>19</sup>, at each occurrence, is independently selected from H, methyl, and ethyl;

provided, when R<sup>13</sup> is H,

then Z is butyl substituted with 1-3 R<sup>12</sup>;


25 C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 1-3 R<sup>12</sup>; or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 1-3 R<sup>12</sup>.

15. A compound of Claim 14 of Formula (Ib):




or a pharmaceutically acceptable salt or prodrug thereof  
wherein:

35 Ring B is selected from:



16. A compound according to Claim 15 of Formula (Ic):



(Ic)

or a pharmaceutically acceptable salt or prodrug thereof  
wherein

$R^3$  is  $R^4$ ,

$R^4$  is  $C_1$ - $C_4$  alkyl substituted with 0-1  $R^{4a}$ ,  
 $C_2$ - $C_4$  alkenyl substituted with 0-1  $R^{4a}$ , or  
 $C_2$ - $C_4$  alkynyl substituted with 0-1  $R^{4a}$ ;

- R<sup>4a</sup>, at each occurrence, is independently selected from  
H, F, CF<sub>3</sub>,  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,  
phenyl substituted with 0-3 R<sup>4b</sup>, or  
5 5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>4b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
10 pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;
- R<sup>4b</sup>, at each occurrence, is independently selected from H,  
15 OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;
- R<sup>5</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>5b</sup>;  
20 C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>5b</sup>;
- R<sup>5b</sup>, at each occurrence, is independently selected from:  
H, methyl, ethyl, propyl, butyl, CF<sub>3</sub>, OR<sup>14</sup>, =O;  
25 C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-2 R<sup>5c</sup>;  
phenyl substituted with 0-3 R<sup>5c</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
30 substituted with 0-3 R<sup>5c</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;  
35
- R<sup>5c</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,

S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

5 R<sup>11</sup>, at each occurrence, is independently selected from  
H, =O, NR<sup>18</sup>R<sup>19</sup>, CF<sub>3</sub>;  
C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with 0-1 R<sup>11a</sup>;  
phenyl substituted with 0-3 R<sup>11b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>11b</sup>; or  
10 5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>11b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
15 pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>11a</sup>, at each occurrence, is independently selected from H,  
C<sub>1</sub>-C<sub>4</sub> alkyl, OR<sup>14</sup>, F, Cl, =O, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, or phenyl  
20 substituted with 0-3 R<sup>11b</sup>;

R<sup>11b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, methyl, ethyl, propyl, butyl,  
methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub>  
25 haloalkoxy;

Z is C<sub>1</sub>-C<sub>3</sub> alkyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>3</sub> alkenyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>3</sub> alkynyl substituted with 1-3 R<sup>12</sup>;  
30 C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>12b</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
35 substituted with 0-3 R<sup>12b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,

pyrrolyl, piperaziny, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;


- 5  $R^{12}$ , at each occurrence, is independently selected from  
 $C_6-C_{10}$  aryl substituted with 0-4  $R^{12b}$ ;  
 $C_3-C_6$  carbocycle substituted with 0-3  $R^{12b}$ ; or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
10 substituted with 0-3  $R^{12b}$ ; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperaziny, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;  
15  $R^{12b}$ , at each occurrence, is independently selected from  
H, OH, Cl, F,  $NR^{15}R^{16}$ ,  $CF_3$ , acetyl,  $SCH_3$ ,  $S(=O)CH_3$ ,  
 $S(=O)_2CH_3$ , methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy,  $C_1-C_2$  haloalkyl, and  $C_1-C_2$  haloalkoxy;  
20  $R^{13}$ , at each occurrence, is independently selected from  
H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
Cl, F, Br, CN,  $NR^{15}R^{16}$ , and  $CF_3$ ;  
25  $R^{14}$  is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;  
 $R^{15}$ , at each occurrence, is independently selected from H,  
methyl, ethyl, propyl, and butyl;  
30  $R^{16}$ , at each occurrence, is independently selected from  
H, OH, methyl, ethyl, propyl, butyl, benzyl,  
phenethyl, methyl-C(=O)-, ethyl-C(=O)-,  
methyl-S(=O)<sub>2</sub>-, and ethyl-S(=O)<sub>2</sub>-;  
35  $R^{18}$ , at each occurrence, is independently selected from  
H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and  
phenethyl; and

R<sup>19</sup>, at each occurrence, is independently selected from H, methyl, and ethyl.

provided, when R<sup>13</sup> is H,

- 5 then Z is C<sub>2</sub>-C<sub>3</sub> alkenyl substituted with 1-3 R<sup>12</sup>; or  
C<sub>2</sub>-C<sub>3</sub> alkynyl substituted with 1-3 R<sup>12</sup>.

17. A compound according to Claim 15 of Formula (Id):



10

(Id)

or a pharmaceutically acceptable salt or prodrug thereof  
wherein:

15 R<sup>3</sup> is R<sup>4</sup>,

R<sup>4</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>4a</sup>,  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>4a</sup>, or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>4a</sup>;

20

R<sup>4a</sup>, at each occurrence, is independently selected from  
H, F, CF<sub>3</sub>,

C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,  
phenyl substituted with 0-3 R<sup>4b</sup>, or

25

5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>4b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
30 pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>4b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

5

R<sup>5</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>5b</sup>;

10 R<sup>5b</sup>, at each occurrence, is independently selected from:  
H, methyl, ethyl, propyl, butyl, CF<sub>3</sub>, OR<sup>14</sup>, =O;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-2 R<sup>5c</sup>;  
phenyl substituted with 0-3 R<sup>5c</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
15 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>5c</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
20 pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>5c</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
25 S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

R<sup>11</sup>, at each occurrence, is independently selected from  
H, =O, NR<sup>18</sup>R<sup>19</sup>, CF<sub>3</sub>;  
30 C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with 0-1 R<sup>11a</sup>;  
phenyl substituted with 0-3 R<sup>11b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>11b</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
35 sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>11b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,

pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

5       R<sup>11a</sup>, at each occurrence, is independently selected from H,  
C<sub>1</sub>-C<sub>4</sub> alkyl, OR<sup>14</sup>, F, Cl, =O, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, or phenyl  
substituted with 0-3 R<sup>11b</sup>;

10       R<sup>11b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, methyl, ethyl, propyl, butyl,  
methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub>  
haloalkoxy;

15       Z is C<sub>1</sub>-C<sub>3</sub> alkyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>3</sub> alkenyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>3</sub> alkynyl substituted with 1-3 R<sup>12</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>12b</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
20       sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>12b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
25       imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

30       R<sup>12</sup>, at each occurrence, is independently selected from  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>12b</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>12b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
35       pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;



R<sup>12b</sup>, at each occurrence, is independently selected from  
H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

5

R<sup>13</sup>, at each occurrence, is independently selected from  
H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
Cl, F, Br, CN, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;

10 R<sup>14</sup> is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;

R<sup>15</sup>, at each occurrence, is independently selected from H,  
methyl, ethyl, propyl, and butyl;

15 R<sup>16</sup>, at each occurrence, is independently selected from  
H, OH, methyl, ethyl, propyl, butyl, benzyl,  
phenethyl, methyl-C(=O)-, ethyl-C(=O)-,  
methyl-S(=O)<sub>2</sub>-, and ethyl-S(=O)<sub>2</sub>-;


20 R<sup>18</sup>, at each occurrence, is independently selected from  
H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and  
phenethyl; and

25 R<sup>19</sup>, at each occurrence, is independently selected from  
H, methyl, and ethyl.

provided, when R<sup>13</sup> is H,  
then Z is C<sub>2</sub>-C<sub>3</sub> alkenyl substituted with 1-3 R<sup>12</sup>; or  
C<sub>2</sub>-C<sub>3</sub> alkynyl substituted with 1-3 R<sup>12</sup>.

30

18. A compound according to Claim 15 of Formula (Ie):



(Ie)

or a pharmaceutically acceptable salt or prodrug thereof  
wherein:

5

$R^3$  is  $R^4$ ,

$R^4$  is  $C_1$ - $C_4$  alkyl substituted with 0-1  $R^{4a}$ ,  
 $C_2$ - $C_4$  alkenyl substituted with 0-1  $R^{4a}$ , or  
10  $C_2$ - $C_4$  alkynyl substituted with 0-1  $R^{4a}$ ;

$R^{4a}$ , at each occurrence, is independently selected from  
H, F,  $CF_3$ ,

15  $C_3$ - $C_6$  carbocycle substituted with 0-3  $R^{4b}$ ,  
phenyl substituted with 0-3  $R^{4b}$ , or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3  $R^{4b}$ ; wherein said 5 to 6  
20 membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

25  $R^{4b}$ , at each occurrence, is independently selected from H,  
OH, Cl, F,  $NR^{15}R^{16}$ ,  $CF_3$ , acetyl,  $SCH_3$ ,  $S(=O)CH_3$ ,  
 $S(=O)_2CH_3$ , methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy,  $C_1$ - $C_2$  haloalkyl, and  $C_1$ - $C_2$  haloalkoxy;

30  $R^5$  is  $C_1$ - $C_4$  alkyl substituted with 0-1  $R^{5b}$ ;  
 $C_2$ - $C_4$  alkenyl substituted with 0-1  $R^{5b}$ ;  
 $C_2$ - $C_4$  alkynyl substituted with 0-1  $R^{5b}$ ;

R<sup>5b</sup>, at each occurrence, is independently selected from:

H, methyl, ethyl, propyl, butyl, CF<sub>3</sub>, OR<sup>14</sup>, =O;

C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-2 R<sup>5c</sup>;

phenyl substituted with 0-3 R<sup>5c</sup>; or

- 5 5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>5c</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
10 pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>5c</sup>, at each occurrence, is independently selected from H,

- 15 OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

R<sup>10</sup> is H, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>;

- 20 C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with 0-1 R<sup>10a</sup>;  
phenyl substituted with 0-4 R<sup>10b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>10b</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
25 sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>10b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
30 imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>10a</sup>, at each occurrence, is independently selected from H,  
methyl, ethyl, propyl, butyl, OR<sup>14</sup>, Cl, F, =O, NR<sup>15</sup>R<sup>16</sup>,  
CF<sub>3</sub>, or phenyl substituted with 0-4 R<sup>10b</sup>;

35

R<sup>10b</sup>, at each occurrence, is independently selected from H,  
OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
propoxy, Cl, F, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;

Z is C<sub>1</sub>-C<sub>3</sub> alkyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>3</sub> alkenyl, substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>3</sub> alkynyl substituted with 1-3 R<sup>12</sup>;  
5 C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>12b</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
10 substituted with 0-3 R<sup>12b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

15 R<sup>12</sup>, at each occurrence, is independently selected from  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>12b</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
20 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>12b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
25 pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>12b</sup>, at each occurrence, is independently selected from  
H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
30 S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

R<sup>13</sup>, at each occurrence, is independently selected from  
H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
35 Cl, F, Br, CN, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;

R<sup>14</sup> is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;

R<sup>15</sup>, at each occurrence, is independently selected from H, methyl, ethyl, propyl, and butyl;

R<sup>16</sup>, at each occurrence, is independently selected from  
 5 H, OH, methyl, ethyl, propyl, butyl, benzyl, phenethyl, methyl-C(=O)-, ethyl-C(=O)-, methyl-S(=O)<sub>2</sub>-, and ethyl-S(=O)<sub>2</sub>-;

R<sup>17</sup> is H, methyl, ethyl, propyl, butyl, methoxymethyl, ethoxymethyl, methoxyethyl, ethoxyethyl, phenyl substituted by 0-3 R<sup>17a</sup>, or  
 10 -CH<sub>2</sub>-phenyl substituted by 0-3 R<sup>17a</sup>;

R<sup>17a</sup> is H, methyl, methoxy, -OH, F, Cl, CF<sub>3</sub>, or OCF<sub>3</sub>;  
 15

R<sup>18</sup>, at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl; and


20 R<sup>19</sup>, at each occurrence, is independently selected from H, methyl, and ethyl.

provided, when R<sup>13</sup> is H,

then Z is C<sub>2</sub>-C<sub>3</sub> alkenyl substituted with 1-3 R<sup>12</sup>; or

25 C<sub>2</sub>-C<sub>3</sub> alkynyl substituted with 1-3 R<sup>12</sup>.

19. A compound according to Claim 15 of Formula (If):



30

(If)

or a pharmaceutically acceptable salt or prodrug thereof  
 wherein:

R<sup>3</sup> is R<sup>4</sup>,

R<sup>4</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>4a</sup>,  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>4a</sup>, or  
5 C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>4a</sup>;

R<sup>4a</sup>, at each occurrence, is independently selected from  
H, F, CF<sub>3</sub>,  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,  
10 phenyl substituted with 0-3 R<sup>4b</sup>, or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>4b</sup>; wherein said 5 to 6  
15 membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

20 R<sup>4b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

25 R<sup>5</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>5b</sup>;

R<sup>5b</sup>, at each occurrence, is independently selected from:  
30 H, methyl, ethyl, propyl, butyl, CF<sub>3</sub>, OR<sup>14</sup>, =O;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-2 R<sup>5c</sup>;  
phenyl substituted with 0-3 R<sup>5c</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
35 sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>5c</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,

pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

5  $R^{5c}$ , at each occurrence, is independently selected from H,  
OH, Cl, F,  $NR^{15}R^{16}$ ,  $CF_3$ , acetyl,  $SCH_3$ ,  $S(=O)CH_3$ ,  
 $S(=O)_2CH_3$ , methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy,  $C_1$ - $C_2$  haloalkyl, and  $C_1$ - $C_2$  haloalkoxy;

10 Z is  $C_1$ - $C_3$  alkyl substituted with 1-3  $R^{12}$ ;  
 $C_2$ - $C_3$  alkenyl substituted with 1-3  $R^{12}$ ;  
 $C_2$ - $C_3$  alkynyl substituted with 1-3  $R^{12}$ ;  
 $C_6$ - $C_{10}$  aryl substituted with 0-4  $R^{12b}$ ;  
 $C_3$ - $C_6$  carbocycle substituted with 0-3  $R^{12b}$ ; or  
5 to 6 membered heterocycle containing 1 to 4  
15 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3  $R^{12b}$ ; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
20 pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

$R^{12}$ , at each occurrence, is independently selected from  
 $C_6$ - $C_{10}$  aryl substituted with 0-4  $R^{12b}$ ;  
25  $C_3$ - $C_6$  carbocycle substituted with 0-3  $R^{12b}$ ; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3  $R^{12b}$ ; wherein said 5 to 6  
30 membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

35  $R^{12b}$ , at each occurrence, is independently selected from  
H, OH, Cl, F,  $NR^{15}R^{16}$ ,  $CF_3$ , acetyl,  $SCH_3$ ,  $S(=O)CH_3$ ,  
 $S(=O)_2CH_3$ , methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy,  $C_1$ - $C_2$  haloalkyl, and  $C_1$ - $C_2$  haloalkoxy;

R<sup>13</sup>, at each occurrence, is independently selected from  
H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
Cl, F, Br, CN, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;

5

R<sup>14</sup> is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;

R<sup>15</sup>, at each occurrence, is independently selected from H,  
methyl, ethyl, propyl, and butyl;

10

R<sup>16</sup>, at each occurrence, is independently selected from  
H, OH, methyl, ethyl, propyl, butyl, benzyl,  
phenethyl, methyl-C(=O)-, ethyl-C(=O)-,  
methyl-S(=O)<sub>2</sub>-, and ethyl-S(=O)<sub>2</sub>-;

15

R<sup>18</sup>, at each occurrence, is independently selected from  
H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and  
phenethyl; and

20 R<sup>19</sup>, at each occurrence, is independently selected from  
H, methyl, and ethyl.

provided, when R<sup>13</sup> is H,

then Z is C<sub>2</sub>-C<sub>3</sub> alkenyl substituted with 1-3 R<sup>12</sup>; or

25

C<sub>2</sub>-C<sub>3</sub> alkynyl substituted with 1-3 R<sup>12</sup>.

20. A compound according to one of Claims 16, 17, 18, 19,  
wherein:

30 R<sup>3</sup> is -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>,  
-CH<sub>2</sub>(CH<sub>3</sub>)<sub>2</sub>, -CH(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>C(CH<sub>3</sub>)<sub>3</sub>,  
-CF<sub>3</sub>, -CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CF<sub>3</sub>,  
-CH=CH<sub>2</sub>, -CH<sub>2</sub>CH=CH<sub>2</sub>, -CH<sub>2</sub>C(CH<sub>3</sub>)=CH<sub>2</sub>, -CH<sub>2</sub>CH=C(CH<sub>3</sub>)<sub>2</sub>,  
-CH<sub>2</sub>CH<sub>2</sub>CH=CH<sub>2</sub>, -CH<sub>2</sub>CH<sub>2</sub>C(CH<sub>3</sub>)=CH<sub>2</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH=C(CH<sub>3</sub>)<sub>2</sub>,  
35 cis-CH<sub>2</sub>CH=CH(CH<sub>3</sub>), cis-CH<sub>2</sub>CH<sub>2</sub>CH=CH(CH<sub>3</sub>),  
trans-CH<sub>2</sub>CH=CH(CH<sub>3</sub>), trans-CH<sub>2</sub>CH<sub>2</sub>CH=CH(CH<sub>3</sub>);  
-C≡CH, -CH<sub>2</sub>C≡CH, -CH<sub>2</sub>C≡C(CH<sub>3</sub>),



- cyclopropyl-CH<sub>2</sub>-, cyclobutyl-CH<sub>2</sub>-, cyclopentyl-CH<sub>2</sub>-,  
 cyclohexyl-CH<sub>2</sub>-, cyclopropyl-CH<sub>2</sub>CH<sub>2</sub>-,  
 cyclobutyl-CH<sub>2</sub>CH<sub>2</sub>-, cyclopentyl-CH<sub>2</sub>CH<sub>2</sub>-,  
 cyclohexyl-CH<sub>2</sub>CH<sub>2</sub>-, phenyl-CH<sub>2</sub>-,
- 5 (2-F-phenyl)CH<sub>2</sub>-, (3-F-phenyl)CH<sub>2</sub>-, (4-F-phenyl)CH<sub>2</sub>-,  
 (2-Cl-phenyl)CH<sub>2</sub>-, (3-Cl-phenyl)CH<sub>2</sub>-, (4-Cl-phenyl)CH<sub>2</sub>-,  
 (2,3-diF-phenyl)CH<sub>2</sub>-, (2,4-diF-phenyl)CH<sub>2</sub>-,  
 (2,5-diF-phenyl)CH<sub>2</sub>-, (2,6-diF-phenyl)CH<sub>2</sub>-,  
 (3,4-diF-phenyl)CH<sub>2</sub>-, (3,5-diF-phenyl)CH<sub>2</sub>-,
- 10 (2,3-diCl-phenyl)CH<sub>2</sub>-, (2,4-diCl-phenyl)CH<sub>2</sub>-,  
 (2,5-diCl-phenyl)CH<sub>2</sub>-, (2,6-diCl-phenyl)CH<sub>2</sub>-,  
 (3,4-diCl-phenyl)CH<sub>2</sub>-, (3,5-diCl-phenyl)CH<sub>2</sub>-,  
 (3-F-4-Cl-phenyl)CH<sub>2</sub>-, (3-F-5-Cl-phenyl)CH<sub>2</sub>-,  
 (3-Cl-4-F-phenyl)CH<sub>2</sub>-, phenyl-CH<sub>2</sub>CH<sub>2</sub>-,
- 15 (2-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (4-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (3-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (4-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2,3-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2,4-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2,5-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2,6-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,
- 20 (3,4-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3,5-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2,3-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2,4-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2,5-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2,6-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (3,4-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3,5-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (3-F-4-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, or (3-F-5-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,
- 25
- R<sup>5</sup> is -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>,  
 -CH(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>C(CH<sub>3</sub>)<sub>3</sub>,  
 -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -CH(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>3</sub>,  
 -CH<sub>2</sub>CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>, -CF<sub>3</sub>, -CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CF<sub>3</sub>,  
 30 -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CF<sub>3</sub>, -CH=CH<sub>2</sub>, -CH<sub>2</sub>CH=CH<sub>2</sub>,  
 -CH=CHCH<sub>3</sub>, cis-CH<sub>2</sub>CH=CH(CH<sub>3</sub>), trans-CH<sub>2</sub>CH=CH(CH<sub>3</sub>),  
 trans-CH<sub>2</sub>CH=CH(C<sub>6</sub>H<sub>5</sub>), -CH<sub>2</sub>CH=C(CH<sub>3</sub>)<sub>2</sub>, cis-CH<sub>2</sub>CH=CHCH<sub>2</sub>CH<sub>3</sub>,  
 trans-CH<sub>2</sub>CH=CHCH<sub>2</sub>CH<sub>3</sub>, cis-CH<sub>2</sub>CH<sub>2</sub>CH=CH(CH<sub>3</sub>),  
 trans-CH<sub>2</sub>CH<sub>2</sub>CH=CH(CH<sub>3</sub>), trans-CH<sub>2</sub>CH=CHCH<sub>2</sub>(C<sub>6</sub>H<sub>5</sub>),
- 35 -C≡CH, -CH<sub>2</sub>C≡CH, -CH<sub>2</sub>C≡C(CH<sub>3</sub>), -CH<sub>2</sub>C≡C(C<sub>6</sub>H<sub>5</sub>)  
 -CH<sub>2</sub>CH<sub>2</sub>C≡CH, -CH<sub>2</sub>CH<sub>2</sub>C≡C(CH<sub>3</sub>), -CH<sub>2</sub>CH<sub>2</sub>C≡C(C<sub>6</sub>H<sub>5</sub>)  
 -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C≡CH, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C≡C(CH<sub>3</sub>), -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C≡C(C<sub>6</sub>H<sub>5</sub>)  
 cyclopropyl-CH<sub>2</sub>-, cyclobutyl-CH<sub>2</sub>-, cyclopentyl-CH<sub>2</sub>-,

- cyclohexyl-CH<sub>2</sub>-, (2-CH<sub>3</sub>-cyclopropyl)CH<sub>2</sub>-,  
 (3-CH<sub>3</sub>-cyclobutyl)CH<sub>2</sub>-,  
 cyclopropyl-CH<sub>2</sub>CH<sub>2</sub>-, cyclobutyl-CH<sub>2</sub>CH<sub>2</sub>-,  
 cyclopentyl-CH<sub>2</sub>CH<sub>2</sub>-, cyclohexyl-CH<sub>2</sub>CH<sub>2</sub>-,  
 5 (2-CH<sub>3</sub>-cyclopropyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-CH<sub>3</sub>-cyclobutyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 phenyl-CH<sub>2</sub>-, (2-F-phenyl)CH<sub>2</sub>-, (3-F-phenyl)CH<sub>2</sub>-,  
 (4-F-phenyl)CH<sub>2</sub>-, furanyl-CH<sub>2</sub>-, thienyl-CH<sub>2</sub>-,  
 pyridyl-CH<sub>2</sub>-, 1-imidazolyl-CH<sub>2</sub>-, oxazolyl-CH<sub>2</sub>-,  
 isoxazolyl-CH<sub>2</sub>-,  
 10 phenyl-CH<sub>2</sub>CH<sub>2</sub>-, (2-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (4-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, furanyl-CH<sub>2</sub>CH<sub>2</sub>-, thienyl-CH<sub>2</sub>CH<sub>2</sub>-,  
 pyridyl-CH<sub>2</sub>CH<sub>2</sub>-, 1-imidazolyl-CH<sub>2</sub>CH<sub>2</sub>-, oxazolyl-CH<sub>2</sub>CH<sub>2</sub>-,  
 isoxazolyl-CH<sub>2</sub>CH<sub>2</sub>-,  
 15 Z is phenyl, 2-F-phenyl, 3-F-phenyl, 4-F-phenyl,  
 2-Cl-phenyl, 3-Cl-phenyl, 4-Cl-phenyl, 2,3-diF-phenyl,  
 2,4-diF-phenyl, 2,5-diF-phenyl, 2,6-diF-phenyl,  
 3,4-diF-phenyl, 3,5-diF-phenyl, 2,3-diCl-phenyl,  
 2,4-diCl-phenyl, 2,5-diCl-phenyl, 2,6-diCl-phenyl,  
 20 3,4-diCl-phenyl, 3,5-diCl-phenyl, 3-F-4-Cl-phenyl,  
 3-F-5-Cl-phenyl, 3-Cl-4-F-phenyl, 2-MeO-phenyl,  
 3-MeO-phenyl, 4-MeO-phenyl, 2-Me-phenyl, 3-Me-phenyl,  
 4-Me-phenyl, 2-MeS-phenyl, 3-MeS-phenyl, 4-MeS-phenyl,  
 2-CF<sub>3</sub>O-phenyl, 3-CF<sub>3</sub>O-phenyl, 4-CF<sub>3</sub>O-phenyl,  
 25 furanyl, thienyl, pyridyl, 2-Me-pyridyl, 3-Me-pyridyl,  
 4-Me-pyridyl, 1-imidazolyl, oxazolyl, isoxazolyl,  
 1-benzimidazolyl,  
 cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl,  
 morpholino, N-piperinyl,  
 30 phenyl-CH<sub>2</sub>-, (2-F-phenyl)CH<sub>2</sub>-, (3-F-phenyl)CH<sub>2</sub>-,  
 (4-F-phenyl)CH<sub>2</sub>-, (2-Cl-phenyl)CH<sub>2</sub>-, (3-Cl-phenyl)CH<sub>2</sub>-,  
 (4-Cl-phenyl)CH<sub>2</sub>-, (2,3-diF-phenyl)CH<sub>2</sub>-,  
 (2,4-diF-phenyl)CH<sub>2</sub>-, (2,5-diF-phenyl)CH<sub>2</sub>-,  
 (2,6-diF-phenyl)CH<sub>2</sub>-, (3,4-diF-phenyl)CH<sub>2</sub>-,  
 35 (3,5-diF-phenyl)CH<sub>2</sub>-, (2,3-diCl-phenyl)CH<sub>2</sub>-,  
 (2,4-diCl-phenyl)CH<sub>2</sub>-, (2,5-diCl-phenyl)CH<sub>2</sub>-,  
 (2,6-diCl-phenyl)CH<sub>2</sub>-, (3,4-diCl-phenyl)CH<sub>2</sub>-,  
 (3,5-diCl-phenyl)CH<sub>2</sub>-, (3-F-4-Cl-phenyl)CH<sub>2</sub>-,

- (3-F-5-Cl-phenyl)CH<sub>2</sub>-, (3-Cl-4-F-phenyl)CH<sub>2</sub>-,  
 (2-MeO-phenyl)CH<sub>2</sub>-, (3-MeO-phenyl)CH<sub>2</sub>-,  
 (4-MeO-phenyl)CH<sub>2</sub>-, (2-Me-phenyl)CH<sub>2</sub>-,  
 (3-Me-phenyl)CH<sub>2</sub>-, (4-Me-phenyl)CH<sub>2</sub>-,  
 5 (2-MeS-phenyl)CH<sub>2</sub>-, (3-MeS-phenyl)CH<sub>2</sub>-,  
 4-MeS-phenyl)CH<sub>2</sub>-, (2-CF<sub>3</sub>O-phenyl)CH<sub>2</sub>-,  
 (3-CF<sub>3</sub>O-phenyl)CH<sub>2</sub>-, (4-CF<sub>3</sub>O-phenyl)CH<sub>2</sub>-,  
 (furanyl)CH<sub>2</sub>-, (thienyl)CH<sub>2</sub>-, (pyridyl)CH<sub>2</sub>-,  
 (2-Me-pyridyl)CH<sub>2</sub>-, (3-Me-pyridyl)CH<sub>2</sub>-,  
 10 (4-Me-pyridyl)CH<sub>2</sub>-, (1-imidazolyl)CH<sub>2</sub>-,  
 (oxazolyl)CH<sub>2</sub>-, (isoxazolyl)CH<sub>2</sub>-,  
 (1-benzimidazolyl)CH<sub>2</sub>-, (cyclopropyl)CH<sub>2</sub>-,  
 (cyclobutyl)CH<sub>2</sub>-, (cyclopentyl)CH<sub>2</sub>-,  
 (cyclohexyl)CH<sub>2</sub>-, (morpholino)CH<sub>2</sub>-, (N-pipridinyl)CH<sub>2</sub>-,  
 15 phenyl-CH<sub>2</sub>CH<sub>2</sub>-, (phenyl)<sub>2</sub>CHCH<sub>2</sub>-, (2-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (3-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (4-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (4-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2,3-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 20 (2,4-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2,5-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2,6-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3,4-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (3,5-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2,3-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2,4-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2,5-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2,6-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3,4-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 25 (3,5-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-F-4-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (3-F-5-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-Cl-4-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2-MeO-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-MeO-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (4-MeO-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2-Me-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (3-Me-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (4-Me-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 30 (2-MeS-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-MeS-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (4-MeS-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2-CF<sub>3</sub>O-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (3-CF<sub>3</sub>O-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (4-CF<sub>3</sub>O-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (furanyl)CH<sub>2</sub>CH<sub>2</sub>-, (thienyl)CH<sub>2</sub>CH<sub>2</sub>-, (pyridyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2-Me-pyridyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-Me-pyridyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 35 (4-Me-pyridyl)CH<sub>2</sub>CH<sub>2</sub>-, (imidazolyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (oxazolyl)CH<sub>2</sub>CH<sub>2</sub>-, (isoxazolyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (benzimidazolyl)CH<sub>2</sub>CH<sub>2</sub>-, (cyclopropyl)CH<sub>2</sub>CH<sub>2</sub>-,


(cyclobutyl)CH<sub>2</sub>CH<sub>2</sub>-, (cyclopentyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (cyclohexyl)CH<sub>2</sub>CH<sub>2</sub>-, (morpholino)CH<sub>2</sub>CH<sub>2</sub>-, or  
 (N-piperidiny)CH<sub>2</sub>CH<sub>2</sub>-;

- 5 R<sup>10</sup> is H, methyl, ethyl, phenyl, benzyl, phenethyl,  
 4-F-phenyl, (4-F-phenyl)CH<sub>2</sub>-, (4-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 4-Cl-phenyl, (4-Cl-phenyl)CH<sub>2</sub>-, (4-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 4-CH<sub>3</sub>-phenyl, (4-CH<sub>3</sub>-phenyl)CH<sub>2</sub>-, (4-CH<sub>3</sub>-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 4-CF<sub>3</sub>-phenyl, (4-CF<sub>3</sub>-phenyl)CH<sub>2</sub>-, or  
 10 (4-CF<sub>3</sub>-phenyl)CH<sub>2</sub>CH<sub>2</sub>-;

- R<sup>11</sup>, at each occurrence, is independently selected from  
 H, =O, methyl, ethyl, phenyl, benzyl, phenethyl,  
 4-F-phenyl, (4-F-phenyl)CH<sub>2</sub>-, (4-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 15 3-F-phenyl, (3-F-phenyl)CH<sub>2</sub>-, (3-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 2-F-phenyl, (2-F-phenyl)CH<sub>2</sub>-, (2-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 4-Cl-phenyl, (4-Cl-phenyl)CH<sub>2</sub>-, (4-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 3-Cl-phenyl, (3-Cl-phenyl)CH<sub>2</sub>-, (3-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 4-CH<sub>3</sub>-phenyl, (4-CH<sub>3</sub>-phenyl)CH<sub>2</sub>-, (4-CH<sub>3</sub>-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 20 3-CH<sub>3</sub>-phenyl, (3-CH<sub>3</sub>-phenyl)CH<sub>2</sub>-, (3-CH<sub>3</sub>-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 4-CF<sub>3</sub>-phenyl, (4-CF<sub>3</sub>-phenyl)CH<sub>2</sub>-, (4-CF<sub>3</sub>-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 pyrid-2-yl, pyrid-3-yl, or pyrid-4-yl, and

- R<sup>13</sup>, at each occurrence, is independently selected from  
 25 H, F, Cl, OH, -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -OCH<sub>3</sub>, or -CF<sub>3</sub>.

21. A method for the treatment of neurological disorders  
 associated with  $\beta$ -amyloid production comprising  
 administering to a host in need of such treatment a  
 30 therapeutically effective amount of a compound of Formula  
 (I):



or a pharmaceutically acceptable salt or prodrug thereof,  
wherein:

Q is  $-NR^1R^2$ ;

5

$R^1$  is  $OR^{14}$ ;

$R^2$  is independently selected from H,  $C_1$ - $C_6$  alkyl,  $C_3$ - $C_{10}$   
carbocycle,  $C_6$ - $C_{10}$  aryl, and 5 to 10 membered  
10 heterocycle containing 1 to 4 heteroatoms selected  
from nitrogen, oxygen, and sulphur;

$R^3$  is  $-(CR^7R^{7a})_n-R^4$ ,

$-(CR^7R^{7a})_n-S-(CR^7R^{7a})_m-R^4$ ,

15  $-(CR^7R^{7a})_n-O-(CR^7R^{7a})_m-R^4$ ,

$-(CR^7R^{7a})_n-N(R^{7b})-(CR^7R^{7a})_m-R^4$ ,

$-(CR^7R^{7a})_n-S(=O)-(CR^7R^{7a})_m-R^4$ ,

$-(CR^7R^{7a})_n-S(=O)_2-(CR^7R^{7a})_m-R^4$ ,

$-(CR^7R^{7a})_n-C(=O)-(CR^7R^{7a})_m-R^4$ ,

20  $-(CR^7R^{7a})_n-N(R^{7b})C(=O)-(CR^7R^{7a})_m-R^4$ ,

$-(CR^7R^{7a})_n-C(=O)N(R^{7b})-(CR^7R^{7a})_m-R^4$ ,

$-(CR^7R^{7a})_n-N(R^{7b})S(=O)_2-(CR^7R^{7a})_m-R^4$ , or

$-(CR^7R^{7a})_n-S(=O)_2N(R^{7b})-(CR^7R^{7a})_m-R^4$ ;

25  $n$  is 0, 1, 2, or 3;

$m$  is 0, 1, 2, or 3;

$R^{3a}$  is H, OH,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  alkoxy,  $C_2$ - $C_4$  alkenyl  
30 or  $C_2$ - $C_4$  alkenyloxy;

$R^4$  is H, OH,  $OR^{14a}$ ,

$C_1$ - $C_6$  alkyl substituted with 0-3  $R^{4a}$ ,

$C_2$ - $C_6$  alkenyl substituted with 0-3  $R^{4a}$ ,

35  $C_2$ - $C_6$  alkynyl substituted with 0-3  $R^{4a}$ ,

$C_3$ - $C_{10}$  carbocycle substituted with 0-3  $R^{4b}$ ,

$C_6$ - $C_{10}$  aryl substituted with 0-3  $R^{4b}$ , or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R<sup>4b</sup>;

5

R<sup>4a</sup>, at each occurrence, is independently selected from is H, F, Cl, Br, I, CF<sub>3</sub>,

C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,

C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>4b</sup>, or

10

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R<sup>4b</sup>;

15

R<sup>4b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>,

C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl,

C<sub>1</sub>-C<sub>4</sub> haloalkoxy, and C<sub>1</sub>-C<sub>4</sub> halothioalkyl-S-;

20

R<sup>5</sup> is H, OR<sup>14</sup>;

C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 0-3 R<sup>5b</sup>;

C<sub>1</sub>-C<sub>6</sub> alkoxy substituted with 0-3 R<sup>5b</sup>;

C<sub>2</sub>-C<sub>6</sub> alkenyl substituted with 0-3 R<sup>5b</sup>;

25

C<sub>2</sub>-C<sub>6</sub> alkynyl substituted with 0-3 R<sup>5b</sup>;

C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>5c</sup>;

C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>5c</sup>; or

5 to 10 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

30

sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R<sup>5c</sup>;

R<sup>5a</sup> is H, OH, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>2</sub>-C<sub>4</sub> alkenyl, or C<sub>2</sub>-C<sub>4</sub> alkenyloxy;

35

R<sup>5b</sup>, at each occurrence, is independently selected from:

H, C<sub>1</sub>-C<sub>6</sub> alkyl, CF<sub>3</sub>, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>,

NR<sup>15</sup>R<sup>16</sup>;

C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>5c</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>5c</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
5 sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>5c</sup>;

R<sup>5c</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>,  
10 S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>,  
C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl,  
C<sub>1</sub>-C<sub>4</sub> haloalkoxy, and C<sub>1</sub>-C<sub>4</sub> halothioalkyl-S-;

R<sup>6</sup> is H;  
15 C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 0-3 R<sup>6a</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>6b</sup>; or  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>6b</sup>;

R<sup>6a</sup>, at each occurrence, is independently selected from H,  
20 C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>,  
aryl or CF<sub>3</sub>;

R<sup>6b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl,  
25 C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy;

R<sup>7</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, CF<sub>3</sub>, phenyl and C<sub>1</sub>-C<sub>4</sub> alkyl;

30 R<sup>7a</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, CF<sub>3</sub>, and C<sub>1</sub>-C<sub>4</sub> alkyl;

R<sup>7b</sup> is independently selected from H and C<sub>1</sub>-C<sub>4</sub> alkyl;

35 Ring B is a 7 membered lactam,  
wherein the lactam is saturated, partially saturated  
or unsaturated;

wherein each additional lactam carbon is substituted  
with 0-2 R<sup>11</sup>; and,  
optionally, the lactam contains a heteroatom selected  
from -O-, -S-, -S(=O)-, -S(=O)<sub>2</sub>-, -N=, -NH-, and -  
5 N(R<sup>10</sup>)-;

additionally, two R<sup>11</sup> substituents on adjacent atoms may be  
combined to form a benzo fused radical; wherein said  
benzo fused radical is substituted with 0-4 R<sup>13</sup>;

10 additionally, two R<sup>11</sup> substituents on adjacent atoms may be  
combined to form a 5 to 6 membered heteroaryl fused  
radical, wherein said 5 to 6 membered heteroaryl fused  
radical comprises 1 or 2 heteroatoms selected from N,  
15 O, and S; wherein said 5 to 6 membered heteroaryl  
fused radical is substituted with 0-3 R<sup>13</sup>;

additionally, two R<sup>11</sup> substituents on the same or adjacent  
carbon atoms may be combined to form a C<sub>3</sub>-C<sub>6</sub>  
20 carbocycle substituted with 0-3 R<sup>13</sup>;

R<sup>10</sup> is H, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>,  
S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, S(=O)<sub>2</sub>R<sup>17</sup>;  
C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 0-3 R<sup>10a</sup>;  
25 C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>10b</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>10b</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
30 is substituted with 0-3 R<sup>10b</sup>;

R<sup>10a</sup>, at each occurrence, is independently selected from H,  
C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>,  
CF<sub>3</sub>, or aryl substituted with 0-4 R<sup>10b</sup>;

35 R<sup>10b</sup>, at each occurrence, is independently selected from H,  
OH, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, CN, NO<sub>2</sub>,  
NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub>



alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl, C<sub>1</sub>-C<sub>4</sub> haloalkoxy,  
and C<sub>1</sub>-C<sub>4</sub> haloalkyl-S-;

- 5 R<sup>11</sup>, at each occurrence, is independently selected from  
H, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>18</sup>R<sup>19</sup>,  
C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, CF<sub>3</sub>;  
C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 0-3 R<sup>11a</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>11b</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>11b</sup>; or  
10 5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>11b</sup>;
- 15 R<sup>11a</sup>, at each occurrence, is independently selected from  
H, C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>,  
NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>;  
phenyl substituted with 0-3 R<sup>11b</sup>;  
C<sub>3</sub>-C<sub>6</sub> cycloalkyl substituted with 0-3 R<sup>11b</sup>; and  
20 5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>11b</sup>;
- 25 R<sup>11b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>,  
S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>,  
C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl,  
C<sub>1</sub>-C<sub>4</sub> haloalkoxy, and C<sub>1</sub>-C<sub>4</sub> halothioalkyl-S-;  
30 Z is H;  
C<sub>1</sub>-C<sub>8</sub> alkyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 1-3 R<sup>12</sup>;  
35 C<sub>1</sub>-C<sub>8</sub> alkyl substituted with 0-3 R<sup>12a</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>12a</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>12a</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;

C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-4 R<sup>12b</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
5 is substituted with 0-3 R<sup>12b</sup>;

R<sup>12</sup>, at each occurrence, is independently selected from  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-4 R<sup>12b</sup>; or  
10 5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>12b</sup>;

15 R<sup>12a</sup>, at each occurrence, is independently selected from  
H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, -C(=O)NR<sup>15</sup>R<sup>16</sup>,  
CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>,  
C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl,  
C<sub>1</sub>-C<sub>4</sub> haloalkoxy, or C<sub>1</sub>-C<sub>4</sub> halothioalkyl-S-;

20 R<sup>12b</sup>, at each occurrence, is independently selected from  
H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl,  
SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>,  
C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl,  
25 C<sub>1</sub>-C<sub>4</sub> haloalkoxy, and C<sub>1</sub>-C<sub>4</sub> halothioalkyl-S-;

R<sup>13</sup>, at each occurrence, is independently selected from  
H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, CN,  
NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;

30 R<sup>14</sup> is H, phenyl, benzyl, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkoxyalkyl, or  
C<sub>3</sub>-C<sub>6</sub> cycloalkyl;

R<sup>14a</sup> is H, phenyl, benzyl, or C<sub>1</sub>-C<sub>4</sub> alkyl;

35

R<sup>15</sup>, at each occurrence, is independently selected from H,  
C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-,  
and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-;

R<sup>16</sup>, at each occurrence, is independently selected from  
H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, phenethyl,  
(C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-;

5

R<sup>17</sup> is H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkoxyalkyl,  
aryl substituted by 0-4 R<sup>17a</sup>, or  
-CH<sub>2</sub>-aryl substituted by 0-4 R<sup>17a</sup>;

10 R<sup>17a</sup> is H, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
propoxy, butoxy, -OH, F, Cl, Br, I, CF<sub>3</sub>, OCF<sub>3</sub>, SCH<sub>3</sub>,  
S(O)CH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, -NH<sub>2</sub>, -N(CH<sub>3</sub>)<sub>2</sub>, or C<sub>1</sub>-C<sub>4</sub> haloalkyl;

15 R<sup>18</sup>, at each occurrence, is independently selected from  
H, C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl, benzyl, phenethyl,  
(C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-; and

R<sup>19</sup>, at each occurrence, is independently selected from  
H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl, benzyl, phenethyl,  
20 (C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-;

provided, when R<sup>13</sup> is H,

then Z is H;

25 C<sub>4</sub>-C<sub>8</sub> alkyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 1-3 R<sup>12</sup>;  
C<sub>1</sub>-C<sub>8</sub> alkyl substituted with 0-3 R<sup>12a</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>12a</sup>; or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>12a</sup>; and

30

provided, when ring B is a 1,3,4,5-tetrahydro-1-(Z)-5-  
(R<sup>10</sup>)-6,6,7,7-tetra(R<sup>11</sup>)-2,4-dioxo-2H-1,5-diazepin-3-yl  
core, and R<sup>13</sup> is H; then

35 R<sup>10</sup> is H, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>,  
S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, S(=O)<sub>2</sub>R<sup>17</sup>; or  
C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 0-3 R<sup>10a</sup>;

R<sup>10a</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>.

- 5    22. A pharmaceutical composition comprising a compound according to one of Claims 1-20 and a pharmaceutically acceptable carrier.
- 10    23. A method for the treatment of neurological disorders associated with  $\beta$ -amyloid production comprising administering to a host in need of such treatment a therapeutically effective amount of a compound according to one of Claims 1-20.
- 15    24. A method for inhibiting  $\gamma$ -secretase activity comprising administering to a host in need of such inhibition a therapeutically effective amount of a compound according to one of Claims 1-20 that inhibits  $\gamma$ -secretase activity.

20

TITLESUCCINOYLAMINO BENZODIAZEPINES AS INHIBITORS OF A $\beta$  PROTEIN  
PRODUCTION5                   FIELD OF THE INVENTION

This invention relates to novel lactams having drug and bio-affecting properties, their pharmaceutical compositions and methods of use. These novel compounds inhibit the processing of amyloid precursor protein and, 10 more specifically, inhibit the production of A $\beta$ -peptide, thereby acting to prevent the formation of neurological deposits of amyloid protein. More particularly, the present invention relates to the treatment of neurological disorders related to  $\beta$ -amyloid production such as 15 Alzheimer's disease and Down's Syndrome.

BACKGROUND OF THE INVENTION

Alzheimer's disease (AD) is a degenerative brain disorder characterized clinically by progressive loss of 20 memory, temporal and local orientation, cognition, reasoning, judgment and emotional stability. AD is a common cause of progressive dementia in humans and is one of the major causes of death in the United States. AD has been observed in all races and ethnic groups worldwide, and 25 is a major present and future health problem. No treatment that effectively prevents AD or reverses the clinical symptoms and underlying pathophysiology is currently available (for review, Selkoe, 1994).

Histopathological examination of brain tissue derived 30 upon autopsy or from neurosurgical specimens in effected individuals revealed the occurrence of amyloid plaques and neurofibrillar tangles in the cerebral cortex of such patients. Similar alterations were observed in patients with Trisomy 21 (Down's syndrome), and hereditary cerebral 35 hemorrhage with amyloidosis of the Dutch-type. Neurofibrillar tangles are nonmembrane-bound bundles of abnormal proteinaceous filaments and biochemical and immunochemical studies led to the conclusion that their

principle protein subunit is an altered phosphorylated form of the tau protein (reviewed in Selkoe, 1994).

Biochemical and immunological studies revealed that the dominant proteinaceous component of the amyloid plaque is an approximately 4.2 kilodalton (kD) protein of about 39 to 43 amino acids. This protein was designated A $\beta$ ,  $\beta$ -amyloid peptide, and sometimes  $\beta$ /A4; referred to herein as A $\beta$ . In addition to its deposition in amyloid plaques, A $\beta$  is also found in the walls of meningeal and parenchymal arterioles, small arteries, capillaries, and sometimes, venules. A $\beta$  was first purified and a partial amino acid reported in 1984 (Glennner and Wong, Biochem. Biophys. Res. Commun. 120: 885-890). The isolation and sequence data for the first 28 amino acids are described in U.S. Pat. No 4,666,829.

Compelling evidence accumulated during the last decade revealed that A $\beta$  is an internal polypeptide derived from a type 1 integral membrane protein, termed  $\beta$  amyloid precursor protein (APP).  $\beta$  APP is normally produced by many cells both in vivo and in cultured cells, derived from various animals and humans. A $\beta$  is derived from cleavage of  $\beta$  APP by as yet unknown enzyme (protease) system(s), collectively termed secretases.

The existence of at least four proteolytic activities has been postulated. They include  $\beta$  secretase(s), generating the N-terminus of A $\beta$ ,  $\alpha$  secretase(s) cleaving around the 16/17 peptide bond in A $\beta$ , and  $\gamma$  secretases, generating C-terminal A $\beta$  fragments ending at position 38, 39, 40, 42, and 43 or generating C-terminal extended precursors which are subsequently truncated to the above polypeptides.

Several lines of evidence suggest that abnormal accumulation of A $\beta$  plays a key role in the pathogenesis of AD. Firstly, A $\beta$  is the major protein found in amyloid plaques. Secondly, A $\beta$  is neurotoxic and may be causally related to neuronal death observed in AD patients. Thirdly, missense DNA mutations at position 717 in the 770 isoform of  $\beta$  APP can be found in effected members but not


unaffected members of several families with a genetically determined (familiar) form of AD. In addition, several other  $\beta$  APP mutations have been described in familiar forms of AD. Fourthly, similar neuropathological changes have  
5 been observed in transgenic animals overexpressing mutant forms of human  $\beta$  APP. Fifthly, individuals with Down's syndrome have an increased gene dosage of  $\beta$  APP and develop early-onset AD. Taken together, these observations strongly suggest that  $A\beta$  depositions may be causally  
10 related to the AD.

It is hypothesized that inhibiting the production of  $A\beta$  will prevent and reduce neurological degeneration, by controlling the formation of amyloid plaques, reducing neurotoxicity and, generally, mediating the pathology  
15 associated with  $A\beta$  production. One method of treatment methods would therefore be based on drugs that inhibit the formation of  $A\beta$  in vivo.

Methods of treatment could target the formation of  $A\beta$  through the enzymes involved in the proteolytic processing  
20 of  $\beta$  amyloid precursor protein. Compounds that inhibit  $\beta$  or  $\gamma$  secretase activity, either directly or indirectly, could control the production of  $A\beta$ . Advantageously, compounds that specifically target  $\gamma$  secretases, could control the production of  $A\beta$ . Such inhibition of  $\beta$  or  
25  $\gamma$  secretases could thereby reduce production of  $A\beta$ , which, thereby, could reduce or prevent the neurological disorders associated with  $A\beta$  protein.


PCT publication number WO 96/29313 discloses the general formula:

30




covering metalloprotease inhibiting compounds useful for the treatment of diseases associated with excess and/or  
35 unwanted matrix metalloprotease activity, particularly collagenase and or stromelysin activity.

Compounds of general formula:




- 5 are disclosed in PCT publication number WO 95/22966 relating to matrix metalloprotease inhibitors. The compounds of the invention are useful for the treatment of conditions associated with the destruction of cartilage, including corneal ulceration, osteoporosis, periodontitis  
10 and cancer.

European Patent Application number EP 0652009A1 relates to the general formula:



- 15 and discloses compounds that are protease inhibitors that inhibit A $\beta$  production.

US Patent Number 5703129 discloses the general formula:




20

which covers 5-amino-6-cyclohexyl-4-hydroxy-hexanamide derivatives that inhibit A $\beta$  production and are useful in the treatment of Alzheimer's disease.

- Copending, commonly assigned U.S. patent application  
25 Serial Number 09/370089 filed August 7, 1999 (equivalent to international application PCT US99/17717) discloses lactams of general formula:





wherein the lactam ring B is substituted by succinamide and a carbocyclic, aryl, or heteroaryl group. These compounds inhibit the processing of amyloid precursor protein and, more specifically, inhibit the production of A $\beta$ -peptide, thereby acting to prevent the formation of neurological deposits of amyloid protein.

None of the above references teaches or suggests the compounds of the present invention which are described in detail below.


#### SUMMARY OF THE INVENTION

One object of the present invention is to provide novel compounds which are useful as inhibitors of the production of A $\beta$  protein or pharmaceutically acceptable salts or prodrugs thereof.

It is another object of the present invention to provide pharmaceutical compositions comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of at least one of the compounds of the present invention or a pharmaceutically acceptable salt or prodrug form thereof.

It is another object of the present invention to provide a method for treating degenerative neurological disorders comprising administering to a host in need of such treatment a therapeutically effective amount of at least one of the compounds of the present invention or a pharmaceutically acceptable salt or prodrug form thereof.

These and other objects, which will become apparent during the following detailed description, have been achieved by the inventors' discovery that compounds of Formula (I):




(I)

or pharmaceutically acceptable salt or prodrug forms thereof, wherein  $R^3$ ,  $R^{3a}$ ,  $R^5$ ,  $R^{5a}$ ,  $R^6$ ,  $Q$ ,  $B$ , and  $Z$  are defined below, are effective inhibitors of the production of  $A\beta$ .

#### DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

Thus, in a first embodiment, the present invention provides a novel compound of Formula (I):



(I)

or a pharmaceutically acceptable salt or prodrug thereof, wherein:

$Q$  is  $-NR^1R^2$ ;

$R^1$ , at each occurrence, is independently selected from:

H;

$C_1$ - $C_6$  alkyl substituted with 0-3  $R^{1a}$ ;

$C_3$ - $C_{10}$  carbocycle substituted with 0-3  $R^{1b}$ ;

$C_6$ - $C_{10}$  aryl substituted with 0-3  $R^{1b}$ ; and

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3  $R^{1b}$ ;

$R^{1a}$ , at each occurrence, is independently selected from H,

$C_1$ - $C_6$  alkyl Cl, F, Br, I, =O, CN,  $NO_2$ ,  $NR^{15}R^{16}$ ,  $CF_3$ ;

$C_3$ - $C_{10}$  carbocycle substituted with 0-3  $R^{1b}$ ;

$C_6$ - $C_{10}$  aryl substituted with 0-3  $R^{1b}$ ; and

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R<sup>1b</sup>;

5

R<sup>1b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>6</sub> haloalkyl, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy;

10 R<sup>2</sup> is independently selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>10</sub> carbocycle, C<sub>6</sub>-C<sub>10</sub> aryl, and 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur;

15 R<sup>3</sup> is -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-R<sup>4</sup>,  
 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-S-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>,  
 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-O-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>,  
 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-N(R<sup>7b</sup>)-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>,  
 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-S(=O)-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>,  
 20 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-S(=O)<sub>2</sub>-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>,  
 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-C(=O)-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>,  
 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-N(R<sup>7b</sup>)C(=O)-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>,  
 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-C(=O)N(R<sup>7b</sup>)-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>,  
 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-N(R<sup>7b</sup>)S(=O)<sub>2</sub>-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>, or  
 25 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-S(=O)<sub>2</sub>N(R<sup>7b</sup>)-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>;

n is 0, 1, 2, or 3;

m is 0, 1, 2, or 3;

30

R<sup>3a</sup> is H, OH, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>2</sub>-C<sub>4</sub> alkenyl or C<sub>2</sub>-C<sub>4</sub> alkenyloxy;

R<sup>4</sup> is H, OH, OR<sup>14a</sup>,

35

C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 0-3 R<sup>4a</sup>,  
 C<sub>2</sub>-C<sub>6</sub> alkenyl substituted with 0-3 R<sup>4a</sup>,  
 C<sub>2</sub>-C<sub>6</sub> alkynyl substituted with 0-3 R<sup>4a</sup>,  
 C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,

- 5 C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>4b</sup>, or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>4b</sup>;
- R<sup>4a</sup>, at each occurrence, is independently selected from is  
H, F, Cl, Br, I, CF<sub>3</sub>,  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,  
10 C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>4b</sup>, or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>4b</sup>;
- 15 R<sup>4b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>,  
S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>,  
C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl,  
20 C<sub>1</sub>-C<sub>4</sub> haloalkoxy, and C<sub>1</sub>-C<sub>4</sub> halothioalkyl-S-;
- R<sup>5</sup> is H, OR<sup>14</sup>;  
C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 0-3 R<sup>5b</sup>;  
C<sub>1</sub>-C<sub>6</sub> alkoxy substituted with 0-3 R<sup>5b</sup>;  
25 C<sub>2</sub>-C<sub>6</sub> alkenyl substituted with 0-3 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>6</sub> alkynyl substituted with 0-3 R<sup>5b</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>5c</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>5c</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
30 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>5c</sup>;
- R<sup>5a</sup> is H, OH, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>2</sub>-C<sub>4</sub> alkenyl, or  
35 C<sub>2</sub>-C<sub>4</sub> alkenyloxy;
- R<sup>5b</sup>, at each occurrence, is independently selected from:

- H, C<sub>1</sub>-C<sub>6</sub> alkyl, CF<sub>3</sub>, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>5c</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>5c</sup>; or  
5 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R<sup>5c</sup>;
- 10 R<sup>5c</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl, C<sub>1</sub>-C<sub>4</sub> haloalkoxy, and C<sub>1</sub>-C<sub>4</sub> halothioalkyl-S-;
- 15 R<sup>6</sup> is H;  
C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 0-3 R<sup>6a</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>6b</sup>; or  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>6b</sup>;
- 20 R<sup>6a</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, aryl or CF<sub>3</sub>;
- 25 R<sup>6b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy;
- 30 R<sup>7</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, CF<sub>3</sub>, phenyl and C<sub>1</sub>-C<sub>4</sub> alkyl;
- R<sup>7a</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, CF<sub>3</sub>, and C<sub>1</sub>-C<sub>4</sub> alkyl;
- 35 R<sup>7b</sup> is independently selected from H and C<sub>1</sub>-C<sub>4</sub> alkyl;
- Ring B is a 7 membered lactam,

wherein the lactam is saturated, partially saturated or unsaturated;

wherein each additional lactam carbon is substituted with 0-2  $R^{11}$ ; and,

5 optionally, the lactam contains a heteroatom selected from -O-, -S-, -S(=O)-, -S(=O)<sub>2</sub>-, -N=, -NH-, and -N( $R^{10}$ )-;

10 additionally, two  $R^{11}$  substituents on adjacent atoms may be combined to form a benzo fused radical; wherein said benzo fused radical is substituted with 0-4  $R^{13}$ ;

15 additionally, two  $R^{11}$  substituents on adjacent atoms may be combined to form a 5 to 6 membered heteroaryl fused radical, wherein said 5 to 6 membered heteroaryl fused radical comprises 1 or 2 heteroatoms selected from N, O, and S; wherein said 5 to 6 membered heteroaryl fused radical is substituted with 0-3  $R^{13}$ ;

20 additionally, two  $R^{11}$  substituents on the same or adjacent carbon atoms may be combined to form a C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3  $R^{13}$ ;

25  $R^{10}$  is H, C(=O) $R^{17}$ , C(=O)OR<sup>17</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, S(=O)<sub>2</sub>R<sup>17</sup>;  
C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 0-3  $R^{10a}$ ;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4  $R^{10b}$ ;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3  $R^{10b}$ ; or  
5 to 10 membered heterocycle containing 1 to 4  
30 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3  $R^{10b}$ ;

35  $R^{10a}$ , at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, or aryl substituted with 0-4  $R^{10b}$ ;

R<sup>10b</sup>, at each occurrence, is independently selected from H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl, C<sub>1</sub>-C<sub>4</sub> haloalkoxy, and C<sub>1</sub>-C<sub>4</sub> haloalkyl-S-;

R<sup>11</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>18</sup>R<sup>19</sup>, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, CF<sub>3</sub>; C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 0-3 R<sup>11a</sup>; C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>11b</sup>; C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>11b</sup>; or 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R<sup>11b</sup>;

R<sup>11a</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>; phenyl substituted with 0-3 R<sup>11b</sup>; C<sub>3</sub>-C<sub>6</sub> cycloalkyl substituted with 0-3 R<sup>11b</sup>; and 5 to 6 membered heterocycle containing 1 to 3 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R<sup>11b</sup>;

R<sup>11b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl, C<sub>1</sub>-C<sub>4</sub> haloalkoxy, and C<sub>1</sub>-C<sub>4</sub> halothioalkyl-S-;

Z is H;  
C<sub>1</sub>-C<sub>8</sub> alkyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 1-3 R<sup>12</sup>;  
C<sub>1</sub>-C<sub>8</sub> alkyl substituted with 0-3 R<sup>12a</sup>;

- C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>12a</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>12a</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-4 R<sup>12b</sup>; or  
5 5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>12b</sup>;
- 10 R<sup>12</sup>, at each occurrence, is independently selected from  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-4 R<sup>12b</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
15 sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>12b</sup>;
- R<sup>12a</sup>, at each occurrence, is independently selected from  
H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, -C(=O)NR<sup>15</sup>R<sup>16</sup>,  
20 CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>,  
C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl,  
C<sub>1</sub>-C<sub>4</sub> haloalkoxy, or C<sub>1</sub>-C<sub>4</sub> halothioalkyl-S-;
- R<sup>12b</sup>, at each occurrence, is independently selected from  
25 H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl,  
SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>,  
C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl,  
C<sub>1</sub>-C<sub>4</sub> haloalkoxy, and C<sub>1</sub>-C<sub>4</sub> halothioalkyl-S-;
- 30 R<sup>13</sup>, at each occurrence, is independently selected from  
H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, CN,  
NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;
- R<sup>14</sup> is H, phenyl, benzyl, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkoxyalkyl, or  
35 C<sub>3</sub>-C<sub>6</sub> cycloalkyl;
- R<sup>14a</sup> is H, phenyl, benzyl, or C<sub>1</sub>-C<sub>4</sub> alkyl;



- R<sup>15</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-;
- 5 R<sup>16</sup>, at each occurrence, is independently selected from H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-;
- R<sup>17</sup> is H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkoxyalkyl, 10 aryl substituted by 0-4 R<sup>17a</sup>, or -CH<sub>2</sub>-aryl substituted by 0-4 R<sup>17a</sup>;
- R<sup>17a</sup> is H, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, butoxy, -OH, F, Cl, Br, I, CF<sub>3</sub>, OCF<sub>3</sub>, SCH<sub>3</sub>, 15 S(O)CH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, -NH<sub>2</sub>, -N(CH<sub>3</sub>)<sub>2</sub>, or C<sub>1</sub>-C<sub>4</sub> haloalkyl;
- R<sup>18</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-; and 20
- R<sup>19</sup>, at each occurrence, is independently selected from H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-;
- 25 provided, when R<sup>13</sup> is H, then Z is H; C<sub>4</sub>-C<sub>8</sub> alkyl substituted with 1-3 R<sup>12</sup>; C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 1-3 R<sup>12</sup>; C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 1-3 R<sup>12</sup>;
- 30 C<sub>1</sub>-C<sub>8</sub> alkyl substituted with 0-3 R<sup>12a</sup>; C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>12a</sup>; or C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>12a</sup>; and
- provided, when ring B is a 1,3,4,5-tetrahydro-1-(Z)-5-(R<sup>10</sup>)-6,6,7,7-tetra(R<sup>11</sup>)-2,4-dioxo-2H-1,5-diazepin-3-yl core, and R<sup>13</sup> is H; then

R<sup>10</sup> is H, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>,


$S(=O)_2NR^{18}R^{19}$ ,  $S(=O)_2R^{17}$ ; or

$C_1-C_6$  alkyl optionally substituted with 0-3  $R^{10a}$ ;

$R^{10a}$ , at each occurrence, is independently selected from

5 H,  $C_1-C_6$  alkyl,  $OR^{14}$ , Cl, F, Br, I, =O, CN,  $NO_2$ ,  
 $NR^{15}R^{16}$ , and  $CF_3$ .

[2] In a preferred embodiment the present invention  
 provides for a compound of Formula (Ia):



or a pharmaceutically acceptable salt or prodrug thereof,

15 wherein:

Z is H;

$C_1-C_8$  alkyl substituted with 0-3  $R^{12a}$ ;

$C_2-C_4$  alkenyl substituted with 0-3  $R^{12a}$ ; or

20  $C_2-C_4$  alkynyl substituted with 0-3  $R^{12a}$ .

[3] In a more preferred embodiment the present  
 invention provides for a compound of Formula (Ia):

25 wherein:

$R^3$  is  $-(CR^7R^{7a})_n-R^4$ ,

$-(CR^7R^{7a})_n-S-(CR^7R^{7a})_m-R^4$ ,

$-(CR^7R^{7a})_n-O-(CR^7R^{7a})_m-R^4$ , or

30  $-(CR^7R^{7a})_n-N(R^{7b})-(CR^7R^{7a})_m-R^4$ ;

n is 0, 1, or 2;

m is 0, 1, or 2;

35

R<sup>3a</sup> is H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, butoxy, allyl, or 3-buten-1-yl;

R<sup>4</sup> is H, OH, OR<sup>14a</sup>,

- 5 C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 0-3 R<sup>4a</sup>,  
C<sub>2</sub>-C<sub>6</sub> alkenyl substituted with 0-3 R<sup>4a</sup>,  
C<sub>2</sub>-C<sub>6</sub> alkynyl substituted with 0-3 R<sup>4a</sup>,  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>4b</sup>, or  
10 5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>4b</sup>;
- 15 R<sup>4a</sup>, at each occurrence, is independently selected from is  
H, F, Cl, Br, I, CF<sub>3</sub>,  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>4b</sup>, or  
5 to 10 membered heterocycle containing 1 to 4  
20 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>4b</sup>;
- R<sup>4b</sup>, at each occurrence, is independently selected from H,  
25 OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>,  
S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub>  
haloalkyl, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy;

R<sup>5</sup> is H, OR<sup>14</sup>;

- 30 C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 0-3 R<sup>5b</sup>;  
C<sub>1</sub>-C<sub>6</sub> alkoxy substituted with 0-3 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>6</sub> alkenyl substituted with 0-3 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>6</sub> alkynyl substituted with 0-3 R<sup>5b</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>5c</sup>;  
35 C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>5c</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R<sup>5c</sup>;

R<sup>5a</sup> is H or C<sub>1</sub>-C<sub>4</sub> alkyl;

5

R<sup>5b</sup>, at each occurrence, is independently selected from:

H, C<sub>1</sub>-C<sub>6</sub> alkyl, CF<sub>3</sub>, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>,  
NR<sup>15</sup>R<sup>16</sup>;

C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>5c</sup>;

10 C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>5c</sup>; or

5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>5c</sup>;

15

R<sup>5c</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>,  
S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub>  
haloalkyl, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy;

20

R<sup>6</sup> is H, methyl, or ethyl;

R<sup>7</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, CF<sub>3</sub>, phenyl and C<sub>1</sub>-C<sub>4</sub> alkyl;

25

R<sup>7a</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, CF<sub>3</sub>, and C<sub>1</sub>-C<sub>4</sub> alkyl;

R<sup>7b</sup> is independently selected from H, methyl, ethyl,  
30 propyl, and butyl;

Ring B is a 7 membered lactam,  
wherein the lactam is saturated, partially saturated  
or unsaturated;  
35 wherein each additional lactam carbon is substituted  
with 0-2 R<sup>11</sup>; and,

optionally, the lactam contains a heteroatom selected from, -O-, -S-, -S(=O)-, -S(=O)<sub>2</sub>-, -N=, -NH-, and -N(R<sup>10</sup>)-;

5 additionally, two R<sup>11</sup> substituents on adjacent atoms may be combined to form a benzo fused radical; wherein said benzo fused radical is substituted with 0-3 R<sup>13</sup>;

10 additionally, two R<sup>11</sup> substituents on adjacent atoms may be combined to form a 5 to 6 membered heteroaryl fused radical, wherein said 5 to 6 membered heteroaryl fused radical comprises 1 or 2 heteroatoms selected from N, O, and S; wherein said 5 to 6 membered heteroaryl fused radical is substituted with 0-3 R<sup>13</sup>;

15 additionally, two R<sup>11</sup> substituents on the same or adjacent carbon atoms may be combined to form a C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>13</sup>;

20 R<sup>10</sup> is H, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, S(=O)<sub>2</sub>R<sup>17</sup>;  
C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 0-2 R<sup>10a</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>10b</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>10b</sup>; or

25 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R<sup>10b</sup>;

30 R<sup>10a</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, or phenyl substituted with 0-4 R<sup>10b</sup>;

35 R<sup>10b</sup>, at each occurrence, is independently selected from H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, or CF<sub>3</sub>;

R<sup>11</sup>, at each occurrence, is independently selected from

- H, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>18</sup>R<sup>19</sup>,  
C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, CF<sub>3</sub>;  
C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 0-3 R<sup>11a</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>11b</sup>;  
5 C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>11b</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>11b</sup>;  
10  
R<sup>11a</sup>, at each occurrence, is independently selected from H,  
C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>,  
CF<sub>3</sub>, or phenyl substituted with 0-3 R<sup>11b</sup>;  
15 R<sup>11b</sup>, at each occurrence, is independently selected from  
H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl,  
SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy,  
C<sub>1</sub>-C<sub>4</sub> haloalkyl, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy;  
20 Z is H;  
C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 0-3 R<sup>12a</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>12a</sup>; or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>12a</sup>;  
25 R<sup>12a</sup>, at each occurrence, is independently selected from  
H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl,  
SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy,  
C<sub>1</sub>-C<sub>4</sub> haloalkyl, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy;  
30 R<sup>13</sup>, at each occurrence, is independently selected from  
H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, CN,  
NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;  
R<sup>14</sup> is H, phenyl, benzyl, C<sub>1</sub>-C<sub>6</sub> alkyl, or C<sub>2</sub>-C<sub>6</sub> alkoxyalkyl;  
35 R<sup>14a</sup> is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;

R<sup>15</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-;

- 5 R<sup>16</sup>, at each occurrence, is independently selected from H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-;

- R<sup>17</sup> is H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkoxyalkyl,  
10 aryl substituted by 0-4 R<sup>17a</sup>, or  
-CH<sub>2</sub>-aryl substituted by 0-4 R<sup>17a</sup>;

- R<sup>17a</sup> is H, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
propoxy, butoxy, -OH, F, Cl, Br, I, CF<sub>3</sub>, OCF<sub>3</sub>, SCH<sub>3</sub>,  
15 S(O)CH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, -NH<sub>2</sub>, -N(CH<sub>3</sub>)<sub>2</sub>, or C<sub>1</sub>-C<sub>4</sub> haloalkyl;

- R<sup>18</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl, benzyl, phenethyl,  
(C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-; and  
20

- R<sup>19</sup>, at each occurrence, is independently selected from H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl, benzyl, phenethyl,  
(C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-.

- 25 [4] In a further more preferred embodiment the present invention provides for a compound of Formula (Ia) wherein:

R<sup>3</sup> is -(CHR<sup>7</sup>)<sub>n</sub>-R<sup>4</sup>,

- 30 n is 0 or 1;

R<sup>3a</sup> is H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, butoxy, allyl, or 3-buten-1-yl;

- 35 R<sup>4</sup> is H, OH, OR<sup>4a</sup>,  
C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-2 R<sup>4a</sup>,  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-2 R<sup>4a</sup>,  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>4a</sup>,

C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>4b</sup>, or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
5 sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>4b</sup>;

R<sup>4a</sup>, at each occurrence, is independently selected from is  
H, F, Cl, Br, I, CF<sub>3</sub>,  
10 C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,  
phenyl substituted with 0-3 R<sup>4b</sup>, or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
15 substituted with 0-3 R<sup>4b</sup>;

R<sup>4b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>,  
S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkoxy, C<sub>1</sub>-C<sub>2</sub>  
20 haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

R<sup>5</sup> is H, OR<sup>14</sup>;  
C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-3 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>5b</sup>;  
25 C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>5b</sup>;

R<sup>5a</sup> is H, methyl, ethyl, propyl, or butyl;

R<sup>5b</sup>, at each occurrence, is independently selected from:  
30 H, methyl, ethyl, propyl, butyl, CF<sub>3</sub>, OR<sup>14</sup>, Cl, F, Br,  
I, =O;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>5c</sup>;  
phenyl substituted with 0-3 R<sup>5c</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
35 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>5c</sup>;



R<sup>5c</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

5

R<sup>6</sup> is H;

R<sup>7</sup>, at each occurrence, is independently selected from H, F, CF<sub>3</sub>, methyl, and ethyl;

10

Ring B is a 7 membered lactam, wherein the lactam is saturated, partially saturated or unsaturated; wherein each additional lactam carbon is substituted with 0-2 R<sup>11</sup>; and, optionally, the lactam contains a heteroatom selected from -N=, -NH-, and -N(R<sup>10</sup>)-;

15

additionally, two R<sup>11</sup> substituents on adjacent atoms may be combined to form a benzo fused radical; wherein said benzo fused radical is substituted with 0-2 R<sup>13</sup>;

20

additionally, two R<sup>11</sup> substituents on adjacent atoms may be combined to form a 5 to 6 membered heteroaryl fused radical, wherein said 5 to 6 membered heteroaryl fused radical comprises 1 or 2 heteroatoms selected from N, O, and S; wherein said 5 to 6 membered heteroaryl fused radical is substituted with 0-2 R<sup>13</sup>;

25

additionally, two R<sup>11</sup> substituents on the same or adjacent carbon atoms may be combined to form a C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-2 R<sup>13</sup>;

30

R<sup>10</sup> is H, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>; C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with 0-1 R<sup>10a</sup>; phenyl substituted with 0-4 R<sup>10b</sup>; C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>10b</sup>; or

35

5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R<sup>10b</sup>;

5

R<sup>10a</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>4</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, or phenyl substituted with 0-4 R<sup>10b</sup>;

10 R<sup>10b</sup>, at each occurrence, is independently selected from H, OH, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkoxy, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, or CF<sub>3</sub>;

15 R<sup>11</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, =O, NR<sup>18</sup>R<sup>19</sup>, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>, CF<sub>3</sub>;

C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 0-3 R<sup>11a</sup>;

C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>11b</sup>;

C<sub>3</sub>-6 carbocycle substituted with 0-3 R<sup>11b</sup>; or

20 5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R<sup>11b</sup>;

25 R<sup>11a</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>4</sub> alkyl, OR<sup>14</sup>, F, =O, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, or phenyl substituted with 0-3 R<sup>11b</sup>;

30 R<sup>11b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

Z is H;

C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-3 R<sup>12a</sup>;

35 C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>12a</sup>; or

C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>12a</sup>;

R<sup>12a</sup>, at each occurrence, is independently selected from

H, OH, Cl, F,  $\text{NR}^{15}\text{R}^{16}$ ,  $\text{CF}_3$ , acetyl,  $\text{SCH}_3$ ,  $\text{S(=O)CH}_3$ ,  $\text{S(=O)}_2\text{CH}_3$ ,  $\text{C}_1\text{-C}_4$  alkyl,  $\text{C}_1\text{-C}_3$  alkoxy,  $\text{C}_1\text{-C}_2$  haloalkyl, and  $\text{C}_1\text{-C}_2$  haloalkoxy;

- 5  $\text{R}^{13}$ , at each occurrence, is independently selected from H, OH,  $\text{C}_1\text{-C}_6$  alkyl,  $\text{C}_1\text{-C}_4$  alkoxy, Cl, F, Br, I, CN,  $\text{NO}_2$ ,  $\text{NR}^{15}\text{R}^{16}$ , and  $\text{CF}_3$ ;

$\text{R}^{14}$  is H, phenyl, benzyl,  $\text{C}_1\text{-C}_4$  alkyl, or  $\text{C}_2\text{-C}_4$  alkoxyalkyl;

10

- $\text{R}^{15}$ , at each occurrence, is independently selected from H,  $\text{C}_1\text{-C}_4$  alkyl, benzyl, phenethyl,  $(\text{C}_1\text{-C}_4 \text{ alkyl})\text{-C(=O)-}$ , and  $(\text{C}_1\text{-C}_4 \text{ alkyl})\text{-S(=O)}_2\text{-}$ ;

- 15  $\text{R}^{16}$ , at each occurrence, is independently selected from H, OH,  $\text{C}_1\text{-C}_4$  alkyl, benzyl, phenethyl,  $(\text{C}_1\text{-C}_4 \text{ alkyl})\text{-C(=O)-}$ , and  $(\text{C}_1\text{-C}_4 \text{ alkyl})\text{-S(=O)}_2\text{-}$ ;

- 20  $\text{R}^{17}$  is H, methyl, ethyl, propyl, butyl, methoxymethyl, ethoxymethyl, methoxyethyl, ethoxyethyl, phenyl substituted by 0-3  $\text{R}^{17a}$ , or  $\text{-CH}_2\text{-phenyl}$  substituted by 0-3  $\text{R}^{17a}$ ;

$\text{R}^{17a}$  is H, methyl, methoxy, -OH, F, Cl,  $\text{CF}_3$ , or  $\text{OCF}_3$ ;


25

- $\text{R}^{18}$ , at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl;

- 30  $\text{R}^{19}$ , at each occurrence, is independently selected from H, methyl, and ethyl.


[5] In an even more preferred embodiment the present invention provides for a compound of Formula (Ib):

35




or a pharmaceutically acceptable salt or prodrug thereof  
 5 wherein:

Ring B is selected from:



15 [6] In an even further more preferred embodiment the present invention provides for a compound of Formula (Ic):



or a pharmaceutically acceptable salt or prodrug thereof  
wherein

R<sup>3</sup> is R<sup>4</sup>,

5

R<sup>4</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>4a</sup>,  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>4a</sup>, or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>4a</sup>;

10 R<sup>4a</sup>, at each occurrence, is independently selected from  
H, F, CF<sub>3</sub>,  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,  
phenyl substituted with 0-3 R<sup>4b</sup>, or  
5 to 6 membered heterocycle containing 1 to 4  
15 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>4b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
20 pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>4b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
25 S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

R<sup>5</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>5b</sup>;  
30 C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>5b</sup>;

R<sup>5b</sup>, at each occurrence, is independently selected from:  
H, methyl, ethyl, propyl, butyl, CF<sub>3</sub>, OR<sup>14</sup>, =O;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-2 R<sup>5c</sup>;  
35 phenyl substituted with 0-3 R<sup>5c</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is

- substituted with 0-3 R<sup>5c</sup>; wherein said 5 to 6 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;
- R<sup>5c</sup>, at each occurrence, is independently selected from H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;
- R<sup>11</sup>, at each occurrence, is independently selected from H, =O, NR<sup>18</sup>R<sup>19</sup>, CF<sub>3</sub>; C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with 0-1 R<sup>11a</sup>; phenyl substituted with 0-3 R<sup>11b</sup>; C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>11b</sup>; and 5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R<sup>11b</sup>; wherein said 5 to 6 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;
- R<sup>11a</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>4</sub> alkyl, OR<sup>14</sup>, F, Cl, =O, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, or phenyl substituted with 0-3 R<sup>11b</sup>;
- R<sup>11b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;
- Z is H; C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-3 R<sup>12a</sup>; C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>12a</sup>; or C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>12a</sup>;

R<sup>13</sup>, at each occurrence, is independently selected from H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy, Cl, F, Br, CN, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;

R<sup>14</sup> is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;


R<sup>15</sup>, at each occurrence, is independently selected from H, methyl, ethyl, propyl, and butyl;

R<sup>16</sup>, at each occurrence, is independently selected from H, OH, methyl, ethyl, propyl, butyl, benzyl, phenethyl, methyl-C(=O)-, ethyl-C(=O)-, methyl-S(=O)<sub>2</sub>-, and ethyl-S(=O)<sub>2</sub>-;

R<sup>18</sup>, at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl; and

25 R<sup>19</sup>, at each occurrence, is independently selected from  
H, methyl, and ethyl.

[7] In another even further more preferred embodiment the present invention provides for a compound of Formula (Id):



(Id)

or a pharmaceutically acceptable salt or prodrug thereof  
wherein:

R<sup>3</sup> is R<sup>4</sup>,

5

R<sup>4</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>4a</sup>,  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>4a</sup>, or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>4a</sup>;

10 R<sup>4a</sup>, at each occurrence, is independently selected from  
H, F, CF<sub>3</sub>,  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,  
phenyl substituted with 0-3 R<sup>4b</sup>, or  
5 to 6 membered heterocycle containing 1 to 4  
15 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>4b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
20 pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>4b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
25 S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

R<sup>5</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>5b</sup>;  
30 C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>5b</sup>;

R<sup>5b</sup>, at each occurrence, is independently selected from:  
H, methyl, ethyl, propyl, butyl, CF<sub>3</sub>, OR<sup>14</sup>, =O;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-2 R<sup>5c</sup>;  
35 phenyl substituted with 0-3 R<sup>5c</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is



substituted with 0-3 R<sup>5c</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
5 imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>5c</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
10 ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

R<sup>11</sup>, at each occurrence, is independently selected from  
H, =O, NR<sup>18</sup>R<sup>19</sup>, CF<sub>3</sub>;  
C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with 0-1 R<sup>11a</sup>;  
15 phenyl substituted with 0-3 R<sup>11b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>11b</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
20 substituted with 0-3 R<sup>11b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

25 R<sup>11a</sup>, at each occurrence, is independently selected from H,  
C<sub>1</sub>-C<sub>4</sub> alkyl, OR<sup>14</sup>, F, Cl, =O, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, or phenyl  
substituted with 0-3 R<sup>11b</sup>;

30 R<sup>11b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, methyl, ethyl, propyl, butyl,  
methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub>  
haloalkoxy;

35 Z is H;  
C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-3 R<sup>12a</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>12a</sup>; or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>12a</sup>;

R<sup>12a</sup>, at each occurrence, is independently selected from  
 H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
 S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
 5 ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

R<sup>13</sup>, at each occurrence, is independently selected from  
 H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
 Cl, F, Br, CN, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;

10 R<sup>14</sup> is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;


R<sup>15</sup>, at each occurrence, is independently selected from H,  
 methyl, ethyl, propyl, and butyl;

15 R<sup>16</sup>, at each occurrence, is independently selected from  
 H, OH, methyl, ethyl, propyl, butyl, benzyl,  
 phenethyl, methyl-C(=O)-, ethyl-C(=O)-,  
 methyl-S(=O)<sub>2</sub>-, and ethyl-S(=O)<sub>2</sub>-;

20 R<sup>18</sup>, at each occurrence, is independently selected from  
 H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and  
 phenethyl; and

25 R<sup>19</sup>, at each occurrence, is independently selected from  
 H, methyl, and ethyl.

[8] In another even further more preferred embodiment  
 the present invention provides for a compound of Formula  
 30 (Ie):



(Ie)

or a pharmaceutically acceptable salt or prodrug thereof  
wherein:

R<sup>3</sup> is R<sup>4</sup>,

5

R<sup>4</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>4a</sup>,  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>4a</sup>, or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>4a</sup>;

10 R<sup>4a</sup>, at each occurrence, is independently selected from  
H, F, CF<sub>3</sub>,

C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,

phenyl substituted with 0-3 R<sup>4b</sup>, or

5 to 6 membered heterocycle containing 1 to 4

15 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>4b</sup>; wherein said 5 to 6

membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,

20 pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>4b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,

25 S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

R<sup>5</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>5b</sup>;

C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>5b</sup>;

30 C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>5b</sup>;

R<sup>5b</sup>, at each occurrence, is independently selected from:

H, methyl, ethyl, propyl, butyl, CF<sub>3</sub>, OR<sup>14</sup>, =O;

C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-2 R<sup>5c</sup>;

35 phenyl substituted with 0-3 R<sup>5c</sup>; or

5 to 6 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 6 membered heterocycle is

substituted with 0-3 R<sup>5c</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
5 imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>5c</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
10 ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

R<sup>10</sup> is H, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>;  
C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with 0-1 R<sup>10a</sup>;  
phenyl substituted with 0-4 R<sup>10b</sup>;  
15 C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>10b</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>10b</sup>; wherein said 5 to 6  
20 membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

25 R<sup>10a</sup>, at each occurrence, is independently selected from H,  
methyl, ethyl, propyl, butyl, OR<sup>14</sup>, Cl, F, =O, NR<sup>15</sup>R<sup>16</sup>,  
CF<sub>3</sub>, or phenyl substituted with 0-4 R<sup>10b</sup>;

R<sup>10b</sup>, at each occurrence, is independently selected from H,  
30 OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
propoxy, Cl, F, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;

Z is H;  
C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-3 R<sup>12a</sup>;  
35 C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>12a</sup>; or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>12a</sup>;

R<sup>12a</sup>, at each occurrence, is independently selected from

H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

- 5 R<sup>13</sup>, at each occurrence, is independently selected from  
H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
Cl, F, Br, CN, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;

10 R<sup>14</sup> is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;

- R<sup>15</sup>, at each occurrence, is independently selected from H,  
methyl, ethyl, propyl, and butyl;

- 15 R<sup>16</sup>, at each occurrence, is independently selected from  
H, OH, methyl, ethyl, propyl, butyl, benzyl,  
phenethyl, methyl-C(=O)-, ethyl-C(=O)-,  
methyl-S(=O)<sub>2</sub>-, and ethyl-S(=O)<sub>2</sub>-;


- 20 R<sup>17</sup> is H, methyl, ethyl, propyl, butyl, methoxymethyl,  
ethoxymethyl, methoxyethyl, ethoxyethyl,  
phenyl substituted by 0-3 R<sup>17a</sup>, or  
-CH<sub>2</sub>-phenyl substituted by 0-3 R<sup>17a</sup>;

- 25 R<sup>17a</sup> is H, methyl, methoxy, -OH, F, Cl, CF<sub>3</sub>, or OCF<sub>3</sub>;

- R<sup>18</sup>, at each occurrence, is independently selected from  
H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and  
phenethyl; and

- 30 R<sup>19</sup>, at each occurrence, is independently selected from  
H, methyl, and ethyl.

[9] In another even further more preferred embodiment  
the present invention provides for a compound of Formula  
35 (If):



(If)

or a pharmaceutically acceptable salt or prodrug thereof  
wherein:

5

$R^3$  is  $R^4$ ,

$R^4$  is  $C_1$ - $C_4$  alkyl substituted with 0-1  $R^{4a}$ ,  
 $C_2$ - $C_4$  alkenyl substituted with 0-1  $R^{4a}$ , or  
10  $C_2$ - $C_4$  alkynyl substituted with 0-1  $R^{4a}$ ;

$R^{4a}$ , at each occurrence, is independently selected from  
H, F,  $CF_3$ ,

15

$C_3$ - $C_6$  carbocycle substituted with 0-3  $R^{4b}$ ,

phenyl substituted with 0-3  $R^{4b}$ , or

5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3  $R^{4b}$ ; wherein said 5 to 6

20

membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

25  $R^{4b}$ , at each occurrence, is independently selected from H,  
OH, Cl, F,  $NR^{15}R^{16}$ ,  $CF_3$ , acetyl,  $SCH_3$ ,  $S(=O)CH_3$ ,  
 $S(=O)_2CH_3$ , methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy,  $C_1$ - $C_2$  haloalkyl, and  $C_1$ - $C_2$  haloalkoxy;

30  $R^5$  is  $C_1$ - $C_4$  alkyl substituted with 0-1  $R^{5b}$ ;  
 $C_2$ - $C_4$  alkenyl substituted with 0-1  $R^{5b}$ ;  
 $C_2$ - $C_4$  alkynyl substituted with 0-1  $R^{5b}$ ;

- R<sup>5b</sup>, at each occurrence, is independently selected from:  
H, methyl, ethyl, propyl, butyl, CF<sub>3</sub>, OR<sup>14</sup>, =O;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-2 R<sup>5c</sup>;  
phenyl substituted with 0-3 R<sup>5c</sup>; or
- 5 5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>5c</sup>; wherein said 5 to 6  
10 membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;
- 15 R<sup>5c</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;
- Z is H;
- 20 C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-3 R<sup>12a</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>12a</sup>; or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>12a</sup>;
- 25 R<sup>12a</sup>, at each occurrence, is independently selected from  
H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;
- 30 R<sup>13</sup>, at each occurrence, is independently selected from  
H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
Cl, F, Br, CN, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;
- R<sup>14</sup> is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;
- 35 R<sup>15</sup>, at each occurrence, is independently selected from H,  
methyl, ethyl, propyl, and butyl;
- R<sup>16</sup>, at each occurrence, is independently selected from

H, OH, methyl, ethyl, propyl, butyl, benzyl, phenethyl, methyl-C(=O)-, ethyl-C(=O)-, methyl-S(=O)<sub>2</sub>-, and ethyl-S(=O)<sub>2</sub>-;

5 R<sup>18</sup>, at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl; and

10 R<sup>19</sup>, at each occurrence, is independently selected from H, methyl, and ethyl.

[10] In another even further more preferred embodiment the present invention provides for a compound of one of Formulas (Ic), (Id), (Ie), or (If), wherein:

15

R<sup>3</sup> is -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>,  
-CH<sub>2</sub>(CH<sub>3</sub>)<sub>2</sub>, -CH(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>C(CH<sub>3</sub>)<sub>3</sub>,  
-CF<sub>3</sub>, -CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CF<sub>3</sub>,  
-CH=CH<sub>2</sub>, -CH<sub>2</sub>CH=CH<sub>2</sub>, -CH<sub>2</sub>C(CH<sub>3</sub>)=CH<sub>2</sub>, -CH<sub>2</sub>CH=C(CH<sub>3</sub>)<sub>2</sub>,  
20 -CH<sub>2</sub>CH<sub>2</sub>CH=CH<sub>2</sub>, -CH<sub>2</sub>CH<sub>2</sub>C(CH<sub>3</sub>)=CH<sub>2</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH=C(CH<sub>3</sub>)<sub>2</sub>,  
cis-CH<sub>2</sub>CH=CH(CH<sub>3</sub>), cis-CH<sub>2</sub>CH<sub>2</sub>CH=CH(CH<sub>3</sub>),  
trans-CH<sub>2</sub>CH=CH(CH<sub>3</sub>), trans-CH<sub>2</sub>CH<sub>2</sub>CH=CH(CH<sub>3</sub>);  
-C≡CH, -CH<sub>2</sub>C≡CH, -CH<sub>2</sub>C≡C(CH<sub>3</sub>),  
cyclopropyl-CH<sub>2</sub>-, cyclobutyl-CH<sub>2</sub>-, cyclopentyl-CH<sub>2</sub>-,  
25 cyclohexyl-CH<sub>2</sub>-, cyclopropyl-CH<sub>2</sub>CH<sub>2</sub>-,  
cyclobutyl-CH<sub>2</sub>CH<sub>2</sub>-, cyclopentyl-CH<sub>2</sub>CH<sub>2</sub>-,  
cyclohexyl-CH<sub>2</sub>CH<sub>2</sub>-, phenyl-CH<sub>2</sub>-,  
(2-F-phenyl)CH<sub>2</sub>-, (3-F-phenyl)CH<sub>2</sub>-, (4-F-phenyl)CH<sub>2</sub>-,  
(2-Cl-phenyl)CH<sub>2</sub>-, (3-Cl-phenyl)CH<sub>2</sub>-, (4-Cl-phenyl)CH<sub>2</sub>-,  
30 (2,3-diF-phenyl)CH<sub>2</sub>-, (2,4-diF-phenyl)CH<sub>2</sub>-,  
(2,5-diF-phenyl)CH<sub>2</sub>-, (2,6-diF-phenyl)CH<sub>2</sub>-,  
(3,4-diF-phenyl)CH<sub>2</sub>-, (3,5-diF-phenyl)CH<sub>2</sub>-,  
(2,3-diCl-phenyl)CH<sub>2</sub>-, (2,4-diCl-phenyl)CH<sub>2</sub>-,  
(2,5-diCl-phenyl)CH<sub>2</sub>-, (2,6-diCl-phenyl)CH<sub>2</sub>-,  
35 (3,4-diCl-phenyl)CH<sub>2</sub>-, (3,5-diCl-phenyl)CH<sub>2</sub>-,  
(3-F-4-Cl-phenyl)CH<sub>2</sub>-, (3-F-5-Cl-phenyl)CH<sub>2</sub>-,  
(3-Cl-4-F-phenyl)CH<sub>2</sub>-, phenyl-CH<sub>2</sub>CH<sub>2</sub>-,  
(2-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,



- (4-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (3-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (4-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2,3-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2,4-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2,5-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2,6-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 5 (3,4-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3,5-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2,3-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2,4-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2,5-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2,6-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (3,4-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3,5-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (3-F-4-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, or (3-F-5-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 10
- R<sup>5</sup> is -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>,  
 -CH(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>C(CH<sub>3</sub>)<sub>3</sub>,  
 -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -CH(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>3</sub>,  
 -CH<sub>2</sub>CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>, -CF<sub>3</sub>, -CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CF<sub>3</sub>,  
 15 -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CF<sub>3</sub>, -CH=CH<sub>2</sub>, -CH<sub>2</sub>CH=CH<sub>2</sub>,  
 -CH=CHCH<sub>3</sub>, cis-CH<sub>2</sub>CH=CH(CH<sub>3</sub>), trans-CH<sub>2</sub>CH=CH(CH<sub>3</sub>),  
 trans-CH<sub>2</sub>CH=CH(C<sub>6</sub>H<sub>5</sub>), -CH<sub>2</sub>CH=C(CH<sub>3</sub>)<sub>2</sub>, cis-CH<sub>2</sub>CH=CHCH<sub>2</sub>CH<sub>3</sub>,  
 trans-CH<sub>2</sub>CH=CHCH<sub>2</sub>CH<sub>3</sub>, cis-CH<sub>2</sub>CH<sub>2</sub>CH=CH(CH<sub>3</sub>),  
 trans-CH<sub>2</sub>CH<sub>2</sub>CH=CH(CH<sub>3</sub>), trans-CH<sub>2</sub>CH=CHCH<sub>2</sub>(C<sub>6</sub>H<sub>5</sub>),  
 20 -C≡CH, -CH<sub>2</sub>C≡CH, -CH<sub>2</sub>C≡C(CH<sub>3</sub>), -CH<sub>2</sub>C≡C(C<sub>6</sub>H<sub>5</sub>)  
 -CH<sub>2</sub>CH<sub>2</sub>C≡CH, -CH<sub>2</sub>CH<sub>2</sub>C≡C(CH<sub>3</sub>), -CH<sub>2</sub>CH<sub>2</sub>C≡C(C<sub>6</sub>H<sub>5</sub>)  
 -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C≡CH, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C≡C(CH<sub>3</sub>), -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C≡C(C<sub>6</sub>H<sub>5</sub>)  
 cyclopropyl-CH<sub>2</sub>-, cyclobutyl-CH<sub>2</sub>-, cyclopentyl-CH<sub>2</sub>-,  
 cyclohexyl-CH<sub>2</sub>-, (2-CH<sub>3</sub>-cyclopropyl)CH<sub>2</sub>-,  
 25 (3-CH<sub>3</sub>-cyclobutyl)CH<sub>2</sub>-,  
 cyclopropyl-CH<sub>2</sub>CH<sub>2</sub>-, cyclobutyl-CH<sub>2</sub>CH<sub>2</sub>-,  
 cyclopentyl-CH<sub>2</sub>CH<sub>2</sub>-, cyclohexyl-CH<sub>2</sub>CH<sub>2</sub>-,  
 (2-CH<sub>3</sub>-cyclopropyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-CH<sub>3</sub>-cyclobutyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 phenyl-CH<sub>2</sub>-, (2-F-phenyl)CH<sub>2</sub>-, (3-F-phenyl)CH<sub>2</sub>-,  
 30 (4-F-phenyl)CH<sub>2</sub>-, furanyl-CH<sub>2</sub>-, thienyl-CH<sub>2</sub>-,  
 pyridyl-CH<sub>2</sub>-, 1-imidazolyl-CH<sub>2</sub>-, oxazolyl-CH<sub>2</sub>-,  
 isoxazolyl-CH<sub>2</sub>-,  
 phenyl-CH<sub>2</sub>CH<sub>2</sub>-, (2-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (4-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, furanyl-CH<sub>2</sub>CH<sub>2</sub>-, thienyl-CH<sub>2</sub>CH<sub>2</sub>-,  
 35 pyridyl-CH<sub>2</sub>CH<sub>2</sub>-, 1-imidazolyl-CH<sub>2</sub>CH<sub>2</sub>-, oxazolyl-CH<sub>2</sub>CH<sub>2</sub>-,  
 isoxazolyl-CH<sub>2</sub>CH<sub>2</sub>-,

Z is methyl, ethyl, i-propyl, n-propyl, n-butyl, i-butyl, s-butyl, t-butyl, or allyl;

R<sup>10</sup> is H, methyl, ethyl, phenyl, benzyl, phenethyl,

- 5 4-F-phenyl, (4-F-phenyl)CH<sub>2</sub>-, (4-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
4-Cl-phenyl, (4-Cl-phenyl)CH<sub>2</sub>-, (4-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
4-CH<sub>3</sub>-phenyl, (4-CH<sub>3</sub>-phenyl)CH<sub>2</sub>-, (4-CH<sub>3</sub>-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
4-CF<sub>3</sub>-phenyl, (4-CF<sub>3</sub>-phenyl)CH<sub>2</sub>-, or  
(4-CF<sub>3</sub>-phenyl)CH<sub>2</sub>CH<sub>2</sub>-;

10

R<sup>11</sup>, at each occurrence, is independently selected from

- H, =O, methyl, ethyl, phenyl, benzyl, phenethyl,  
4-F-phenyl, (4-F-phenyl)CH<sub>2</sub>-, (4-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
3-F-phenyl, (3-F-phenyl)CH<sub>2</sub>-, (3-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
15 2-F-phenyl, (2-F-phenyl)CH<sub>2</sub>-, (2-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
4-Cl-phenyl, (4-Cl-phenyl)CH<sub>2</sub>-, (4-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
3-Cl-phenyl, (3-Cl-phenyl)CH<sub>2</sub>-, (3-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
4-CH<sub>3</sub>-phenyl, (4-CH<sub>3</sub>-phenyl)CH<sub>2</sub>-, (4-CH<sub>3</sub>-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
3-CH<sub>3</sub>-phenyl, (3-CH<sub>3</sub>-phenyl)CH<sub>2</sub>-, (3-CH<sub>3</sub>-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
20 4-CF<sub>3</sub>-phenyl, (4-CF<sub>3</sub>-phenyl)CH<sub>2</sub>-, (4-CF<sub>3</sub>-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
pyrid-2-yl, pyrid-3-yl, or pyrid-4-yl, and

R<sup>13</sup>, at each occurrence, is independently selected from

H, F, Cl, OH, -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -OCH<sub>3</sub>, or -CF<sub>3</sub>.

25

[11] In another even further more preferred embodiment the present invention provides for a compound selected from:

- (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-  
30 benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-  
butanediamide;

- (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-  
benzodiazepin-3-yl]-2-(2-methylpropyl)-3-propyl-  
35 butanediamide;

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

5 (2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

(2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-propyl-butanediamide;

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-propyl-butanediamide;

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-methyl-3-allyl-butanediamide;

20 (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-methyl-3-allyl-butanediamide;

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-methyl-3-propyl-butanediamide;

25 (2R) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-methyl-butanediamide;

(2R,3S) N1-[1,3-dihydro-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

5 (2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(2-fluorophenyl)-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

10

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(2-fluorophenyl)-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

15

(2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-(2-fluorophenyl)-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

20

(2S,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

25 (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-propyl-butanediamide;

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(2-fluorophenyl)-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-propyl-butanediamide;

30

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(4-fluorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

35

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-fluorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

- 5 (2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-(4-fluorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

- 10 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(pyrid-2-yl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

- (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(N-morpholino)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;
- 15

- (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(dimethylamino)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;
- 20

- (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(N-methyl-N-phenylamino)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;
- 25

- (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(N-piperidinyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

- 30 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(N-homopiperidinyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

- (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(3-methoxyphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;
- 35

- (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(pyrid-4-yl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;
- 5 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-7-methoxy-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;
- (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(pyrid-3-yl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;
- 10 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-(cyclopropylmethyl)-butanediamide;
- 15 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(3-fluorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;
- 20 (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(3-fluorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;
- 25 (2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-(3-fluorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;
- (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-(3-buten-1-yl)-butanediamide;
- 30 (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-(cyclopentylethyl)-butanediamide;
- 35

- (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-(3-buten-1-yl)-butanediamide;
- 5 (2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-(3-buten-1-yl)-butanediamide;
- (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;
- 10 (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;
- 15 (2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;
- 20 (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-n-butyl-butanediamide;
- (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-propyl-butanediamide;
- 25 (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-chlorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-(3-buten-1-yl)-butanediamide;
- 30 (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-chlorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-n-butyl-butanediamide;
- 35

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-N4-[benzyl]-butanediamide;

5 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-methyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-n-butyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

10

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(2-methylpropyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

15

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(4-chlorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

20

(2R,3S) N1-[1,3-dihydro-1-ethyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

25 (2R,3S) N1-[1,3-dihydro-1-propyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

(2R,3S) N1-[1,3-dihydro-1-(isopropyl)-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide;

30

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3,3-diallyl-butanediamide;


35



(2R,3S) N1-[6,7-dihydro-5-methyl-6-oxo-5H-dibenz[b,d]azepin-7-yl]-2-(2-methylpropyl)-3-allylbutanediarnide; and

5 (2R,3S) N1-[1,3,4,5-tetrahydro-1,5-dimethyl-2,4-dioxo-2H-  
1,5-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-  
butanediamide.

[12] In a preferred embodiment the present invention  
10 provides for a compound of Formula (Ia):



15 or a pharmaceutically acceptable salt or prodrug thereof,  
wherein:

Z is C<sub>1</sub>-C<sub>8</sub> alkyl substituted with 1-3 R<sup>12</sup>;

C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 1-3 R<sup>12</sup>;

20 C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 1-3 R<sup>12</sup>;

C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;

C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-4 R<sup>12b</sup>; or

5 to 10 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

25 sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>12b</sup>;

provided, when  $R^{13}$  is H,

then Z is C<sub>4</sub>-C<sub>8</sub> alkyl substituted with 1-3 R<sup>12</sup>;

30 C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 1-3 R<sup>12</sup>; or

C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 1-3 R<sup>12</sup>; and

provided, when ring B is a 1,3,4,5-tetrahydro-1-(Z)-5-

(R<sup>10</sup>)-6,6,7,7-tetra(R<sup>11</sup>)-2,4-dioxo-2H-1,5-diazepin-3-yl

35 core, and  $R^{13}$  is H; then

$R^{10}$  is H,  $C(=O)R^{17}$ ,  $C(=O)OR^{17}$ ,  $C(=O)NR^{18}R^{19}$ ,  
 $S(=O)_2NR^{18}R^{19}$ ,  $S(=O)_2R^{17}$ ; or  
 $C_1-C_6$  alkyl optionally substituted with 0-3  $R^{10a}$ ;

5

$R^{10a}$ , at each occurrence, is independently selected from  
H,  $C_1-C_6$  alkyl,  $OR^{14}$ , Cl, F, Br, I, =O, CN,  $NO_2$ ,  
 $NR^{15}R^{16}$ , and  $CF_3$ .

10 [13] In a more preferred embodiment the present  
invention provides for a compound of Formula (Ia):

$R^3$  is  $-(CR^7R^{7a})_n-R^4$ ,  
 $-(CR^7R^{7a})_n-S-(CR^7R^{7a})_m-R^4$ ,  
15  $-(CR^7R^{7a})_n-O-(CR^7R^{7a})_m-R^4$ , or  
 $-(CR^7R^{7a})_n-N(R^{7b})-(CR^7R^{7a})_m-R^4$ ;

$n$  is 0, 1, or 2;

20  $m$  is 0, 1, or 2;

$R^{3a}$  is H, OH, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, butoxy, allyl, or 3-buten-1-yl;

25  $R^4$  is H, OH,  $OR^{14a}$ ,  
 $C_1-C_6$  alkyl substituted with 0-3  $R^{4a}$ ,  
 $C_2-C_6$  alkenyl substituted with 0-3  $R^{4a}$ ,  
 $C_2-C_6$  alkynyl substituted with 0-3  $R^{4a}$ ,  
 $C_3-C_{10}$  carbocycle substituted with 0-3  $R^{4b}$ ,  
30  $C_6-C_{10}$  aryl substituted with 0-3  $R^{4b}$ , or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3  $R^{4b}$ ;

35

$R^{4a}$ , at each occurrence, is independently selected from is  
H, F, Cl, Br, I,  $CF_3$ ,  
 $C_3-C_{10}$  carbocycle substituted with 0-3  $R^{4b}$ ,

C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>4b</sup>, or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
5 is substituted with 0-3 R<sup>4b</sup>;

R<sup>4b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>,  
S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub>  
10 haloalkyl, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy;

R<sup>5</sup> is H, OR<sup>14</sup>;  
C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 0-3 R<sup>5b</sup>;  
C<sub>1</sub>-C<sub>6</sub> alkoxy substituted with 0-3 R<sup>5b</sup>;  
15 C<sub>2</sub>-C<sub>6</sub> alkenyl substituted with 0-3 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>6</sub> alkynyl substituted with 0-3 R<sup>5b</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>5c</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>5c</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
20 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>5c</sup>;

R<sup>5a</sup> is H or C<sub>1</sub>-C<sub>4</sub> alkyl;

25 R<sup>5b</sup>, at each occurrence, is independently selected from:  
H, C<sub>1</sub>-C<sub>6</sub> alkyl, CF<sub>3</sub>, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>,  
NR<sup>15</sup>R<sup>16</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>5c</sup>;  
30 C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>5c</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>5c</sup>;

35

R<sup>5c</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>,

$S(=O)CH_3$ ,  $S(=O)_2CH_3$ ,  $C_1-C_6$  alkyl,  $C_1-C_4$  alkoxy,  $C_1-C_4$  haloalkyl, and  $C_1-C_4$  haloalkoxy;

$R^6$  is H, methyl, or ethyl;

5

$R^7$ , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN,  $NO_2$ ,  $CF_3$ , phenyl, and  $C_1-C_4$  alkyl;

10  $R^{7a}$ , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN,  $NO_2$ ,  $CF_3$ , and  $C_1-C_4$  alkyl;

$R^{7b}$  is independently selected from H, methyl, ethyl, propyl, and butyl;

15

Ring B is a 7 membered lactam,

wherein the lactam is saturated, partially saturated or unsaturated;

wherein each additional lactam carbon is substituted with 0-2  $R^{11}$ ; and,

20

optionally, the lactam contains a heteroatom selected from -O-, -S-,  $-S(=O)-$ ,  $-S(=O)_2-$ , -N=, -NH-, and -N( $R^{10}$ )-;

25 additionally, two  $R^{11}$  substituents on adjacent atoms may be combined to form a benzo fused radical; wherein said benzo fused radical is substituted with 0-3  $R^{13}$ ;

30 additionally, two  $R^{11}$  substituents on adjacent atoms may be combined to form a 5 to 6 membered heteroaryl fused radical, wherein said 5 to 6 membered heteroaryl fused radical comprises 1 or 2 heteroatoms selected from N, O, and S; wherein said 5 to 6 membered heteroaryl fused radical is substituted with 0-3  $R^{13}$ ;

35

additionally, two  $R^{11}$  substituents on the same or adjacent carbon atoms may be combined to form a  $C_3-C_6$  carbocycle substituted with 0-3  $R^{13}$ ;

- $R^{10}$  is H,  $C(=O)R^{17}$ ,  $C(=O)OR^{17}$ ,  $C(=O)NR^{18}R^{19}$ ,  
 $S(=O)_2NR^{18}R^{19}$ ,  $S(=O)_2R^{17}$ ;  
 $C_1-C_6$  alkyl optionally substituted with 0-2  $R^{10a}$ ;  
5  $C_6-C_{10}$  aryl substituted with 0-4  $R^{10b}$ ;  
 $C_3-C_{10}$  carbocycle substituted with 0-3  $R^{10b}$ ; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
10 is substituted with 0-3  $R^{10b}$ ;
- $R^{10a}$ , at each occurrence, is independently selected from H,  
 $C_1-C_6$  alkyl,  $OR^{14}$ , Cl, F, Br, I, =O, CN,  $NO_2$ ,  $NR^{15}R^{16}$ ,  
CF<sub>3</sub>, or phenyl substituted with 0-4  $R^{10b}$ ;
- 15  $R^{10b}$ , at each occurrence, is independently selected from H,  
OH,  $C_1-C_6$  alkyl,  $C_1-C_4$  alkoxy, Cl, F, Br, I, CN,  $NO_2$ ,  
 $NR^{15}R^{16}$ , or CF<sub>3</sub>;
- 20  $R^{11}$ , at each occurrence, is independently selected from  
H,  $C_1-C_4$  alkoxy, Cl, F, Br, I, =O, CN,  $NO_2$ ,  $NR^{18}R^{19}$ ,  
 $C(=O)R^{17}$ ,  $C(=O)OR^{17}$ ,  $C(=O)NR^{18}R^{19}$ ,  $S(=O)_2NR^{18}R^{19}$ , CF<sub>3</sub>;  
 $C_1-C_6$  alkyl optionally substituted with 0-3  $R^{11a}$ ;  
 $C_6-C_{10}$  aryl substituted with 0-3  $R^{11b}$ ;  
25  $C_3-C_{10}$  carbocycle substituted with 0-3  $R^{11b}$ ; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3  $R^{11b}$ ;
- 30  $R^{11a}$ , at each occurrence, is independently selected from H,  
 $C_1-C_6$  alkyl,  $OR^{14}$ , Cl, F, Br, I, =O, CN,  $NO_2$ ,  $NR^{15}R^{16}$ ,  
CF<sub>3</sub>, or phenyl substituted with 0-3  $R^{11b}$ ;
- 35  $R^{11b}$ , at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN,  $NO_2$ ,  $NR^{15}R^{16}$ , CF<sub>3</sub>, acetyl, SCH<sub>3</sub>,  
 $S(=O)CH_3$ ,  $S(=O)_2CH_3$ ,  $C_1-C_6$  alkyl,  $C_1-C_4$  alkoxy,  $C_1-C_4$   
haloalkyl, and  $C_1-C_4$  haloalkoxy;

Z is C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 1-3 R<sup>12</sup>;  
5 C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-4 R<sup>12b</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
10 is substituted with 0-3 R<sup>12b</sup>;

R<sup>12</sup>, at each occurrence, is independently selected from  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-4 R<sup>12b</sup>; or  
15 5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>12b</sup>;

20 R<sup>12b</sup>, at each occurrence, is independently selected from  
H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl,  
SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy,  
C<sub>1</sub>-C<sub>4</sub> haloalkyl, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy;

25 R<sup>13</sup>, at each occurrence, is independently selected from  
H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, CN,  
NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;

R<sup>14</sup> is H, phenyl, benzyl, C<sub>1</sub>-C<sub>6</sub> alkyl, or C<sub>2</sub>-C<sub>6</sub> alkoxyalkyl;  
30

R<sup>14a</sup> is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;

R<sup>15</sup>, at each occurrence, is independently selected from H,  
C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-,  
35 and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-;

R<sup>16</sup>, at each occurrence, is independently selected from  
H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, phenethyl,

(C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-;

R<sup>17</sup> is H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkoxyalkyl,  
aryl substituted by 0-4 R<sup>17a</sup>, or  
5 -CH<sub>2</sub>-aryl substituted by 0-4 R<sup>17a</sup>;

R<sup>17a</sup> is H, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
propoxy, butoxy, -OH, F, Cl, Br, I, CF<sub>3</sub>, OCF<sub>3</sub>, SCH<sub>3</sub>,  
S(O)CH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, -NH<sub>2</sub>, -N(CH<sub>3</sub>)<sub>2</sub>, or C<sub>1</sub>-C<sub>4</sub> haloalkyl;  
10

R<sup>18</sup>, at each occurrence, is independently selected from  
H, C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl, benzyl, phenethyl,  
(C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-; and

15 R<sup>19</sup>, at each occurrence, is independently selected from  
H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl, benzyl, phenethyl,  
(C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-;

provided, when R<sup>13</sup> is H,  
20 then Z is C<sub>4</sub>-C<sub>6</sub> alkyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 1-3 R<sup>12</sup>; or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 1-3 R<sup>12</sup>.

[14] In a further more preferred embodiment the  
25 present invention provides for a compound of Formula (Ia)  
wherein:

R<sup>3</sup> is -(CHR<sup>7</sup>)<sub>n</sub>-R<sup>4</sup>,

30 n is 0 or 1;

R<sup>3a</sup> is H, OH, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, butoxy, allyl, or 3-buten-1-yl;

35 R<sup>4</sup> is H, OH, OR<sup>14a</sup>,  
C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-2 R<sup>4a</sup>,  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-2 R<sup>4a</sup>,  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>4a</sup>,

C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>4b</sup>, or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
5 sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>4b</sup>;

R<sup>4a</sup>, at each occurrence, is independently selected from is  
H, F, Cl, Br, I, CF<sub>3</sub>,  
10 C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,  
phenyl substituted with 0-3 R<sup>4b</sup>, or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
15 substituted with 0-3 R<sup>4b</sup>;

R<sup>4b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>,  
S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkoxy, C<sub>1</sub>-C<sub>2</sub>  
20 haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

R<sup>5</sup> is H, OR<sup>14</sup>;  
C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-3 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>5b</sup>;  
25 C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>5b</sup>;

R<sup>5a</sup> is H, methyl, ethyl, propyl, or butyl;

R<sup>5b</sup>, at each occurrence, is independently selected from:  
30 H, methyl, ethyl, propyl, butyl, CF<sub>3</sub>, OR<sup>14</sup>, Cl, F, Br,  
I, =O;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>5c</sup>;  
phenyl substituted with 0-3 R<sup>5c</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
35 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>5c</sup>;



R<sup>5c</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

5

R<sup>6</sup> is H;

R<sup>7</sup>, at each occurrence, is independently selected from H, F, CF<sub>3</sub>, methyl, and ethyl;

10

Ring B is a 7 membered lactam, wherein the lactam is saturated, partially saturated or unsaturated; wherein each additional lactam carbon is substituted with 0-2 R<sup>11</sup>; and, optionally, the lactam contains a heteroatom selected from -N=, -NH-, and -N(R<sup>10</sup>)-;

15

additionally, two R<sup>11</sup> substituents on adjacent atoms may be combined to form a benzo fused radical; wherein said benzo fused radical is substituted with 0-2 R<sup>13</sup>;

20

additionally, two R<sup>11</sup> substituents on adjacent atoms may be combined to form a 5 to 6 membered heteroaryl fused radical, wherein said 5 to 6 membered heteroaryl fused radical comprises 1 or 2 heteroatoms selected from N, O, and S; wherein said 5 to 6 membered heteroaryl fused radical is substituted with 0-2 R<sup>13</sup>;

25

additionally, two R<sup>11</sup> substituents on the same or adjacent carbon atoms may be combined to form a C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-2 R<sup>13</sup>;

30

R<sup>10</sup> is H, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>; C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with 0-1 R<sup>10a</sup>; phenyl substituted with 0-4 R<sup>10b</sup>; C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>10b</sup>; or

35

- 5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R<sup>10b</sup>;
- 5 R<sup>10a</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>4</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, or phenyl substituted with 0-4 R<sup>10b</sup>;
- 10 R<sup>10b</sup>, at each occurrence, is independently selected from H, OH, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkoxy, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, or CF<sub>3</sub>;
- 15 R<sup>11</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, =O, NR<sup>18</sup>R<sup>19</sup>, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>, CF<sub>3</sub>;  
C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 0-3 R<sup>11a</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>11b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>11b</sup>; or
- 20 5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R<sup>11b</sup>;
- 25 R<sup>11a</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>4</sub> alkyl, OR<sup>14</sup>, F, =O, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, or phenyl substituted with 0-3 R<sup>11b</sup>;
- 30 R<sup>11b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;
- Z is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 1-3 R<sup>12</sup>;
- 35 C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 1-3 R<sup>12</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-4 R<sup>12b</sup>; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R<sup>12b</sup>;

5

R<sup>12</sup>, at each occurrence, is independently selected from C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;

C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-4 R<sup>12b</sup>; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R<sup>12b</sup>;

10

R<sup>12b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

15

R<sup>13</sup>, at each occurrence, is independently selected from H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;

20

R<sup>14</sup> is H, phenyl, benzyl, C<sub>1</sub>-C<sub>4</sub> alkyl, or C<sub>2</sub>-C<sub>4</sub> alkoxyalkyl;

R<sup>15</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>4</sub> alkyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>4</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>4</sub> alkyl)-S(=O)<sub>2</sub>-;

25

R<sup>16</sup>, at each occurrence, is independently selected from H, OH, C<sub>1</sub>-C<sub>4</sub> alkyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>4</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>4</sub> alkyl)-S(=O)<sub>2</sub>-;

30

R<sup>17</sup> is H, methyl, ethyl, propyl, butyl, methoxymethyl, ethoxymethyl, methoxyethyl, ethoxyethyl, phenyl substituted by 0-3 R<sup>17a</sup>, or -CH<sub>2</sub>-phenyl substituted by 0-3 R<sup>17a</sup>;

35

R<sup>17a</sup> is H, methyl, methoxy, -OH, F, Cl, CF<sub>3</sub>, or OCF<sub>3</sub>;

R<sup>18</sup>, at each occurrence, is independently selected from  
H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and  
phenethyl; and


5

R<sup>19</sup>, at each occurrence, is independently selected from  
H, methyl, and ethyl;

provided, when R<sup>13</sup> is H,

- 10 then Z is butyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 1-3 R<sup>12</sup>; or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 1-3 R<sup>12</sup>.


[15] In an even more preferred embodiment the present  
15 invention provides for a compound of Formula (Ib):




(Ib)

or a pharmaceutically acceptable salt or prodrug thereof  
20 wherein:

Ring B is selected from:




25



[16] In an even further more preferred embodiment the present invention provides for a compound of Formula (Ic):

5



(Ic)

or a pharmaceutically acceptable salt or prodrug thereof wherein

10

$R^3$  is  $R^4$ ,

$R^4$  is  $C_1$ - $C_4$  alkyl substituted with 0-1  $R^{4a}$ ,  
 $C_2$ - $C_4$  alkenyl substituted with 0-1  $R^{4a}$ , or  
 $C_2$ - $C_4$  alkynyl substituted with 0-1  $R^{4a}$ ;

15

$R^{4a}$ , at each occurrence, is independently selected from  
H, F,  $CF_3$ ,

20

$C_3$ - $C_6$  carbocycle substituted with 0-3  $R^{4b}$ ,

phenyl substituted with 0-3  $R^{4b}$ , or

5 to 6 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3  $R^{4b}$ ; wherein said 5 to 6

25

membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>4b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

5

R<sup>5</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>5b</sup>;

10 R<sup>5b</sup>, at each occurrence, is independently selected from:  
H, methyl, ethyl, propyl, butyl, CF<sub>3</sub>, OR<sup>14</sup>, =O;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-2 R<sup>5c</sup>;  
phenyl substituted with 0-3 R<sup>5c</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
15 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>5c</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
20 pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>5c</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
25 S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

R<sup>11</sup>, at each occurrence, is independently selected from  
H, =O, NR<sup>18</sup>R<sup>19</sup>, CF<sub>3</sub>;  
30 C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with 0-1 R<sup>11a</sup>;  
phenyl substituted with 0-3 R<sup>11b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>11b</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
35 sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>11b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,

pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

5 R<sup>11a</sup>, at each occurrence, is independently selected from H,  
C<sub>1</sub>-C<sub>4</sub> alkyl, OR<sup>14</sup>, F, Cl, =O, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, or phenyl  
substituted with 0-3 R<sup>11b</sup>;

10 R<sup>11b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, methyl, ethyl, propyl, butyl,  
methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub>  
haloalkoxy;

15 Z is C<sub>1</sub>-C<sub>3</sub> alkyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>3</sub> alkenyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>3</sub> alkynyl substituted with 1-3 R<sup>12</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>12b</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
20 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>12b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
25 imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>12</sup>, at each occurrence, is independently selected from  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>12b</sup>; or  
30 5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>12b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
35 pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>12b</sup>, at each occurrence, is independently selected from  
H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

5

R<sup>13</sup>, at each occurrence, is independently selected from  
H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
Cl, F, Br, CN, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;

10 R<sup>14</sup> is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;

R<sup>15</sup>, at each occurrence, is independently selected from H,  
methyl, ethyl, propyl, and butyl;

15 R<sup>16</sup>, at each occurrence, is independently selected from  
H, OH, methyl, ethyl, propyl, butyl, benzyl,  
phenethyl, methyl-C(=O)-, ethyl-C(=O)-,  
methyl-S(=O)<sub>2</sub>-, and ethyl-S(=O)<sub>2</sub>-;

20 R<sup>18</sup>, at each occurrence, is independently selected from  
H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and  
phenethyl; and

R<sup>19</sup>, at each occurrence, is independently selected from  
25 H, methyl, and ethyl.

provided, when R<sup>13</sup> is H,


then Z is C<sub>2</sub>-C<sub>3</sub> alkenyl substituted with 1-3 R<sup>12</sup>; or

C<sub>2</sub>-C<sub>3</sub> alkynyl substituted with 1-3 R<sup>12</sup>.

30

[17] In another even further more preferred embodiment  
the present invention provides for a compound of Formula  
(Id):





(Id)

or a pharmaceutically acceptable salt or prodrug thereof  
wherein:

5

$R^3$  is  $R^4$ ,

$R^4$  is  $C_1$ - $C_4$  alkyl substituted with 0-1  $R^{4a}$ ,  
 $C_2$ - $C_4$  alkenyl substituted with 0-1  $R^{4a}$ , or  
10  $C_2$ - $C_4$  alkynyl substituted with 0-1  $R^{4a}$ ;

$R^{4a}$ , at each occurrence, is independently selected from  
H, F,  $CF_3$ ,  
 $C_3$ - $C_6$  carbocycle substituted with 0-3  $R^{4b}$ ,  
15 phenyl substituted with 0-3  $R^{4b}$ , or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3  $R^{4b}$ ; wherein said 5 to 6  
20 membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

25  $R^{4b}$ , at each occurrence, is independently selected from H,  
OH, Cl, F,  $NR^{15}R^{16}$ ,  $CF_3$ , acetyl,  $SCH_3$ ,  $S(=O)CH_3$ ,  
 $S(=O)_2CH_3$ , methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy,  $C_1$ - $C_2$  haloalkyl, and  $C_1$ - $C_2$  haloalkoxy;

30  $R^5$  is  $C_1$ - $C_4$  alkyl substituted with 0-1  $R^{5b}$ ;  
 $C_2$ - $C_4$  alkenyl substituted with 0-1  $R^{5b}$ ;  
 $C_2$ - $C_4$  alkynyl substituted with 0-1  $R^{5b}$ ;

R<sup>5b</sup>, at each occurrence, is independently selected from:

H, methyl, ethyl, propyl, butyl, CF<sub>3</sub>, OR<sup>14</sup>, =O;

C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-2 R<sup>5c</sup>;

phenyl substituted with 0-3 R<sup>5c</sup>; or

- 5 5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>5c</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
10 pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>5c</sup>, at each occurrence, is independently selected from H,

- 15 OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

R<sup>11</sup>, at each occurrence, is independently selected from

- 20 H, =O, NR<sup>18</sup>R<sup>19</sup>, CF<sub>3</sub>;  
C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with 0-1 R<sup>11a</sup>;  
phenyl substituted with 0-3 R<sup>11b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>11b</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
25 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>11b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
30 pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>11a</sup>, at each occurrence, is independently selected from H,

- 35 C<sub>1</sub>-C<sub>4</sub> alkyl, OR<sup>14</sup>, F, Cl, =O, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, or phenyl  
substituted with 0-3 R<sup>11b</sup>;

R<sup>11b</sup>, at each occurrence, is independently selected from H,

OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, methyl, ethyl, propyl, butyl,

methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

Z is C<sub>1</sub>-C<sub>3</sub> alkyl substituted with 1-3 R<sup>12</sup>;

5 C<sub>2</sub>-C<sub>3</sub> alkenyl substituted with 1-3 R<sup>12</sup>;

C<sub>2</sub>-C<sub>3</sub> alkynyl substituted with 1-3 R<sup>12</sup>;

C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;

C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>12b</sup>; or

5 to 6 membered heterocycle containing 1 to 4

10 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R<sup>12b</sup>; wherein said 5 to 6 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, 15 pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>12</sup>, at each occurrence, is independently selected from C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;

20 C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>12b</sup>; or

5 to 6 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R<sup>12b</sup>; wherein said 5 to 6 25 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

30 R<sup>12b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

35 R<sup>13</sup>, at each occurrence, is independently selected from H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy, Cl, F, Br, CN, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;

R<sup>14</sup> is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;

R<sup>15</sup>, at each occurrence, is independently selected from H, methyl, ethyl, propyl, and butyl;

5

R<sup>16</sup>, at each occurrence, is independently selected from H, OH, methyl, ethyl, propyl, butyl, benzyl, phenethyl, methyl-C(=O)-, ethyl-C(=O)-, methyl-S(=O)<sub>2</sub>-, and ethyl-S(=O)<sub>2</sub>-;

10

R<sup>18</sup>, at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl; and

15 R<sup>19</sup>, at each occurrence, is independently selected from H, methyl, and ethyl.

provided, when R<sup>13</sup> is H,


then Z is C<sub>2</sub>-C<sub>3</sub> alkenyl substituted with 1-3 R<sup>12</sup>; or

20

C<sub>2</sub>-C<sub>3</sub> alkynyl substituted with 1-3 R<sup>12</sup>.

[18] In another even further more preferred embodiment the present invention provides for a compound of Formula (Ie):

25



(Ie)

or a pharmaceutically acceptable salt or prodrug thereof wherein:

30

R<sup>3</sup> is R<sup>4</sup>,

R<sup>4</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>4a</sup>,

C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>4a</sup>, or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>4a</sup>;

R<sup>4a</sup>, at each occurrence, is independently selected from  
5 H, F, CF<sub>3</sub>,  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,

---

R<sup>5c</sup>, at each occurrence, is independently selected from H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

5

R<sup>10</sup> is H, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>;

C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with 0-1 R<sup>10a</sup>;

phenyl substituted with 0-4 R<sup>10b</sup>;

C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>10b</sup>; or

10

5 to 6 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 6 membered heterocycle is

substituted with 0-3 R<sup>10b</sup>; wherein said 5 to 6

membered heterocycle is selected from pyridinyl,

15

pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,

pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,

imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

20

R<sup>10a</sup>, at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, OR<sup>14</sup>, Cl, F, =O, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, or phenyl substituted with 0-4 R<sup>10b</sup>;

25

R<sup>10b</sup>, at each occurrence, is independently selected from H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, Cl, F, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;

30

Z is C<sub>1</sub>-C<sub>3</sub> alkyl substituted with 1-3 R<sup>12</sup>;

C<sub>2</sub>-C<sub>3</sub> alkenyl substituted with 1-3 R<sup>12</sup>;

C<sub>2</sub>-C<sub>3</sub> alkynyl substituted with 1-3 R<sup>12</sup>;

C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;

C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>12b</sup>; or

5 to 6 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 6 membered heterocycle is

35

substituted with 0-3 R<sup>12b</sup>; wherein said 5 to 6

membered heterocycle is selected from pyridinyl,

pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,

pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

- 5  $R^{12}$ , at each occurrence, is independently selected from  
 $C_6-C_{10}$  aryl substituted with 0-4  $R^{12b}$ ;  
 $C_3-C_6$  carbocycle substituted with 0-3  $R^{12b}$ ; or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
10 substituted with 0-3  $R^{12b}$ ; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;  
15
- $R^{12b}$ , at each occurrence, is independently selected from  
H, OH, Cl, F,  $NR^{15}R^{16}$ ,  $CF_3$ , acetyl,  $SCH_3$ ,  $S(=O)CH_3$ ,  
 $S(=O)_2CH_3$ , methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy,  $C_1-C_2$  haloalkyl, and  $C_1-C_2$  haloalkoxy;  
20
- $R^{13}$ , at each occurrence, is independently selected from  
H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
Cl, F, Br, CN,  $NR^{15}R^{16}$ , and  $CF_3$ ;
- 25  $R^{14}$  is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;
- $R^{15}$ , at each occurrence, is independently selected from H,  
methyl, ethyl, propyl, and butyl;
- 30  $R^{16}$ , at each occurrence, is independently selected from  
H, OH, methyl, ethyl, propyl, butyl, benzyl,  
phenethyl, methyl-C(=O)-, ethyl-C(=O)-,  
methyl-S(=O)<sub>2</sub>-, and ethyl-S(=O)<sub>2</sub>-;
- 35  $R^{17}$  is H, methyl, ethyl, propyl, butyl, methoxymethyl,  
ethoxymethyl, methoxyethyl, ethoxyethyl,  
phenyl substituted by 0-3  $R^{17a}$ , or  
-CH<sub>2</sub>-phenyl substituted by 0-3  $R^{17a}$ ;

R<sup>17a</sup> is H, methyl, methoxy, -OH, F, Cl, CF<sub>3</sub>, or OCF<sub>3</sub>;


5 R<sup>18</sup>, at each occurrence, is independently selected from  
H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and  
phenethyl; and

R<sup>19</sup>, at each occurrence, is independently selected from  
H, methyl, and ethyl.

10

provided, when R<sup>13</sup> is H,  
then Z is C<sub>2</sub>-C<sub>3</sub> alkenyl substituted with 1-3 R<sup>12</sup>; or  
C<sub>2</sub>-C<sub>3</sub> alkynyl substituted with 1-3 R<sup>12</sup>.

15 [19] In another even further more preferred embodiment  
the present invention provides for a compound of Formula  
(If):



20

(If)

or a pharmaceutically acceptable salt or prodrug thereof  
wherein:

R<sup>3</sup> is R<sup>4</sup>,

25

R<sup>4</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>4a</sup>,  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>4a</sup>, or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>4a</sup>;

30 R<sup>4a</sup>, at each occurrence, is independently selected from  
H, F, CF<sub>3</sub>,  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,  
phenyl substituted with 0-3 R<sup>4b</sup>, or



- 5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R<sup>4b</sup>; wherein said 5 to 6 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;
- 10 R<sup>4b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;
- 15 R<sup>5</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>5b</sup>; C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>5b</sup>; C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>5b</sup>;
- R<sup>5b</sup>, at each occurrence, is independently selected from:
- 20 H, methyl, ethyl, propyl, butyl, CF<sub>3</sub>, OR<sup>14</sup>, =O; C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-2 R<sup>5c</sup>; phenyl substituted with 0-3 R<sup>5c</sup>; or 5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R<sup>5c</sup>; wherein said 5 to 6 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;
- 25 30 R<sup>5c</sup>, at each occurrence, is independently selected from H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;
- 35 Z is C<sub>1</sub>-C<sub>3</sub> alkyl substituted with 1-3 R<sup>12</sup>; C<sub>2</sub>-C<sub>3</sub> alkenyl substituted with 1-3 R<sup>12</sup>;

- C<sub>2</sub>-C<sub>3</sub> alkynyl substituted with 1-3 R<sup>12</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>12b</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
5 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>12b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
10 pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;
- R<sup>12</sup>, at each occurrence, is independently selected from  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
15 C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>12b</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>12b</sup>; wherein said 5 to 6  
20 membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;
- 25 R<sup>12b</sup>, at each occurrence, is independently selected from  
H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;
- 30 R<sup>13</sup>, at each occurrence, is independently selected from  
H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
Cl, F, Br, CN, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;
- R<sup>14</sup> is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;  
35
- R<sup>15</sup>, at each occurrence, is independently selected from H,  
methyl, ethyl, propyl, and butyl;

R<sup>16</sup>, at each occurrence, is independently selected from  
H, OH, methyl, ethyl, propyl, butyl, benzyl,  
phenethyl, methyl-C(=O)-, ethyl-C(=O)-,  
methyl-S(=O)<sub>2</sub>-, and ethyl-S(=O)<sub>2</sub>-;

5

R<sup>18</sup>, at each occurrence, is independently selected from  
H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and  
phenethyl; and

10 R<sup>19</sup>, at each occurrence, is independently selected from  
H, methyl, and ethyl.

provided, when R<sup>13</sup> is H,

then Z is C<sub>2</sub>-C<sub>3</sub> alkenyl substituted with 1-3 R<sup>12</sup>; or

15 C<sub>2</sub>-C<sub>3</sub> alkynyl substituted with 1-3 R<sup>12</sup>.

[20] In another even further more preferred embodiment  
the present invention provides for a compound of one of  
Formulas (Ic), (Id), (Ie), or (If), wherein:

20

R<sup>3</sup> is -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>,  
-CH<sub>2</sub>(CH<sub>3</sub>)<sub>2</sub>, -CH(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>C(CH<sub>3</sub>)<sub>3</sub>,  
-CF<sub>3</sub>, -CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CF<sub>3</sub>,  
-CH=CH<sub>2</sub>, -CH<sub>2</sub>CH=CH<sub>2</sub>, -CH<sub>2</sub>C(CH<sub>3</sub>)=CH<sub>2</sub>, -CH<sub>2</sub>CH=C(CH<sub>3</sub>)<sub>2</sub>,  
25 -CH<sub>2</sub>CH<sub>2</sub>CH=CH<sub>2</sub>, -CH<sub>2</sub>CH<sub>2</sub>C(CH<sub>3</sub>)=CH<sub>2</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH=C(CH<sub>3</sub>)<sub>2</sub>,  
cis-CH<sub>2</sub>CH=CH(CH<sub>3</sub>), cis-CH<sub>2</sub>CH<sub>2</sub>CH=CH(CH<sub>3</sub>),  
trans-CH<sub>2</sub>CH=CH(CH<sub>3</sub>), trans-CH<sub>2</sub>CH<sub>2</sub>CH=CH(CH<sub>3</sub>);  
-C≡CH, -CH<sub>2</sub>C≡CH, -CH<sub>2</sub>C≡C(CH<sub>3</sub>),  
cyclopropyl-CH<sub>2</sub>-, cyclobutyl-CH<sub>2</sub>-, cyclopentyl-CH<sub>2</sub>-,  
30 cyclohexyl-CH<sub>2</sub>-, cyclopropyl-CH<sub>2</sub>CH<sub>2</sub>-,  
cyclobutyl-CH<sub>2</sub>CH<sub>2</sub>-, cyclopentyl-CH<sub>2</sub>CH<sub>2</sub>-,  
cyclohexyl-CH<sub>2</sub>CH<sub>2</sub>-, phenyl-CH<sub>2</sub>-,  
(2-F-phenyl)CH<sub>2</sub>-, (3-F-phenyl)CH<sub>2</sub>-, (4-F-phenyl)CH<sub>2</sub>-,  
(2-Cl-phenyl)CH<sub>2</sub>-, (3-Cl-phenyl)CH<sub>2</sub>-, (4-Cl-phenyl)CH<sub>2</sub>-,  
35 (2,3-diF-phenyl)CH<sub>2</sub>-, (2,4-diF-phenyl)CH<sub>2</sub>-,  
(2,5-diF-phenyl)CH<sub>2</sub>-, (2,6-diF-phenyl)CH<sub>2</sub>-,  
(3,4-diF-phenyl)CH<sub>2</sub>-, (3,5-diF-phenyl)CH<sub>2</sub>-,  
(2,3-diCl-phenyl)CH<sub>2</sub>-, (2,4-diCl-phenyl)CH<sub>2</sub>-,

- (2,5-diCl-phenyl)CH<sub>2</sub>-, (2,6-diCl-phenyl)CH<sub>2</sub>-,  
 (3,4-diCl-phenyl)CH<sub>2</sub>-, (3,5-diCl-phenyl)CH<sub>2</sub>-,  
 (3-F-4-Cl-phenyl)CH<sub>2</sub>-, (3-F-5-Cl-phenyl)CH<sub>2</sub>-,  
 (3-Cl-4-F-phenyl)CH<sub>2</sub>-, phenyl-CH<sub>2</sub>CH<sub>2</sub>-,  
 5 (2-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (4-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (3-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (4-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2,3-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2,4-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2,5-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2,6-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 10 (3,4-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3,5-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2,3-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2,4-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2,5-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2,6-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (3,4-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3,5-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (3-F-4-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, or (3-F-5-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 15  
 R<sup>5</sup> is -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>,  
 -CH(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH<sub>2</sub>C(CH<sub>3</sub>)<sub>3</sub>,  
 -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -CH(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, -CH<sub>2</sub>CH(CH<sub>3</sub>)CH<sub>2</sub>CH<sub>3</sub>,  
 -CH<sub>2</sub>CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, -CH(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>, -CF<sub>3</sub>, -CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CF<sub>3</sub>,  
 20 -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CF<sub>3</sub>, -CH=CH<sub>2</sub>, -CH<sub>2</sub>CH=CH<sub>2</sub>,  
 -CH=CHCH<sub>3</sub>, cis-CH<sub>2</sub>CH=CH(CH<sub>3</sub>), trans-CH<sub>2</sub>CH=CH(CH<sub>3</sub>),  
 trans-CH<sub>2</sub>CH=CH(C<sub>6</sub>H<sub>5</sub>), -CH<sub>2</sub>CH=C(CH<sub>3</sub>)<sub>2</sub>, cis-CH<sub>2</sub>CH=CHCH<sub>2</sub>CH<sub>3</sub>,  
 trans-CH<sub>2</sub>CH=CHCH<sub>2</sub>CH<sub>3</sub>, cis-CH<sub>2</sub>CH<sub>2</sub>CH=CH(CH<sub>3</sub>),  
 trans-CH<sub>2</sub>CH<sub>2</sub>CH=CH(CH<sub>3</sub>), trans-CH<sub>2</sub>CH=CHCH<sub>2</sub>(C<sub>6</sub>H<sub>5</sub>),  
 25 -C≡CH, -CH<sub>2</sub>C≡CH, -CH<sub>2</sub>C≡C(CH<sub>3</sub>), -CH<sub>2</sub>C≡C(C<sub>6</sub>H<sub>5</sub>)  
 -CH<sub>2</sub>CH<sub>2</sub>C≡CH, -CH<sub>2</sub>CH<sub>2</sub>C≡C(CH<sub>3</sub>), -CH<sub>2</sub>CH<sub>2</sub>C≡C(C<sub>6</sub>H<sub>5</sub>)  
 -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C≡CH, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C≡C(CH<sub>3</sub>), -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C≡C(C<sub>6</sub>H<sub>5</sub>)  
 cyclopropyl-CH<sub>2</sub>-, cyclobutyl-CH<sub>2</sub>-, cyclopentyl-CH<sub>2</sub>-,  
 cyclohexyl-CH<sub>2</sub>-, (2-CH<sub>3</sub>-cyclopropyl)CH<sub>2</sub>-,  
 30 (3-CH<sub>3</sub>-cyclobutyl)CH<sub>2</sub>-,  
 cyclopropyl-CH<sub>2</sub>CH<sub>2</sub>-, cyclobutyl-CH<sub>2</sub>CH<sub>2</sub>-,  
 cyclopentyl-CH<sub>2</sub>CH<sub>2</sub>-, cyclohexyl-CH<sub>2</sub>CH<sub>2</sub>-,  
 (2-CH<sub>3</sub>-cyclopropyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-CH<sub>3</sub>-cyclobutyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 phenyl-CH<sub>2</sub>-, (2-F-phenyl)CH<sub>2</sub>-, (3-F-phenyl)CH<sub>2</sub>-,  
 35 (4-F-phenyl)CH<sub>2</sub>-, furanyl-CH<sub>2</sub>-, thienyl-CH<sub>2</sub>-,  
 pyridyl-CH<sub>2</sub>-, 1-imidazolyl-CH<sub>2</sub>-, oxazolyl-CH<sub>2</sub>-,  
 isoxazolyl-CH<sub>2</sub>-,  
 phenyl-CH<sub>2</sub>CH<sub>2</sub>-, (2-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,

(4-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, furanyl-CH<sub>2</sub>CH<sub>2</sub>-, thienyl-CH<sub>2</sub>CH<sub>2</sub>-,  
 pyridyl-CH<sub>2</sub>CH<sub>2</sub>-, 1-imidazolyl-CH<sub>2</sub>CH<sub>2</sub>-, oxazolyl-CH<sub>2</sub>CH<sub>2</sub>-,  
 isoxazolyl-CH<sub>2</sub>CH<sub>2</sub>-,

- 5 Z is phenyl, 2-F-phenyl, 3-F-phenyl, 4-F-phenyl,  
 2-Cl-phenyl, 3-Cl-phenyl, 4-Cl-phenyl, 2,3-diF-phenyl,  
 2,4-diF-phenyl, 2,5-diF-phenyl, 2,6-diF-phenyl,  
 3,4-diF-phenyl, 3,5-diF-phenyl, 2,3-diCl-phenyl,  
 2,4-diCl-phenyl, 2,5-diCl-phenyl, 2,6-diCl-phenyl,  
 10 3,4-diCl-phenyl, 3,5-diCl-phenyl, 3-F-4-Cl-phenyl,  
 3-F-5-Cl-phenyl, 3-Cl-4-F-phenyl, 2-MeO-phenyl,  
 3-MeO-phenyl, 4-MeO-phenyl, 2-Me-phenyl, 3-Me-phenyl,  
 4-Me-phenyl, 2-MeS-phenyl, 3-MeS-phenyl, 4-MeS-phenyl,  
 2-CF<sub>3</sub>O-phenyl, 3-CF<sub>3</sub>O-phenyl, 4-CF<sub>3</sub>O-phenyl,  
 15 furanyl, thienyl, pyridyl, 2-Me-pyridyl, 3-Me-pyridyl,  
 4-Me-pyridyl, 1-imidazolyl, oxazolyl, isoxazolyl,  
 1-benzimidazolyl,  
 cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl,  
 morpholino, N-piperinyl,  
 20 phenyl-CH<sub>2</sub>-, (2-F-phenyl)CH<sub>2</sub>-, (3-F-phenyl)CH<sub>2</sub>-,  
 (4-F-phenyl)CH<sub>2</sub>-, (2-Cl-phenyl)CH<sub>2</sub>-, (3-Cl-phenyl)CH<sub>2</sub>-,  
 (4-Cl-phenyl)CH<sub>2</sub>-, (2,3-diF-phenyl)CH<sub>2</sub>-,  
 (2,4-diF-phenyl)CH<sub>2</sub>-, (2,5-diF-phenyl)CH<sub>2</sub>-,  
 (2,6-diF-phenyl)CH<sub>2</sub>-, (3,4-diF-phenyl)CH<sub>2</sub>-,  
 25 (3,5-diF-phenyl)CH<sub>2</sub>-, (2,3-diCl-phenyl)CH<sub>2</sub>-,  
 (2,4-diCl-phenyl)CH<sub>2</sub>-, (2,5-diCl-phenyl)CH<sub>2</sub>-,  
 (2,6-diCl-phenyl)CH<sub>2</sub>-, (3,4-diCl-phenyl)CH<sub>2</sub>-,  
 (3,5-diCl-phenyl)CH<sub>2</sub>-, (3-F-4-Cl-phenyl)CH<sub>2</sub>-,  
 (3-F-5-Cl-phenyl)CH<sub>2</sub>-, (3-Cl-4-F-phenyl)CH<sub>2</sub>-,  
 30 (2-MeO-phenyl)CH<sub>2</sub>-, (3-MeO-phenyl)CH<sub>2</sub>-,  
 (4-MeO-phenyl)CH<sub>2</sub>-, (2-Me-phenyl)CH<sub>2</sub>-,  
 (3-Me-phenyl)CH<sub>2</sub>-, (4-Me-phenyl)CH<sub>2</sub>-,  
 (2-MeS-phenyl)CH<sub>2</sub>-, (3-MeS-phenyl)CH<sub>2</sub>-,  
 4-MeS-phenyl)CH<sub>2</sub>-, (2-CF<sub>3</sub>O-phenyl)CH<sub>2</sub>-,  
 35 (3-CF<sub>3</sub>O-phenyl)CH<sub>2</sub>-, (4-CF<sub>3</sub>O-phenyl)CH<sub>2</sub>-,  
 (furanyl)CH<sub>2</sub>-, (thienyl)CH<sub>2</sub>-, (pyridyl)CH<sub>2</sub>-,  
 (2-Me-pyridyl)CH<sub>2</sub>-, (3-Me-pyridyl)CH<sub>2</sub>-,  
 (4-Me-pyridyl)CH<sub>2</sub>-, (1-imidazolyl)CH<sub>2</sub>-,


- (oxazolyl)CH<sub>2</sub>-, (isoxazolyl)CH<sub>2</sub>-,  
 (1-benzimidazolyl)CH<sub>2</sub>-, (cyclopropyl)CH<sub>2</sub>-,  
 (cyclobutyl)CH<sub>2</sub>-, (cyclopentyl)CH<sub>2</sub>-,  
 (cyclohexyl)CH<sub>2</sub>-, (morpholino)CH<sub>2</sub>-, (N-pipridinyl)CH<sub>2</sub>-,  
 5 phenyl-CH<sub>2</sub>CH<sub>2</sub>-, (phenyl)<sub>2</sub>CHCH<sub>2</sub>-, (2-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (3-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (4-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (4-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2,3-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 10 (2,4-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2,5-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2,6-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3,4-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (3,5-diF-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2,3-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2,4-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2,5-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2,6-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3,4-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 15 (3,5-diCl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-F-4-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (3-F-5-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-Cl-4-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2-MeO-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-MeO-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (4-MeO-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2-Me-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (3-Me-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (4-Me-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 20 (2-MeS-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-MeS-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (4-MeS-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (2-CF<sub>3</sub>O-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (3-CF<sub>3</sub>O-phenyl)CH<sub>2</sub>CH<sub>2</sub>-, (4-CF<sub>3</sub>O-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (furanyl)CH<sub>2</sub>CH<sub>2</sub>-, (thienyl)CH<sub>2</sub>CH<sub>2</sub>-, (pyridyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (2-Me-pyridyl)CH<sub>2</sub>CH<sub>2</sub>-, (3-Me-pyridyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 25 (4-Me-pyridyl)CH<sub>2</sub>CH<sub>2</sub>-, (imidazolyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (oxazolyl)CH<sub>2</sub>CH<sub>2</sub>-, (isoxazolyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (benzimidazolyl)CH<sub>2</sub>CH<sub>2</sub>-, (cyclopropyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (cyclobutyl)CH<sub>2</sub>CH<sub>2</sub>-, (cyclopentyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 (cyclohexyl)CH<sub>2</sub>CH<sub>2</sub>-, (morpholino)CH<sub>2</sub>CH<sub>2</sub>-, or  
 30 (N-pipridinyl)CH<sub>2</sub>CH<sub>2</sub>-;
- R<sup>10</sup> is H, methyl, ethyl, phenyl, benzyl, phenethyl,  
 4-F-phenyl, (4-F-phenyl)CH<sub>2</sub>-, (4-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 4-Cl-phenyl, (4-Cl-phenyl)CH<sub>2</sub>-, (4-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 35 4-CH<sub>3</sub>-phenyl, (4-CH<sub>3</sub>-phenyl)CH<sub>2</sub>-, (4-CH<sub>3</sub>-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 4-CF<sub>3</sub>-phenyl, (4-CF<sub>3</sub>-phenyl)CH<sub>2</sub>-, or  
 (4-CF<sub>3</sub>-phenyl)CH<sub>2</sub>CH<sub>2</sub>-;

R<sup>11</sup>, at each occurrence, is independently selected from  
 H, =O, methyl, ethyl, phenyl, benzyl, phenethyl,  
 4-F-phenyl, (4-F-phenyl)CH<sub>2</sub>-, (4-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 3-F-phenyl, (3-F-phenyl)CH<sub>2</sub>-, (3-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 5 2-F-phenyl, (2-F-phenyl)CH<sub>2</sub>-, (2-F-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 4-Cl-phenyl, (4-Cl-phenyl)CH<sub>2</sub>-, (4-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 3-Cl-phenyl, (3-Cl-phenyl)CH<sub>2</sub>-, (3-Cl-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 4-CH<sub>3</sub>-phenyl, (4-CH<sub>3</sub>-phenyl)CH<sub>2</sub>-, (4-CH<sub>3</sub>-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 3-CH<sub>3</sub>-phenyl, (3-CH<sub>3</sub>-phenyl)CH<sub>2</sub>-, (3-CH<sub>3</sub>-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 10 4-CF<sub>3</sub>-phenyl, (4-CF<sub>3</sub>-phenyl)CH<sub>2</sub>-, (4-CF<sub>3</sub>-phenyl)CH<sub>2</sub>CH<sub>2</sub>-,  
 pyrid-2-yl, pyrid-3-yl, or pyrid-4-yl, and

R<sup>13</sup>, at each occurrence, is independently selected from  
 H, F, Cl, OH, -CH<sub>3</sub>, -CH<sub>2</sub>CH<sub>3</sub>, -OCH<sub>3</sub>, or -CF<sub>3</sub>.

15

[21] In another embodiment the present invention  
 provides for a method for the treatment of neurological  
 disorders associated with  $\beta$ -amyloid production comprising  
 administering to a host in need of such treatment a  
 20 therapeutically effective amount of a compound of Formula  
 (I):



25 or a pharmaceutically acceptable salt or prodrug thereof,  
 wherein:

Q is -NR<sup>1</sup>R<sup>2</sup>;

30 R<sup>1</sup> is OR<sup>14</sup>;

R<sup>2</sup> is independently selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>10</sub>  
 carbocycle, C<sub>6</sub>-C<sub>10</sub> aryl, and 5 to 10 membered  
 heterocycle containing 1 to 4 heteroatoms selected  
 35 from nitrogen, oxygen, and sulphur;

$R^3$  is  $-(CR^7R^{7a})_n-R^4$ ,  
 $-(CR^7R^{7a})_n-S-(CR^7R^{7a})_m-R^4$ ,  
 $-(CR^7R^{7a})_n-O-(CR^7R^{7a})_m-R^4$ ,  
5  $-(CR^7R^{7a})_n-N(R^{7b})-(CR^7R^{7a})_m-R^4$ ,  
 $-(CR^7R^{7a})_n-S(=O)-(CR^7R^{7a})_m-R^4$ ,  
 $-(CR^7R^{7a})_n-S(=O)_2-(CR^7R^{7a})_m-R^4$ ,  
 $-(CR^7R^{7a})_n-C(=O)-(CR^7R^{7a})_m-R^4$ ,  
 $-(CR^7R^{7a})_n-N(R^{7b})C(=O)-(CR^7R^{7a})_m-R^4$ ,  
10  $-(CR^7R^{7a})_n-C(=O)N(R^{7b})-(CR^7R^{7a})_m-R^4$ ,  
 $-(CR^7R^{7a})_n-N(R^{7b})S(=O)_2-(CR^7R^{7a})_m-R^4$ , or  
 $-(CR^7R^{7a})_n-S(=O)_2N(R^{7b})-(CR^7R^{7a})_m-R^4$ ;

$n$  is 0, 1, 2, or 3;

15

$m$  is 0, 1, 2, or 3;

$R^{3a}$  is H, OH,  $C_1$ - $C_4$  alkyl,  $C_1$ - $C_4$  alkoxy,  $C_2$ - $C_4$  alkenyl  
or  $C_2$ - $C_4$  alkenyloxy;

20

$R^4$  is H, OH,  $OR^{14a}$ ,  
 $C_1$ - $C_6$  alkyl substituted with 0-3  $R^{4a}$ ,  
 $C_2$ - $C_6$  alkenyl substituted with 0-3  $R^{4a}$ ,  
 $C_2$ - $C_6$  alkynyl substituted with 0-3  $R^{4a}$ ,  
25  $C_3$ - $C_{10}$  carbocycle substituted with 0-3  $R^{4b}$ ,  
 $C_6$ - $C_{10}$  aryl substituted with 0-3  $R^{4b}$ , or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
30 is substituted with 0-3  $R^{4b}$ ;

$R^{4a}$ , at each occurrence, is independently selected from is  
H, F, Cl, Br, I,  $CF_3$ ,

35  $C_3$ - $C_{10}$  carbocycle substituted with 0-3  $R^{4b}$ ,  
 $C_6$ - $C_{10}$  aryl substituted with 0-3  $R^{4b}$ , or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and



sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R<sup>4b</sup>;

5 R<sup>4b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl, C<sub>1</sub>-C<sub>4</sub> haloalkoxy, and C<sub>1</sub>-C<sub>4</sub> halothioalkyl-S-;

10 R<sup>5</sup> is H, OR<sup>14</sup>;  
C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 0-3 R<sup>5b</sup>;  
C<sub>1</sub>-C<sub>6</sub> alkoxy substituted with 0-3 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>6</sub> alkenyl substituted with 0-3 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>6</sub> alkynyl substituted with 0-3 R<sup>5b</sup>;  
15 C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>5c</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>5c</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
20 is substituted with 0-3 R<sup>5c</sup>;

R<sup>5a</sup> is H, OH, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>2</sub>-C<sub>4</sub> alkenyl, or  
C<sub>2</sub>-C<sub>4</sub> alkenyloxy;

25 R<sup>5b</sup>, at each occurrence, is independently selected from:  
H, C<sub>1</sub>-C<sub>6</sub> alkyl, CF<sub>3</sub>, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>,  
NR<sup>15</sup>R<sup>16</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>5c</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>5c</sup>; or  
30 5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>5c</sup>;

35 R<sup>5c</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl,

C<sub>1</sub>-C<sub>4</sub> haloalkoxy, and C<sub>1</sub>-C<sub>4</sub> halothioalkyl-S-;

R<sup>6</sup> is H;

C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 0-3 R<sup>6a</sup>;

5 C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>6b</sup>; or

C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>6b</sup>;

10 R<sup>6a</sup>, at each occurrence, is independently selected from H,  
C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>,  
aryl or CF<sub>3</sub>;

15 R<sup>6b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl,  
C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy;

R<sup>7</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, CF<sub>3</sub>, phenyl and C<sub>1</sub>-C<sub>4</sub> alkyl;

20 R<sup>7a</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, CF<sub>3</sub>, and C<sub>1</sub>-C<sub>4</sub> alkyl;

R<sup>7b</sup> is independently selected from H and C<sub>1</sub>-C<sub>4</sub> alkyl;

Ring B is a 7 membered lactam,

25 wherein the lactam is saturated, partially saturated  
or unsaturated;

wherein each additional lactam carbon is substituted  
with 0-2 R<sup>11</sup>; and,

30 optionally, the lactam contains a heteroatom selected  
from -O-, -S-, -S(=O)-, -S(=O)<sub>2</sub>-, -N=, -NH-, and -  
N(R<sup>10</sup>)-;

35 additionally, two R<sup>11</sup> substituents on adjacent atoms may be  
combined to form a benzo fused radical; wherein said  
benzo fused radical is substituted with 0-4 R<sup>13</sup>;

additionally, two R<sup>11</sup> substituents on adjacent atoms may be  
combined to form a 5 to 6 membered heteroaryl fused

radical, wherein said 5 to 6 membered heteroaryl fused radical comprises 1 or 2 heteroatoms selected from N, O, and S; wherein said 5 to 6 membered heteroaryl fused radical is substituted with 0-3 R<sup>13</sup>;

5

additionally, two R<sup>11</sup> substituents on the same or adjacent carbon atoms may be combined to form a C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>13</sup>;

10 R<sup>10</sup> is H, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, S(=O)<sub>2</sub>R<sup>17</sup>;

C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 0-3 R<sup>10a</sup>;

C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>10b</sup>;

C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>10b</sup>; or

15 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R<sup>10b</sup>;

20 R<sup>10a</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, or aryl substituted with 0-4 R<sup>10b</sup>;

25 R<sup>10b</sup>, at each occurrence, is independently selected from H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl, C<sub>1</sub>-C<sub>4</sub> haloalkoxy, and C<sub>1</sub>-C<sub>4</sub> haloalkyl-S-;

30 R<sup>11</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>18</sup>R<sup>19</sup>, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, CF<sub>3</sub>; C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 0-3 R<sup>11a</sup>;

C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>11b</sup>;

35 C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>11b</sup>; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R<sup>11b</sup>;

5 R<sup>11a</sup>, at each occurrence, is independently selected from  
H, C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>,  
NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>;  
phenyl substituted with 0-3 R<sup>11b</sup>;  
C<sub>3</sub>-C<sub>6</sub> cycloalkyl substituted with 0-3 R<sup>11b</sup>; and  
10 5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>11b</sup>;

15 R<sup>11b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>,  
S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>,  
C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl,  
C<sub>1</sub>-C<sub>4</sub> haloalkoxy, and C<sub>1</sub>-C<sub>4</sub> halothioalkyl-S-;

20 Z is H;  
C<sub>1</sub>-C<sub>8</sub> alkyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 1-3 R<sup>12</sup>;  
C<sub>1</sub>-C<sub>8</sub> alkyl substituted with 0-3 R<sup>12a</sup>;  
25 C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>12a</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>12a</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-4 R<sup>12b</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
30 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>12b</sup>;

R<sup>12</sup>, at each occurrence, is independently selected from  
35 C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-4 R<sup>12b</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R<sup>12b</sup>;

- 5 R<sup>12a</sup>, at each occurrence, is independently selected from  
H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, -C(=O)NR<sup>15</sup>R<sup>16</sup>,  
CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>,  
C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl,  
C<sub>1</sub>-C<sub>4</sub> haloalkoxy, or C<sub>1</sub>-C<sub>4</sub> halothioalkyl-S-;
- 10 R<sup>12b</sup>, at each occurrence, is independently selected from  
H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl,  
SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>,  
C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl,  
C<sub>1</sub>-C<sub>4</sub> haloalkoxy, and C<sub>1</sub>-C<sub>4</sub> halothioalkyl-S-;
- 15 R<sup>13</sup>, at each occurrence, is independently selected from  
H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, CN,  
NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;
- 20 R<sup>14</sup> is H, phenyl, benzyl, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkoxyalkyl, or  
C<sub>3</sub>-C<sub>6</sub> cycloalkyl;
- R<sup>14a</sup> is H, phenyl, benzyl, or C<sub>1</sub>-C<sub>4</sub> alkyl;
- 25 R<sup>15</sup>, at each occurrence, is independently selected from H,  
C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-,  
and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-;
- 30 R<sup>16</sup>, at each occurrence, is independently selected from  
H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, phenethyl,  
(C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-;
- R<sup>17</sup> is H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkoxyalkyl,  
aryl substituted by 0-4 R<sup>17a</sup>, or  
35 -CH<sub>2</sub>-aryl substituted by 0-4 R<sup>17a</sup>;

R<sup>17a</sup> is H, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, butoxy, -OH, F, Cl, Br, I, CF<sub>3</sub>, OCF<sub>3</sub>, SCH<sub>3</sub>, S(O)CH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, -NH<sub>2</sub>, -N(CH<sub>3</sub>)<sub>2</sub>, or C<sub>1</sub>-C<sub>4</sub> haloalkyl;

5 R<sup>18</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-; and

10 R<sup>19</sup>, at each occurrence, is independently selected from H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-;

provided, when R<sup>13</sup> is H,  
then Z is H;

15 C<sub>4</sub>-C<sub>8</sub> alkyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 1-3 R<sup>12</sup>;  
C<sub>1</sub>-C<sub>8</sub> alkyl substituted with 0-3 R<sup>12a</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>12a</sup>; or  
20 C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>12a</sup>; and

provided, when ring B is a 1,3,4,5-tetrahydro-1-(Z)-5-(R<sup>10</sup>)-6,6,7,7-tetra(R<sup>11</sup>)-2,4-dioxo-2H-1,5-diazepin-3-yl core, and R<sup>13</sup> is H; then

25 R<sup>10</sup> is H, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, S(=O)<sub>2</sub>R<sup>17</sup>; or  
C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 0-3 R<sup>10a</sup>;

30 R<sup>10a</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>.

In a more preferred embodiment of the present  
35 invention, Q is NH<sub>2</sub>.

In a most preferred embodiment the total number of carbon atoms in R<sup>3</sup>, R<sup>3a</sup>, R<sup>5</sup>, and R<sup>5a</sup> equals seven or more.

It is understood that any and all embodiments of the present invention may be taken in conjunction with any other embodiment to describe additional even more preferred  
5   embodiments of the present invention.

In a second embodiment, the present invention provides a pharmaceutical composition comprising a compound of Formula (I) and a pharmaceutically acceptable carrier.  
10

In a third embodiment, the present invention provides a method for the treatment of neurological disorders associated with  $\beta$ -amyloid production comprising administering to a host in need of such treatment a  
15   therapeutically effective amount of a compound of Formula (I).

In a preferred embodiment the neurological disorder associated with  $\beta$ -amyloid production is Alzheimer's  
20   Disease.

In a fourth embodiment, the present invention provides a method for inhibiting  $\gamma$ -secretase activity for the treatment of a physiological disorder associated with  
25   inhibiting  $\gamma$ -secretase activity comprising administering to a host in need of such inhibition a therapeutically effective amount of a compound of Formula (I) that inhibits  $\gamma$ -secretase activity.

Thus, the present invention provides a method for inhibiting  $\gamma$ -secretase activity comprising administering to a host in need of such inhibition a therapeutically effective amount of a compound of Formula (I) that inhibits  $\gamma$ -secretase activity.  
30  
35

In a preferred embodiment the physiological disorder associated with inhibiting  $\gamma$ -secretase activity is Alzheimer's Disease.

In a fifth embodiment, the present invention provides a compound of Formula (I) for use in therapy.

5 In a preferred embodiment the present invention provides a compound of Formula (I) for use in therapy of Alzheimer's Disease.

10 In a sixth embodiment, the present invention provides for the use of a compound of Formula (I) for the manufacture of a medicament for the treatment of Alzheimer's Disease.

#### 15 DEFINITIONS

As used herein, the term "A $\beta$ " denotes the protein designated A $\beta$ ,  $\beta$ -amyloid peptide, and sometimes  $\beta$ /A4, in the art. A $\beta$  is an approximately 4.2 kilodalton (kD) protein of about 39 to 43 amino acids found in amyloid  
20 plaques, the walls of meningeal and parenchymal arterioles, small arteries, capillaries, and sometimes, venules. The isolation and sequence data for the first 28 amino acids are described in U.S. Pat. No 4,666,829. The 43 amino acid sequence is:

25

```

1
Asp  Ala  Glu  Phe  Arg  His  Asp  Ser  Gly  Tyr
11
Glu  Val  His  His  Gln  Lys  Leu  Val  Phe  Phe
21
Ala  Glu  Asp  Val  Gly  Ser  Asn  Lys  Gly  Ala
31
Ile  Ile  Gly  Leu  Met  Val  Gly  Gly  Val  Val
41
Ile  Ala  Thr

```

The term "APP", as used herein, refers to the protein known in the art as  $\beta$  amyloid precursor protein. This



protein is the precursor for A $\beta$  and through the activity of "secretase" enzymes, as used herein, it is processed into A $\beta$ . Differing secretase enzymes, known in the art, have been designated  $\beta$  secretase, generating the N-terminus of A $\beta$ ,  $\alpha$  secretase cleaving around the 16/17 peptide bond in A $\beta$ , and " $\gamma$  secretases", as used herein, generating C-terminal A $\beta$  fragments ending at position 38, 39, 40, 42, and 43 or generating C-terminal extended precursors which are subsequently truncated to the above polypeptides.

10       The compounds herein described may have asymmetric centers. Compounds of the present invention containing an asymmetrically substituted atom may be isolated in optically active or racemic forms. It is well known in the art how to prepare optically active forms, such as by  
15 resolution of racemic forms or by synthesis from optically active starting materials. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the present invention. Cis and  
20 trans geometric isomers of the compounds of the present invention are described and may be isolated as a mixture of isomers or as separated isomeric forms. All chiral, diastereomeric, racemic forms and all geometric isomeric forms of a structure are intended, unless the specific  
25 stereochemistry or isomeric form is specifically indicated.

      The term "substituted," as used herein, means that any one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valency is not exceeded, and  
30 that the substitution results in a stable compound. When a substituent is keto (i.e., =O), then 2 hydrogens on the atom are replaced.

      When any variable (e.g., R<sup>5b</sup>) occurs more than one time in any constituent or formula for a compound, its  
35 definition at each occurrence is independent of its definition at every other occurrence. Thus, for example, if a group is shown to be substituted with 0-2 R<sup>5b</sup>, then said group may optionally be substituted with up to two R<sup>5b</sup>

groups and R<sup>5b</sup> at each occurrence is selected independently from the definition of R<sup>5b</sup>. Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

- 5        When a bond to a substituent is shown to cross a bond connecting two atoms in a ring, then such substituent may be bonded to any atom on the ring. When a substituent is listed without indicating the atom via which such substituent is bonded to the rest of the compound of a  
10    given formula, then such substituent may be bonded via any atom in such substituent. Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

- As used herein, "alkyl" or "alkylene" is intended to  
15    include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms; for example, "C<sub>1</sub>-C<sub>6</sub> alkyl" denotes alkyl having 1, 2, 3, 4, 5, or 6 carbon atoms. Examples of alkyl include, but are not limited to, methyl, ethyl, n-propyl,  
20    i-propyl, n-butyl, i-butyl, sec-butyl, t-butyl, pentyl, and hexyl. Preferred "alkyl" group, unless otherwise specified, is "C<sub>1</sub>-C<sub>4</sub> alkyl". Additionally, unless otherwise specified, "propyl" denotes n-propyl or i-propyl; "butyl" denotes n-butyl, i-butyl, sec-butyl, or t-butyl.

- 25        As used herein, "alkenyl" or "alkenylene" is intended to include hydrocarbon chains of either a straight or branched configuration and one or more unsaturated carbon-carbon bonds which may occur in any stable point along the chain. Examples of "C<sub>2</sub>-C<sub>6</sub> alkenyl" include, but  
30    are not limited to, ethenyl, 1-propenyl, 2-propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 3-methyl-2-butenyl, 2-pentenyl, 3-pentenyl, hexenyl, and the like.

- As used herein, "alkynyl" or "alkynylene" is intended to include hydrocarbon chains of either a straight or  
35    branched configuration and one or more carbon-carbon triple bonds which may occur in any stable point along the chain, such as ethynyl, 1-propynyl, 2-propynyl, 1-butynyl, 2-butynyl, 3-butynyl, and the like.

"Alkoxy" or "alkyloxy" represents an alkyl group as defined above with the indicated number of carbon atoms attached through an oxygen bridge. Examples of alkoxy include, but are not limited to, methoxy, ethoxy, 5 n-propoxy, i-propoxy, n-butoxy, s-butoxy, t-butoxy, n-pentoxo, and s-pentoxo. Preferred alkoxy groups are methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, s-butoxy, t-butoxy. Similarly, "alkylthio" or "thioalkoxy" is 10 represents an alkyl group as defined above with the indicated number of carbon atoms attached through a sulphur bridge.

"Halo" or "halogen" as used herein refers to fluoro, chloro, bromo, and iodo. Unless otherwise specified, preferred halo is fluoro and chloro. "Counterion" is used 15 to represent a small, negatively charged species such as chloride, bromide, hydroxide, acetate, sulfate, and the like.

"Haloalkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups 20 having the specified number of carbon atoms, substituted with 1 or more halogen (for example  $-C_vF_w$  where  $v = 1$  to  $3$  and  $w = 1$  to  $(2v+1)$ ). Examples of haloalkyl include, but are not limited to, trifluoromethyl, trichloromethyl, pentafluoroethyl, pentachloroethyl, 2,2,2-trifluoroethyl, 25 2,2-difluoroethyl, heptafluoropropyl, and heptachloropropyl. "Haloalkoxy" is intended to mean a haloalkyl group as defined above with the indicated number of carbon atoms attached through an oxygen bridge; for example trifluoromethoxy, pentafluoroethoxy, 2,2,2- 30 trifluoroethoxy, and the like. "Halothioalkoxy" is intended to mean a haloalkyl group as defined above with the indicated number of carbon atoms attached through a sulphur bridge.

"Cycloalkyl" is intended to include saturated ring 35 groups, having the specified number of carbon atoms. For example, "C<sub>3</sub>-C<sub>6</sub> cycloalkyl" denotes such as cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl.

As used herein, "carbocycle" is intended to mean any stable 3- to 7-membered monocyclic or bicyclic or 7- to 13-membered bicyclic or tricyclic, any of which may be saturated, partially unsaturated, or aromatic. Examples of such carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, adamantyl, cyclooctyl, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane (decalin), [2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, or tetrahydronaphthyl (tetralin). Preferred "carbocycle" are cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

As used herein, the term "heterocycle" or "heterocyclic ring" is intended to mean a stable 5- to 7-membered monocyclic or bicyclic or 7- to 14-membered bicyclic heterocyclic ring which is saturated partially unsaturated or unsaturated (aromatic), and which consists of carbon atoms and 1, 2, 3 or 4 heteroatoms independently selected from the group consisting of N, O and S and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The nitrogen and sulfur heteroatoms may optionally be oxidized. The heterocyclic ring may be attached to its pendant group at any heteroatom or carbon atom which results in a stable structure. The heterocyclic rings described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. If specifically noted, a nitrogen in the heterocycle may optionally be quaternized. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one another. It is preferred that the total number of S and O atoms in the heterocycle is not more than 1.


Examples of heterocycles include, but are not limited to, 1H-indazole, 2-pyrrolidonyl, 2H,6H-1,5,2-dithiazinyl, 2H-pyrrolyl, 3H-indolyl, 4-piperidonyl, 4aH-carbazole, 4H-quinoliziny, 6H-1,2,5-thiadiazinyl, acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl,

- benzothiophenyl, benzoxazolyl, benzthiazolyl,  
benztriazolyl, benztetrazolyl, benzisoxazolyl,  
benzisothiazolyl, benzimidazolyl, carbazolyl,  
4aH-carbazolyl, b-carbolinyl, chromanyl, chromenyl,  
5 cinnolyl, decahydroquinolyl, 2H,6H-1,5,2-dithiazinyl,  
dihydrofuro[2,3-b]tetrahydrofuran, furanyl, furazanyl,  
imidazolidinyl, imidazolyl, imidazolyl, 1H-indazolyl,  
indolenyl, indolyl, indolizyl, indolyl,  
isobenzofuranyl, isochromanyl, isoindazolyl, isoindolyl,  
10 isoindolyl, isoquinolyl, isothiazolyl, isoxazolyl,  
morpholyl, naphthyridyl, octahydroisoquinolyl,  
oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl,  
1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl,  
oxazolyl, oxazolidinylperimidyl, phenanthridyl,  
15 phenanthrolinyl, phenarsazinyl, phenazinyl, phenothiazinyl,  
phenoxathiinyl, phenoxazinyl, phthalazinyl, piperazinyl,  
piperidinyl, pteridinyl, piperidonyl, 4-piperidonyl,  
pteridinyl, purinyl, pyranyl, pyrazinyl, pyrazolidinyl,  
pyrazolyl, pyrazolyl, pyridazinyl, pyridoxazole,  
20 pyridoimidazole, pyridothiazole, pyridinyl, pyridyl,  
pyrimidinyl, pyrrolidinyl, pyrrolinyl, pyrrolyl,  
quinazolyl, quinolyl, 4H-quinolizyl, quinoxalyl,  
quinuclidyl, carbolinyl, tetrahydrofuranyl,  
tetrahydroisoquinolyl, tetrahydroquinolyl,  
25 6H-1,2,5-thiadiazinyl, 1,2,3-thiadiazolyl,  
1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl,  
thianthrenyl, thiazolyl, thienyl, thienothiazolyl,  
thienooxazolyl, thienoimidazolyl, thiophenyl, triazinyl,  
1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl,  
30 1,3,4-triazolyl, xanthenyl. Preferred 5 to 10 membered  
heterocycles include, but are not limited to, pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl,  
tetrazolyl, benzofuranyl, benzothiofuranyl, indolyl,  
35 benzimidazolyl, 1H-indazolyl, oxazolidinyl, isoxazolidinyl,  
benzotriazolyl, benzisoxazolyl, oxindolyl, benzoxazolyl,  
quinolyl, and isoquinolyl. Preferred 5 to 6 membered  
heterocycles include, but are not limited to, pyridinyl,

pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, tetrazolyl; more preferred 5 to 6 membered heterocycles include, but are not limited to, pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, and tetrazolyl. Also included are fused ring and spiro compounds containing, for example, the above heterocycles.

As used herein, the term "aryl", "C<sub>6</sub>-C<sub>10</sub> aryl" or aromatic residue, is intended to mean an aromatic moiety containing the specified number of carbon atoms; for example phenyl, pyridinyl or naphthyl. Preferred "aryl" is phenyl. Unless otherwise specified, "aryl" may be unsubstituted or substituted with 0 to 3 groups selected from H, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, butoxy, amino, hydroxy, Cl, F, Br, I, CF<sub>3</sub>, SCH<sub>3</sub>, S(O)CH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, -N(CH<sub>3</sub>)<sub>2</sub>, N(CH<sub>3</sub>)H, CN, NO<sub>2</sub>, OCF<sub>3</sub>, C(=O)CH<sub>3</sub>, CO<sub>2</sub>H, CO<sub>2</sub>CH<sub>3</sub>, or C<sub>1</sub>-C<sub>4</sub> haloalkyl.


The phrase "additional lactam carbons", as used herein, is intended to denote the number of optional carbon atoms in the lactam ring B of Formula (I). Formula (I"):




(I")


represents the lactam ring B of Formula (I). Additional lactam carbons are carbons in lactam ring B other than the carbons numbered 2 and 3 in the backbone of the formula. The additional lactam carbons may be optionally replaced by a heteroatom selected from oxygen, nitrogen and sulfur. Lactam ring B contains 1, 2, 3, 4, 5, 6 or 7 optional carbons, wherein one optional carbon may optionally be replaced by a heteroatom, such that the total number of members of lactam ring B, including atoms numbered 1, 2 and 3 in the backbone, does not exceed 10. It is preferred

that the total number of atoms of lactam ring B is 6, 7 or 8; it is more preferred that the total number of atoms of lactam ring B is seven. It is further understood that lactam ring B may optionally be unsaturated or partially unsaturated (i.e. two adjacent atoms in the ring form a double bond) wherein the backbone of lactam ring B may contain one, two or three double bonds. Examples of lactam ring B include:






B15



B16


but are not intended to limit the invention. Preferred  
 5 examples of lactam ring B are B1, B2, B5, B6, B8, B9, B13,  
 and B16; more preferred examples of lactam ring B are B1,  
 B6, B8, B9, and B13. Preferred examples of substituent R<sup>10</sup>  
 or R<sup>11</sup> on lactam B are methyl, ethyl, phenyl, 4-  
 fluorophenyl, 4-chlorophenyl, 4-trifluoromethylphenyl, (4-  
 10 fluorophenyl)methyl, (4-chlorophenyl)methyl, (4-  
 trifluoromethylphenyl)methyl, and 2-, 3-, and 4-pyridinyl.  
 Preferred examples of R<sup>13</sup> on lactam B are F, Cl, OH,  
 methyl, ethyl, methoxy, and trifluoromethyl.

15 The compounds herein described may have asymmetric  
 centers. One enantiomer of a compound of Formula (I) may  
 display superior biological activity over the opposite  
 enantiomer. For example carbon 3 of lactam ring B Formula  
 (I") may exist in either an S or R configuration. Thus, an  
 20 R or S configuration at carbon 3 in Formula (I") is  
 considered part of the invention. An example of such  
 configuration includes,



25

and





but is not intended to be limited to this example of ring B. When required, separation of the racemic material can be achieved by methods known in the art. Additionally, the carbon atoms to which  $R^3$  and  $R^5$  are attached may describe chiral carbons which may display superior biological activity over the opposite enantiomer. For example, where  $R^3$  and  $R^5$  are not H, then the configuration of the two centers may be described as (2R,3R), (2R,3S), (2S,3R), or (2S,3S). All configurations are considered part of the invention; however, the (2R,3S) and the (2S,3R) are preferred and the (2R,3S) is more preferred.

The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic,

phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, and the like.

- 5     The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a
- 10 stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical*
- 15 *Sciences*, 17th ed., Mack Publishing Company, Easton, PA, 1985, p. 1418, the disclosure of which is hereby incorporated by reference.

- "Prodrugs" are intended to include any covalently bonded carriers which release the active parent drug
- 20 according to formula (I) *in vivo* when such prodrug is administered to a mammalian subject. Prodrugs of a compound of formula (I) are prepared by modifying functional groups present in the compound in such a way that the modifications are cleaved, either in routine
- 25 manipulation or *in vivo*, to the parent compound. Prodrugs include compounds of formula (I) wherein a hydroxy, amino, or sulfhydryl group is bonded to any group that, when the prodrug or compound of formula (I) is administered to a mammalian subject, cleaves to form a free hydroxyl, free
- 30 amino, or free sulfhydryl group, respectively. Examples of prodrugs include, but are not limited to, acetate, formate and benzoate derivatives of alcohol and amine functional groups in the compounds of formula (I), and the like.

- "Stable compound" and "stable structure" are meant to
- 35 indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

### SYNTHESIS

The compounds of the present invention can be prepared in a number of ways well known to one skilled in the art of organic synthesis. The compounds of the present invention  
5 can be synthesized using the methods described below, together with synthetic methods known in the art of synthetic organic chemistry, or variations thereon as appreciated by those skilled in the art. Preferred methods include, but are not limited to, those described below.  
10 All references cited herein are hereby incorporated in their entirety herein by reference.

The novel compounds of this invention may be prepared using the reactions and techniques described in this section. The reactions are performed in solvents  
15 appropriate to the reagents and materials employed and are suitable for the transformations being effected. Also, in the description of the synthetic methods described below, it is to be understood that all proposed reaction conditions, including choice of solvent, reaction  
20 atmosphere, reaction temperature, duration of the experiment and workup procedures, are chosen to be the conditions standard for that reaction, which should be readily recognized by one skilled in the art. It is understood by one skilled in the art of organic synthesis  
25 that the functionality present on various portions of the molecule must be compatible with the reagents and reactions proposed. Such restrictions to the substituents which are compatible with the reaction conditions will be readily apparent to one skilled in the art and alternate methods  
30 must then be used.

Methods for the synthesis of succinylamino lactams are known in the art and are disclosed in a number of references including PCT publication number WO 96/29313, which is hereby incorporated by reference.


35 Disubstituted succinate derivatives can be prepared by a number of known procedures. The procedure of Evans (D. A. Evans et al, *Org. Synth.* **86**, p83 (1990)) is outlined in Scheme 1 where acylation of an oxazolidinone with an

acylating agent such as an acid chloride provides structures 1. Alkylation to form 2 followed by cleavage of the chiral auxiliary and subsequent alkylation of the dianion of the carboxylic acid 3 provides a variety of

5 disubstituted succinates which can be separated and incorporated into structures of formula (I) by those skilled in the art. Additional examples are found in P. Becket, M. J. Crimmin, M. H. Davis, Z. Spavold, Synlett, (1993), 137-138, incorporated herein by reference.

10

Scheme 1




15 Diastereomerically pure succinate derivatives can be accessed using the chemistry outlined below, adapted from P. Becket, M. J. Crimmin, M. H. Davis, Z. Spavold, Synlett, (1993), 137-138 incorporated herein by reference. This reference provides the synthesis below to obtain compound

20 9, Scheme 2. Compound 11, Scheme 2, is used as an intermediate and is prepared from 9 by hydrogenation of the allyl group followed by coupling of 9-fluorenemethanol under standard conditions using DCC and DMAP in CH<sub>2</sub>Cl<sub>2</sub>. Deprotection of the tert-butyl ester is accomplished by


25 treatment with 50% trifluoroacetic acid.

Scheme 2



A variety of compounds of Formula (I) can be prepared by methods described in Scheme 4. The protected  $\alpha$ -amine 3 of the  $\alpha$ -amino- $\epsilon$ -caprolactam can be prepared by methods well known in the literature for amino protecting groups as discussed in Theodora W. Greene's book "Protective Groups in Organic Synthesis", like *N*-Boc using di-*t*-butyldicarbonate in an appropriate solvent like DMSO. A sulfur atom can be introduced into the ring providing L- $\alpha$ -amino- $\beta$ -thio- $\epsilon$ -caprolactam according to the procedure in S. A. Ahmed et al, FEBS Letters, (1984), vol. 174, pages 76-9 (Scheme 3). One skilled in the art can extend this methodology to the synthesis of  $\beta$ -amino and oxygen containing rings by analogy. The sulfur-containing molecules can also be oxidized to the sulfoxide and sulfone by methods known to one skilled in the art.

Scheme 3




- The lactam nitrogen of compound 13 can be alkylated by
- 5 generating the anion with bases such as LDA, lithium bis(trimethylsilyl)amide or sodium hydride in solvents like THF, with or without cosolvents such as DMPU or HMPA and reacting this with a variety of groups containing leaving groups ( $\text{X}^{\circ}$ ) like bromide, iodide, mesylate or tosylate.
  - 10 Alkylating agents such as  $\alpha$ -bromo amides, ketones and acids can be prepared by a number of literature methods including halogenation of amino acids by diazotization or are commercially available. Other suitable alkylating agents such as alkyl, allylic and benzylic halides can be formed
  - 15 form a variety of precursors such as free-radical addition of halides or activation of alcohols, and other chemistries known to those skilled in the art. For discussion of these types of reactions, see Carey, F.A. and Sundberg, R. J., Advanced Organic Chemistry, Part A, New York: Plenum Press, 1990, pages 304-305, 342-347, 695-698.
  - 20

- The *N*-Boc protecting group can be removed by any number of methods well known in the literature like TFA in methylene chloride to give the compound 15. The amine 15 can be coupled to an appropriately substituted carboxylic acid or acid chloride by methods well described in the literature for making amide bonds, like TBTU in DMF with a base like NMM to give the elaborated compound 16.
- Compounds 16 can be alkylated using standard bases like LDA, NaH, or NaHMDS to deprotonate the amide followed by
- 25 addition of an alkylating agent with an appropriate leaving group like halide, mesylate, or triflate in an appropriate solvent to provide compounds 17 with an  $\text{R}^6$  substituent.
  - 30

The *t*-butyl ester is then removed by treatment with TFA in methylene chloride to give the carboxylic acid **17**.

Scheme 4



The final compounds **18** were prepared by treating the activated carboxylic acid of **17** with an appropriately substituted amine. For instance, activation of the carboxylic acid with HATU (*O*-(7-azabenzotriazol-1-yl)-1,1,3,3,-tetramethyluronium hexafluorophosphate) or PyBOP (benzotriazole-1-yl-oxy-tris-pyrrolidino-phosphonium hexafluorophosphate) or other coupling agents known to those skilled in the art allows condensation with ammonia to form primary amides. Similarly, condensation of the activated acid with hydroxylamine hydrochloride provides the hydroxamic acid, or reaction with a primary or secondary amine provides the substituted amine derivative. Activation of the acid with PyBrOP (bromo-tris-pyrrolidino-phosphonium hexafluorophosphate) followed by addition of an alcohol and 4-dimethylaminopyridine allows formation of the ester directly. For additional acylation reactions see for example Carey, F.A. and Sundberg, R. J., Advanced Organic


Chemistry, Part A, New York: Plenum Press, 1990, pages 475-479.

Additional intermediates in the synthesis of compounds of formula (I) can be prepared as shown in Scheme 5. A suitable resin for solid phase synthesis such as Fmoc (Fluorenylmethylcarbonyl)-protected hydroxylamine bound to polystyrene beads can be purchased from Novabiochem, Inc. Deprotection of the Fmoc group under standard conditions using 20% piperidine in DMF provides trityl-linked hydroxylamine resin. Coupling of a fluorenylmethyl-protected succinic acid derivative such as 20 with a coupling agent such as HATU in a suitable solvent like DMF or N-methylpyrrolidinone provides the support-bound hydroxamate 21. The Fluorenylmethyl ester can be removed using 20% piperidine in DMF to provide the free carboxylic acid which can be coupled to amines like the caprolactam 22 (which is available using chemistry outlined in Scheme 4) using PyBOP (benzotriazole-1-yl-oxy-tris-pyrrolidino-phosphonium hexafluorophosphate) and a suitable base like DIEA in DMF or NMP. The support-bound intermediate 23 can then be elaborated to biaryl structures of the type 24 using typical Suzuki coupling conditions employing a catalyst such as Palladium complexes like tetrakis(triphenylphosphine)-palladium with 2M aqueous sodium carbonate as a base in a suitable solvent like THF or DME and an excess of a boronic acid. The final compounds are liberated from the support employing dilute (5%) trifluoroacetic acid in  $\text{CH}_2\text{Cl}_2$  and purified by conventional chromatography.

30




## Scheme 5



The compounds of formula (I) of the present invention  
 5 can also be prepared from aminolactams (27) and succinic  
 acid derivatives (11) using amide bond syntheses known in  
 the art, including methods commonly used in peptide  
 syntheses, such as HATU, TBTU, BOP, pyBOP, EDC, CDI, DCC,  
 hydroxysuccinimide, mixed carboxylic anhydride, and phenyl  
 10 ester mediated couplings, as illustrated in Scheme 6 for  
 the synthesis of (Ia), a particular embodiment of (I).  
 Depending on the structure of


## Scheme 6

15




the final product, it will be appreciated by those skilled in the art that protecting groups or precursor functionality convertible to the desired groups may be desirable. Protecting groups and their use in synthesis are described in Green and Wuts, *Protective Groups in Organic Synthesis*, (Wiley 1991). This is further illustrated in Scheme 6a, in which the succinate half-ester (9) (Becket et al., *Synlett* 1993, 137-138) is coupled to the aminobenzodiazepine (30) (Sherrill and Sugg, *J. Org. Chem.* 1995, 60, 730-734; Bock et al., *J. Med. Chem.*, 1993, 36, 4276-4292) to give ester (Ib), followed by conversion of the ester group to the primary amide (Ic).

Scheme 6a



Methods for the synthesis of lactams, including amino benzodiazepines, are known in the art and are disclosed in a number of references including PCT publication number WO 98/28268, which is hereby incorporated by reference. Additional examples of the synthesis of ring B lactams within the scope of the invention are shown below in Schemes 7, 8, 9, and 10.

Scheme 7



Step 1: Preparation of 72;

5 Following the literature procedures: Stanetty, P.; Koller, H.; Mihovilovic, M. *J. Org. Chem.*, 1992, 57, 6833-6837; Muchowski, J.M.; Venuti, M.D. *J. Org. Chem.*, 1980, 57, 4798-4801.

10 A stirred mixture of 71 (13.0 g, 67.3 mmol) in diether ether (100 mL) under nitrogen was cooled to below -50° in a dry ice/acetone bath. A solution of t-butyllithium in hexanes (2.5 M, 90 mL) was added dropwise. The cooling bath was removed and the reaction allowed to warm to

approximately -25°C and then the temperature was maintained between -20°C and -10°C for 3.25 hours with a dry ice/ethylene glycol cooling bath. The reaction was cooled again to below -50°C and 4-pyridine carboxaldehyde (8.3 mL, 5 87.5 mmol) was added and the reaction stirred at approximately -15°C for 1.25 hours. The reaction was quenched with a saturated solution of ammonium chloride and the mixture stirred overnight. Hexanes (100 mL) was added. The reaction was filtered, washed with water and dried in 10 vacuo to give **72** (15.0 g, 74% as a tan solid: <sup>1</sup>HNMR (300 MHz, DMSO-d<sub>6</sub>) δ 8.55 (s, 1 H), 8.48 (d, 2 H), 7.50 (d, 1 H), 7.36 (d, 1H), 7.24 (d, 2 H), 7.23 (m, 1 H), 7.09 (t of d, 1 H), 6.68 (d, 1 H), 5.93 (d, 1 H), 1.38 (s, 9 H; API MS m/z = 301 [C<sub>17</sub>H<sub>20</sub>N<sub>2</sub>O<sub>3</sub>+H]<sup>+</sup>).

15

Step 2: Preparation of **73**;

To a vigorously stirred solution of **72** (1.21 g, 4.03 mmol) in methylene chloride (45 mL) was added manganese oxide (5.2 g, 59.8 mmol). After 5 hours the reaction was 20 filtered over celite and the celite washed with methylene chloride. The filtrate was concentrated to an oil and triturated with hexanes. The hexanes were removed via pipette and the solid was dried in vacuo to give a **73** (1.01 g, 84%) as a tan solid: <sup>1</sup>HNMR (300 MHz, CDCl<sub>3</sub>) δ 10.25 (s, 25 1 H), 8.80 (d, 2H), 8.50 (d, 1H), 7.48 (t, 1 H), 7.47 (d, 2 H), 7.43 (d, 1 H), 7.00 (t, 1 H), 1.53 (s, 9 H); API MS m/z = 299 [C<sub>17</sub>H<sub>18</sub>N<sub>2</sub>O<sub>3</sub>+H]<sup>+</sup>.

Step 3: Preparation of **74**;

30 Following the literature procedure; Nudelman, Ayetlet; Bechor, Y.; Falb, E.; Fischer, B.; Wexler, B.A.; Nudelman, Abraham; Syn. Com., 1998, 28(3), 471-474.

Acetyl chloride (57 mL, 801 mmol) was added to ice cold methanol (280 mL) with vigorous stirring. Solid **73** (9.0 g., 30.2 mmol) was added and the reaction stirred overnight. The methanol was evaporated in vacuo and the  
5 residue dissolved in water and cooled in an ice bath. The pH was adjusted to approximately 9 with 6N NaOH. The resulting yellow precipitate was collected by vacuum filtration and washed with water and dried in vacuo at 40°C to give **74** (5.71 g, 95%) as a yellow solid: <sup>1</sup>HNMR (300  
10 MHz, CDCl<sub>3</sub>) δ 8.77 (d, 2 H), 7.43 (d, 2 H), 7.33 (m, 2 H), 6.74 (d, 1 H), 6.49 (t, 1 H), 6.30 (broad s, 2 H); CI MS m/z = 199 [C<sub>12</sub>H<sub>10</sub>N<sub>2</sub>O<sub>1</sub>+H]<sup>+</sup>.

Step 4: Preparation of **76**;

15 The benzotriazole acid **75** (13.6 g, 41.7 mmol) and a catalytic amount of DMF were dissolved in dry THF (130 mL.) The reaction was cooled in an ice/salt bath. Oxalyl chloride (3.6 mL, 41.2 mmol) was added dropwise over 1 hour. The reaction was stirred an additional 2.5 hours. A  
20 solution of **74** (5.5 g, 27.7 mmol) in THF (50 mL) was added dropwise forming a thick precipitate. Water and ether were added and the layers separated. The aqueous layer was extracted with methylene chloride. The combined organics were washed with saturated sodium bicarbonate, dried over  
25 magnesium sulfate and concentrated. The resulting oil was chromatographed over silica gel (1:1 methyl acetate/methylene chloride) to provide **76** (12.2 g, 86%) as a yellow solid: <sup>1</sup>HNMR (300 MHz, CDCl<sub>3</sub>) δ 11.62 (s, 1 H), 8.78 (d, 2 H), 8.63 (d, 1 H), 8.08 (d, 1 H), 7.71 (broad s, 1 H), 7.63 (t, 1 H), 7.54 (m, 1 H), 7.44 (d, 1 H), 7.40-  
30 7.19 (m, 11 H), 7.17 (t, 1 H), 7.02 (broad s, 1 H), 5.24-5.07 (m, 2 H); API MS m/z = 507 [C<sub>28</sub>H<sub>22</sub>N<sub>6</sub>O<sub>4</sub>+H]<sup>+</sup>.

Step 5: Preparation of **77**;

Based on the literature procedures; Semple, G.; Ryder, H.; Ohta, M.; Satoh, M. *Syn. Comm.*, **1996**, 26(4), 721-727. Semple, G.; Ryder, H.; Rooker, D.P.; Batt, A.R.; Kendrick, D.A.; Szelke, M.; Ohta, M.; Satoh, M.; Nishida, A.;  
5 Akuzawa, S.; Miyata, K. *J. Med. Chem.*, **1997**, 40, 331-341.

To stirred ice cold methanol (40 mL) was added **76** (4.0 g, 7.9 mmol) and ammonium hydroxide (20 mL.) The reaction was stirred for 2.5 hours forming a heavy precipitate. The  
10 reaction mixture was added slowly to cold acetic acid (40 mL) and stirred overnight at room temperature. The reaction mixture was evaporated and water added. The solution was adjusted to pH 9 with concentrated ammonium hydroxide and stirred 0.5 hour. The resulting tan solid  
15 was collected by vacuum filtration and dried *in vacuo*. The solid was stirred in ethyl acetate (20 mL) and heated under reflux for 1 hour. The mixture was cooled in ice and filtered. The product was washed with cold ethyl acetate and dried *in vacuo* to give **77** (2.2 g, 72%) as a white  
20 solid:  $^1\text{H NMR}$  (300 MHz, DMSO- $d_6$ )  $\delta$  10.94 (s, 1 H), 8.68 (d, 2 H), 8.51 (d, 1 H), 7.57 (t, 1 H), 7.43-7.25 (m, 10 H), 5.10 (d, 2 H); CI MS  $m/z$  = 387 [ $\text{C}_{22}\text{H}_{18}\text{N}_4\text{O}_3 + \text{H}$ ] $^+$ .

Step 6: Preparation of **78**;


25 Following the literature procedures; Semple, G.; Ryder, H.; Ohta, M.; Satoh, M. *Syn. Comm.*, **1996**, 26(4), 721-727. Semple, G.; Ryder, H.; Rooker, D.P.; Batt, A.R.; Kendrick, D.A.; Szelke, M.; Ohta, M.; Satoh, M.; Nishida, A.; Akuzawa, S.; Miyata, K. *J. Med. Chem.*, **1997**, 40, 331-  
30 341.

A stirred solution of **77** (1.0 g, 2.59 mmol) in DMF (10 mL) under nitrogen was cooled in ice. Sodium hydride (62 mg, 2.59 mmol) was added and the reaction stirred 2 hours.  
35 Methyl iodide (0.16 mL, 2.59 mmol) was added and the

reaction stirred 2 hours at room temperature. The reaction was evaporated and chloroform and water were added. The layers were separated and the aqueous extracted with chloroform. The combined organics were washed with brine, dried over sodium sulfate, decanted, evaporated and dried in vacuo. The crude product was passed through a pad a silica gel (1:1 ethyl acetate/methylene chloride) and evaporated to give **78** (934 mg, 90%) was a tan solid:  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  8.59 (d, 2 H), 7.62 (t, 1 H), 7.52 (d, 1 H), 7.42-7.25 (m, 9 H), 6.72 (d, 1 H), 5.38-5.30 (m, 1 H), 5.15 (s, 2 H), 3.48 (s, 3 H); CI MS  $m/z$  = 401  $[\text{C}_{23}\text{H}_{20}\text{N}_4\text{O}_3+\text{H}]^+$ .

## Scheme 8

15



Using the procedures described in Scheme 7 the 3-pyridyl isomer can be prepared. The following data characterizes the individual intermediates.

Intermediate **82**;

$^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ )  $\delta$  8.61 (broad s, 1 H), 8.49 (s, 1 H), 8.42 (d, 1 H), 7.61 (d, 1 H), 7.49 (d, 1 H), 7.40 (d, 1 H), 7.34-7.21 (m, 2 H), 7.40 (t, 1 H), 6.51 (broad s, 1 H), 6.00 (s, 1 H), 1.39 (s, 9 H).

Intermediate **83**;



$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  10.11 (s, 1 H), 8.92 (s, 1 H), 8.80 (d, 1 H), 8.48 (d, 1 H), 8.02 (d, 1 H), 7.59 (t, 1 H), 7.52-7.42 (m, 2 H), 7.05 (t, 1 H), 1.54 (s, 9 H).

5 Intermediate 84;

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  8.84 (s, 1 H), 8.73 (d, 1 H), 7.93 (d, 1 H), 7.44-7.37 (m, 2 H), 7.32 (t, 1 H), 6.75 (d, 1 H), 6.60 (t, 1 H), 6.22 (broad s, 2 H); CI MS  $m/z$  = 199  $[\text{C}_{12}\text{H}_{10}\text{N}_2\text{O}_1+\text{H}]^+$ .

10

Intermediate 86;

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  11.51 (s, 1 H), 8.81-8.75 (m, 2 H), 8.60 (d, 1 H), 8.07 (d, 1 H), 7.89 (d of t, 1 H), 7.80 (broad s, 1 H), 7.63 (t, 1 H), 7.52 (d, 2 H), 7.42-7.15 (m, 9 H), 6.94 (broad s, 1 H), 5.22-5.02 (m, 2 H); API MS  $m/z$  = 507  $[\text{C}_{28}\text{H}_{22}\text{N}_6\text{O}_4+\text{H}]^+$ .

Intermediate 87;


$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  8.69 (s, 2 H), 8.24 (broad s, 1 H), 7.98 (d, 1 H), 8.48 (t, 1 H), 7.42-7.25 (m, 8 H), 7.17 (d, 1 H), 6.60 (d, 1 H), 5.47 (d, 1 H), 5.18 (s, 2 H); CIMS  $m/z$  = 387  $[\text{C}_{22}\text{H}_{18}\text{N}_4\text{O}_3+\text{H}]^+$ .

Intermediate 88;

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  8.70 (d, 2 H), 8.25 (d, 1 H), 7.62 (t, 1 H), 7.40-7.24 (m, 9 H), 6.68 (d, 1 H), 5.33 (d, 1 H), 5.14 (s, 2 H), 3.46 (s, 3 H).

Scheme 9

30



Step 1: Preparation of 2-Amino-3'-methoxybenzophenone **93**;

Following the literature precedent: Walsh, D. A.

5 Synthesis, 1980, 677-688.

To a solution of bromoanisole **91** (1.38 mL, 10.7 mmol) in THF (5 mL) at 78°C and under nitrogen atmosphere, was added 1.2 M  $t\text{-BuLi}$  in pentane (18.2 mL, 21.9 mmol) dropwise. After stirring for 20 min, anthranilonitrile **92** (0.63 g, 5.3 mmol) was added. The reaction was warmed to rt and stirred overnight. The reaction was cooled to 0°C and quenched with 3N HCl. After stirring 30 min at rt, the acidic extracts were washed with EtOAc (3 X 12 mL). The aqueous layer was made basic (pH > 10) using 6N NaOH and

extracted with  $\text{CH}_2\text{Cl}_2$ . The organic extracts were washed with sat. NaCl, dried over  $\text{NaSO}_4$ , filtered, and concentrated to give a crude mixture containing **93**. The crude oil was purified by chromatography on silica to  
5 provide pure **93** (0.17 g, 14%) as a yellow oil;  $^1\text{H}$  NMR (500 MHz,  $\text{DMSO}-d_6$ )  $\delta$  7.40-7.43 (m, 1 H), 7.27-7.30 (m, 1 H), 7.07-7.17 (m, 4 H), 6.86-6.88 (m, 1 H), 6.49-6.52 (m, 1 H), 3.81 (s, 3 H); CI MS  $m/z$  = 228  $[\text{C}_{14}\text{H}_{13}\text{NO}_2+\text{H}]^+$ .

10 Step 2: Preparation of **95**;

Based on the literature procedures; Semple, G.; Ryder, H.; Ohta, M.; Satoh, M. *Syn. Comm.*, **1996**, 26(4), 721-727. Semple, G.; Ryder, H.; Rooker, D.P.; Batt, A.R.; Kendrick, D.A.; Szelke, M.; Ohta, M.; Satoh, M.; Nishida, A.;  
15 Akuzawa, S.; Miyata, K. *J. Med. Chem.*, **1997**, 40, 331-341.

To a solution of benzotriazole acid **94** (16.47 g, 50.5 mmol) in THF (101 mL) at  $0^\circ\text{C}$  under a nitrogen atmosphere was added oxalyl chloride (4.84 mL, 55.5 mmol) and a  
20 catalytic amount of DMF (1.0 mL). After stirring for 1 h, a solution of **93** (11.47 g, 50.5 mmol) and N-methylmorpholine (12.25 mL, 111 mmol) in THF (51 mL) was added dropwise via dropping funnel. The reaction was allowed to warm to rt and stirred overnight. The solids  
25 were removed by filtration. The filtrate was diluted with MeOH (62 mL), and  $\text{NH}_3$  gas was condensed into the solution for 15 min, the reaction vessel was sealed and stirred at rt for 2 h. After concentration, the crude oil was dissolved in EtOAc, washed with 3N NaOH (3 x 20 mL). The  
30 organic extracts were dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated. The resultant crude oil was dissolved in AcOH (121 mL), and  $\text{NH}_4\text{OAc}$  (7.17 g) was added. After stirring at rt overnight (16 h), the reaction was concentrated and diluted with water and made basic with 50%

NaOH. The basic aqueous solution was extracted with EtOAc (3 x 150 mL). The combined organic extracts were washed with sat. NaCl, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The crude product was purified by  
5 chromatography on silica to provide **95** (6.96 g, 33%) as a solid; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.50 (t, 1 H), 7.10-7.40 (m, 9 H), 7.05 (d, 1 H), 6.97 (dd, 1 H), 6.63 (d, 1 H), 5.30 (d, 1 H), 5.20 (s, 2 H), 3.77 (s, 3 H).


10 Step 3: Preparation of **96**;

To a solution of **95** (2.07 g, 5.4 mmol) and K<sub>2</sub>CO<sub>3</sub> (3.71 g, 26.8 mmol) in DMF (9.7 mL), was added MeI (0.50 mL, 8.1 mmol). After stirring 1 h at rt, the reaction was poured into water (100 mL) and stirred overnight. The solid was  
15 collected by filtration, rinsed with water and Et<sub>2</sub>O. This was dried in a vacuum oven at 55°C to provide **96** (1.5 g, 3.5 mmol) as a red solid; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.58 (t, 1 H), 7.19-7.42 (m, 9 H), 7.07 (d, 1 H), 7.00 (dd, 1 H), 6.69 (d, 1 H), 5.32 (d, 1 H), 5.15 (s, 2 H), 3.83 (s, 3 H), 3.45 (s, 3 H).  
20

Step 3: Preparation of **97**; Cbz deprotection to free amine.

To protected amine **96** (2.13 g, 4.9 mmol) was added 30% HBr in AcOH (50 mL). This was stirred at rt for 1 h. The  
25 reaction was poured into water (200 mL) and washed with EtOAc (200 mL). The aqueous layer was made basic with 50% NaOH and extracted with EtOAc (2 x 200 mL). The extracts were concentrated to give crude **97** (1.14 g, 78%) as a yellow solid; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.58 (t, 1 H),  
30 7.17-7.40 (m, 5 H), 7.12 (d, 1 H), 7.00 (dd, 1 H), 3.83 (s, 3 H), 3.48 (s, 3 H); CI MS m/z = 296 [C<sub>17</sub>H<sub>17</sub>N<sub>3</sub>O<sub>2</sub>+H]<sup>+</sup>.

Scheme 10



Step 1: Preparation of **102**;

- To a stirred mixture of p-anisidine (6 g, 48.7 mmol) in methylene chloride (170 mL) under nitrogen was added DMAP (5.0 mg, mmol) and triethylamine (13.9 mL, 99.7 mmol.) Ethyl chloroformate (4.85 mL, 50.7 mmol) was added and the reaction stirred for 2.5 hours. Dilute phosphoric acid (0.5 N) was added and the mixture stirred for 15 minutes.
- The aqueous layer was neutralized with sodium bicarbonate and then extracted with methylene chloride. The methylene chloride solution was dried over sodium sulfate, filtered and concentrated. Purification by column chromatography (gradient ethyl acetate/hexanes) provided **102** (7.2 g, 75%) as a solid: <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.27 (d, 2 H), 6.82

(d, 2 H), 6.45 (broad s, 1 H), 4.20 (q, 2 H), 3.77 (s, 3 H), 1.31 (t, 3 H).

Step 2: Preparation of **103**;

5 To a stirred suspension of lithium aluminum hydride (1.8 g, 47.4 mmol) in THF (95 mL) was added **102** (3.0 g, 15.4 mmol.) The reaction was heated at reflux for 2 hours. The reaction was cooled and quenched with ethyl acetate (150 mL). Solid sodium sulfate was added and the mixture  
10 filtered. The clear solution was evaporated and dried to give **103** (2.1 g, 100%); <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 6.79 (d, 2 H), 6.56 (d, 2 H), 3.74 (s, 3 H), 2.78 (s, 3 H).

Step 3: Preparation of **105**;

15 Based on the following literature references:  
Sugasawa, T., Toyoda, T.; Adachi, M.; Sasakura, K. *J. Chem. Soc.*, **1978**, 100, 4842-4852. Houpis, I.N.; Moline, A.; Douglas, A.W.; Xavier, L.; Lynch, J.; Volante, R.P.; Reider, P.J. *Tet. Lett.*, **1994**, 35, 6811-6814. Adachi, M.;  
20 Sasakura, K.; Sugasawa, T. *Chem. Pharm. Bull.*, **1985**, 33, 1826-1835. Sugasawa, T.; Adachi, M.; Sasakura, K.; Kitagawa, A. *J. Org. Chem.*, **1979**, 44, 578-586.

To a stirred solution of boron trichloride (1 M in  
25 CH<sub>2</sub>Cl<sub>2</sub>, 29.9 mL) at 0°C was added toluene (15 mL.) A solution of **103** (3.42 g, 24.9 mmol) in toluene (40 mL) was added dropwise over 0.5 h. The reaction was stirred at 0°C for 20 minutes and then heated under reflux for 1 h. Anthranilonitrile (25.7 g, 249 mmol) was added and the  
30 reaction heated under reflux for 6.5 h. 2N HCl (1.5 mL) was added and the reaction heated at 80°C for 0.5 h. Water was added and the reaction was neutralized with solid sodium carbonate and extracted with methylene chloride. The combined organics were dried over magnesium sulfate,  
35 filtered and concentrated. Purification by column

chromatography gave **105** (1.3 g, 33%) as a yellow solid. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 9.98 (broad s, 1 H), 7.61 (d, 2 H), 7.55-7.4 (m, 3 H), 7.1 (d of d, 1 H), 7.02 (d, 1 H), 6.72 (d, 1 H), 3.64 (s, 3 H), 2.93 (d, 3 H).

5

Step 4: Preparation of **107**;

Based on the following literature references: Semple, G.; Ryder, H.; Ohta, M.; Satoh, M. *Syn. Comm.*, **1996**, 26(4), 721-727. Semple, G.; Ryder, H.; Rooker, D.P.; Batt, A.R.; Kendrick, D.A.; Szelke, M.; Ohta, M.; Satoh, M.; Nishida, A.; Akuzawa, S.; Miyata, K. *J. Med. Chem.*, **1997**, 40, 331-341.

To a stirred solution of benzotriazole acid **106** in THF (3 mL) at 0°C was added oxalyl chloride (0.95 mL, 1.1 mmol) and a catalytic amount of DMF and the reaction stirred 1 h. A solution of **105** (240 mg, 1 mmol) and N-methyl morpholine (0.24 mL, 2.2 mmol) in THF 1.5 mL) was added dropwise. The reaction was stirred overnight at room temperature. The solution was filtered and concentrated. Methanol (3 mL) was added ammonia gas was bubbled into the reaction for 25 minutes. The reaction was stirred at room temperature for 1.5 h and then evaporated in vacuo and diluted with ethyl acetate. The solution was washed with 1 N NaOH. The aqueous layer was extracted with ethyl acetate and the combined organics were dried over magnesium sulfate and concentrated. The residue was dissolved in acetic acid (2.4 mL) and ammonium acetate (140 mg) was added and the reaction stirred for 16 h. The reaction was concentrated in vacuo and diluted with 1 N NaOH and extracted with ethyl acetate. The organic layer was dried over sodium sulfate, filtered and concentrated. The residue was chromatographed on silica gel (ethyl acetate/hexanes) to give **107** (98 mg, 45%) as a yellow solid); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.63 (d, 2 H), 7.45 (t, 1 H), 7.34-7.23 (m, 8 H), 7.13 (d, 1 H),

35

6.79 (s, 1 H), 6.67 (d, 1 H), 5.34 (d, 1 H), 5.14 (q, 2 H),  
3.73 (s, 3 H), 3.40 (s, 3 H); API MS  $m/z$  = 430  
[C<sub>25</sub>H<sub>23</sub>N<sub>3</sub>O<sub>4</sub>+H]<sup>+</sup>; mp. 71-73°C; IR (thin film) 3410, 1724,  
1678, 1498, 1289, 1231, 1039, 699 cm<sup>-1</sup>.


5

### Examples

Chemical abbreviations used in the Examples are defined as follows: "DMPU" for 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidone, "TBTU" for O-(1H-benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate, "BOP" for benzotriazol-1-yloxytris-(dimethylamino)-phosphonium hexafluorophosphate, and "HATU" for O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate. "HPLC" is an abbreviation used herein for high pressure  
15 liquid chromatography. Reverse-phase HPLC was carried out using a Vydac C-18 column with gradient elution from 10% to 100 % buffer B in buffer A (buffer A: water containing 0.1% trifluoroacetic acid, buffer B: 10% water, 90% acetonitrile containing 0.1% trifluoroacetic acid).  
20 Alternatively, reverse-phase HPLC was carried out using a Vydac C-18 column with gradient elution from 10% to 90 % acetonitrile in water.


### Example 1

25 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide



30




Step 1

5 A solution of tert-butyl succinate ester (9) (1.0 eq.) in DMF (0.25 M) under N<sub>2</sub> at 0°C was added HATU (1.1 eq.), then Hunig's base (4.0eq.). The mixture was stirred at 0°C for 10 mins. A solution of 1,3-dihydro-1-methyl-3-amino-5-phenyl-2H-1,4-benzodiazepin-2-one (30) in DMF (0.8 M) (1.0 eq.) was added to this solution. The reaction mixture was stirred overnight at room temperature and then transferred to a separatory funnel containing water. 30% n-Hexane in ethyl acetate was added which gave a clear organic layer. The aqueous solution was extracted twice with 30% n-hexane in ethyl acetate. The combined organic layers were washed with water and brine, dried over magnesium sulfate, and concentrated in vacuo. The residue was purified by chromatography on flash grade silica gel using 20-30% ethyl acetate in n-hexane; preferably 30% ethyl acetate in n-hexane. Compound, 1', was isolated as an amorphous white solid (85%). R<sub>f</sub> = 0.25 (7:3 n-hexane:ethyl acetate).

<sup>1</sup>H-NMR:(CDCl<sub>3</sub>): δ7.61-7.21 (m, 10H); 5.77-5.73 (m, 1H); 5.57-5.54 (d, 1H); 5.20-4.97 (m, 2H); 3.47 (s, 3H); 2.63-2.33 (m, 4H); 1.80-1.76 (m, 2H); 1.47-1.46 (d, 9H); 1.43-1.11 (m, 1H); 1.01-0.86 (m, 6H).


MS: C<sub>31</sub>H<sub>39</sub>N<sub>3</sub>O<sub>4</sub> (M+H) 518.3 (M+Na) 540.3.

30 Step 2



A solution of (1') in 50% TFA in methylene chloride (0.15M) was stirred at room temperature overnight. The solution  
 5 was concentrated in vacuo, washed and concentrated four times with toluene in vacuo to give compound (1'') as an amorphous solid (95%). Rf = 0.64 (9.5 : 0.5 methylene chloride : methanol). MS: C<sub>27</sub>H<sub>31</sub>N<sub>3</sub>O<sub>4</sub> (M+H) 462.

### 10 Step 3



To a solution of (1'') (1.0 eq.) in DMF (0.25 M) under N<sub>2</sub> at 0°C was added HATU (1.1 eq.), and then Hunig's base  
 15 (4.0eq.). The mixture was stirred at 0°C for 10 mins, and then anhydrous ammonia bubbled through the solution for two minutes. The reaction mixture was stirred overnight at room temperature and then transferred to a separatory funnel containing water and diluted with 30% n-hexane in ethyl.  
 20 The aqueous solution was extracted twice with 30% n-hexane in ethyl acetate. The combined organic layers were washed with water and brine, dried over magnesium sulfate, and concentrated in vacuo. The residue was purified by chromatography on flash grade silica gel using 4 % methanol  
 25 in methylene chloride. The title compound, Example 1, was

isolated as an amorphous white solid (87%).  $R_f = 0.43$  (9:1 methylene chloride : methanol).

$^1\text{H NMR}$ : ( $\text{CDCl}_3$ ):  $\delta$  7.63-7.22 (m, 10H); 6.25-6.13 (d, 1H);

5 5.88-5.73 (m, 1H); 5.53-5.51 (dd, 1H); 5.44-5.41 (d, 1H);  
5.22-5.04 (m, 2H); 3.47-3.46 (d, 3H); 2.74-2.31 (m, 4H);  
1.81-1.61 (m, 2H); 1.34-1.22 (m, 1H); 0.99-0.87 (m, 6H).


MS:  $\text{C}_{27}\text{H}_{32}\text{N}_4\text{O}_3$  (M+H) 461.

10

#### Example 2

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-N4-[3,5-difluorobenzyl]-butanediamide


15



Following a procedure analogous to the preparation of Example 1 using 3,5-difluorobenzylamine in place of ammonia  
20 in Step 3, the title compound was prepared. MS (M+H)<sup>+</sup> = 587.

#### Example 3

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-propyl-  
25 butanediamide




Following a procedure analogous to the preparation of Example 1 using succinate ester (10) (Scheme 2) in place of succinate ester (9) in Step 1, as well as reagents known to one skilled in the art, the title compound was prepared.

5 MS (M+H)<sup>+</sup> = 463.0.

### Example 4a

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(2-  
fluorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-  
10 methylpropyl)-3-allyl-butanediamide



Following a procedure analogous to the preparation of  
15 Example 1 using 1,3-dihydro-1-methyl-3-amino-5-(2-fluorophenyl)-2H-1,4-benzodiazepine-2-one in place of (30) in Step 1, the mixture of benzodiazepine diastereomers of the title compound was prepared. The isomers were separated using reverse phase chromatography (C-18 column, isocratic  
20 eluent composed of 65% 90:10 water:acetonitrile and 35% 10:90 water:acetonitrile). The first eluting isomer was assigned the (S)-benzodiazepine configuration using the H-NMR method described for Example 12a. The required aminobenzodiazepine was prepared using the method of  
25 Sherrill and Sugg (J. Org. Chem. 1995, 60, 730-734). MS (M+H)<sup>+</sup> = 479.4; (M+Na)<sup>+</sup> = 501.4.


### Example 4b

(2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-(2-  
30 fluorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-  
methylpropyl)-3-allyl-butanediamide

Following the procedure of Example 4a, the second eluting isomer was isolated and assigned the (R)-benzodiazepine configuration using the H-NMR method described for Example 12b. MS (M+H)<sup>+</sup> = 479.4; (M+Na)<sup>+</sup> = 501.4.

### Example 5

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide



Following a procedure analogous to the preparation of Example 1 using (S) isomer of benzodiazepinone (30) in Step 1 in place of racemic benzodiazepinone (30), and reducing the amount of Hunig's base in Step 1 to 2.2 eq., the title compound was prepared.


<sup>1</sup>H-NMR for product of Step 1: (CDCl<sub>3</sub>) : 7.61-7.21 (m, 10H); 5.77-5.73 (m, 1H); 5.57-5.54 (d, 1H); 5.20-5.00 (m, 2H); 3.46 (s, 3H); 2.63-2.33 (m, 4H); 1.80-1.76 (m, 2H); 1.46 (s, 9H); 1.43-1.11 (m, 1H); 0.97-0.95 (d, 3H); 0.88-0.85 (d, 3H).

<sup>1</sup>H-NMR for product of Step 3: (CDCl<sub>3</sub>) : 7.65-7.00 (m, 10H); 6.10-6.00 (s, 1H); 5.85-5.65 (m, 1H); 5.54-5.52 (d, 1H); 5.40-5.30 (s, 1H); 5.23-5.00 (m, 2H); 3.47 (s, 3H); 2.80-2.20 (m, 4H); 1.85-1.60 (m, 2H); 1.40-1.20 (m, 1H),

0.98-0.95 (d, 3H); 0.90-0.87 (d, 3H). MS (M+H)<sup>+</sup> = 461.3;  
MS (M+H)<sup>+</sup> = 483.3.

**Example 6**

5 (2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-  
1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-  
butanediamide



10

Using the method of Example 5, but replacing the (S) isomer of benzodiazepinone (30) with (R) isomer of benzodiazepinone (30) in Step 1, the title compound was prepared. Alternatively, the (R)- and (S)-benzodiazepine  
15 diastereomers may be separated using a Chiralpack AD column, eluting with 0.1% diethylamine in 1:1 n-hexane/i-propanol. Under these conditions, the compound of Example 6 [(R)-benzodiazepine diastereomer] elutes first.


20 <sup>1</sup>H-NMR: (CDCl<sub>3</sub>) : 7.65-7.00 (m, 10H); 6.10-6.00 (s, 1H);  
5.85-5.65 (m, 1H); 5.54-5.52 (d, 1H); 5.40-5.30 (s, 1H);  
5.23-5.00 (m, 2H); 3.47 (s, 3H); 2.80- 2.20 (m, 4H); 1.85-  
1.60 (m, 2H); 1.40-1.20 (m, 1H), 0.99-0.95 (apparent t,  
6H). MS (M+H)<sup>+</sup> = 461.3; MS (M+Na)<sup>+</sup> = 483.3.

25

**Example 7**

(2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-  
1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-propyl-  
30 butanediamide


30



The product of Example 6 was hydrogenated by dissolution in methanol (0.25M), addition of an equal amount by weight of Pearlman's catalyst [ $\text{Pd}(\text{OH})_2 / \text{C}$  (20% Wt, Water %  $\leq 50$ )] and 30 eq. of 1,4-cyclohexadiene at room temperature. The solution was heated to reflux for 1hr, then cooled to room temperature. Removal of volatiles and purification by flash chromatography using 4% methanol in  $\text{CH}_2\text{Cl}_2$  gave the title compound. MS  $(\text{M}+\text{H})^+ = 463.3$ ; MS  $(\text{M}+\text{Na})^+ = 485.3$ .

#### Example 8


(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-propylbutanediamide



Using the product of Example 5 as starting material, and following the method of Example 7, the title compound was prepared. MS  $(\text{M}+\text{H})^+ = 463.3$ ; MS  $(\text{M}+\text{Na})^+ = 485.3$ .

#### Example 9a

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-methyl-3-allylbutanediamide




Following a procedure analogous to the preparation of Example 5, coupling the appropriate succinate ester, prepared using the method of Becket et al. (Synlett, 1993, pp. 137-138) Scheme 1), in place of (9), with (S) isomer of benzodiazepinone-(30) in Step 1, the title compound was prepared. MS (M+H)<sup>+</sup> = 419.1; (M+Na)<sup>+</sup> = 441.1.

10

**Example 10**

(2R) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-methyl-butanediamide



15


Following a procedure analogous to the preparation of Example 1 using the appropriate succinate ester (prepared by methods disclosed in Becket et al. (Synlett, 1993, pp. 137-138) and shown in Scheme 1), in place of ester (9) in Step 1, the title compound was prepared. MS (M+H)<sup>+</sup> = 379.0

**Example 11**

(2R,3S) N1-[1,3-dihydro-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide

25






Following a procedure analogous to the preparation of  
 Example 1 using 1,3-dihydro-3-amino-5-phenyl-2H-1,4-  
 benzodiazepine-2-one in place of (30) in Step 1, the title  
 5 compound was prepared. MS (M+H)<sup>+</sup> = 447.2.

#### Example 12


(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-7-chloro-  
 2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-  
 10 butanediamide



Following a procedure analogous to the preparation of  
 Example 1 using 1,3-dihydro-1-methyl-3-amino-5-phenyl-7-  
 15 chloro-2H-1,4-benzodiazepine-2-one in place of (30) in Step  
 1, the title compound was prepared. MS (M+H)<sup>+</sup> = 495.0.

#### Example 12a


(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-7-  
 20 chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-  
 allyl-butanediamide



The product of Example 12 was separated into the (S)- and (R)-benzodiazepine diastereomers using a Chiralpack AD column, eluting with 1:1 n-hexane - i-propanol. The diastereomers were assigned on the basis of the elution order ((S)-diastereomer elutes second under these conditions), and the i-butyl signals in the H-NMR spectra, consistent with the elution order and H-NMR spectra of the products of Examples 5 and 6. In CDCl<sub>3</sub> the (S)-diastereomer shows two doublets (6H total) above 1 ppm and the (R)-diastereomer shows an apparent triplet (6H) above 1 ppm. MS (M+H)<sup>+</sup> = 495.2; MS (M+Na)<sup>+</sup> = 517.1.

#### Example 12b

(2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide




As described for Example 12a, the product of Example 12 was separated into the (S)- and (R)-benzodiazepine diastereomers using a Chiralpack AD column, eluting with 1:1 n-hexane - i-propanol, with the (R)-diastereomer eluting first under these conditions. In CDCl<sub>3</sub> the i-butyl signals for the (R)-diastereomer appear as an apparent

triplet (6H) above 1 ppm. MS (M+H)<sup>+</sup> = 495.2; MS (M+Na)<sup>+</sup> = 517.1.

### Example 13


- 5 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(2-fluorophenyl)-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide



- 10 Following a procedure analogous to the preparation of Example 1 using 1,3-dihydro-1-methyl-3-amino-5-(2-fluorophenyl)-7-chloro-2H-1,4-benzodiazepine-2-one, prepared by methods known by one skilled in the art, in place of (30) in Step 1, the title compound was prepared.
- 15 MS (M+H)<sup>+</sup> = 513.3; MS (M+Na)<sup>+</sup> = 535.2.

### Example 13a

- (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(2-fluorophenyl)-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide
- 20




- Following a procedure analogous to the preparation of
- 25 Examples 12a and 12b, the (S)-benzodiazepine diastereomer

of Example 13 was isolated. MS (M+H)<sup>+</sup> = 513.3; MS (M+Na)<sup>+</sup> = 535.2.

### Example 13b

5 (2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-(2-fluorophenyl)-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide




10 Following a procedure analogous to the preparation of Examples 12a and 12b, the (R)-benzodiazepine diastereomer of Example 13 was isolated. MS (M+H)<sup>+</sup> = 513.3; MS (M+Na)<sup>+</sup> = 535.2.

15

### Example 14

(2S,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide

20




Following a procedure analogous to the preparation of Example 1 using the appropriate syn-succinate ester (prepared by methods disclosed in Becket et al. (Synlett, 1993, pp. 137-138) and shown in Scheme 1), in place of

25

ester (9) in Step 1, the title compound was prepared. MS  $(M+H)^+ = 461.4$ ;  $(M+Na)^+ = 483.4$ .

#### Example 15

5 (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-propyl-butanediamide




10 Using the product of Example 12a as starting material, and following the reduction procedure of Example 7, the title compound was prepared. MS  $(M+H)^+ = 497.2$ ; MS  $(M+Na)^+ = 519.1$

15

#### Example 16

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(2-fluorophenyl)-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-propyl-butanediamide




20

Using the product of Example 13a as starting material, and following the reduction procedure of Example 7, the title compound was prepared. MS  $(M+H)^+ = 515.3$ ; MS  $(M+Na)^+ = 537.2$

25

#### Example 17

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(4-fluorophenyl)-  
2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-  
butanediamide




5

Following a procedure analogous to the preparation of  
Example 1 using 1,3-dihydro-1-methyl-3-amino-5-(4-  
fluorophenyl)-2H-1,4-benzodiazepine-2-one, prepared by  
methods known to one skilled in the art, in place of (30)

10 in Step 1, the title compound was prepared. MS (M+H)<sup>+</sup> =  
479.3.

#### Example 17a


(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-  
15 fluorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-  
methylpropyl)-3-allyl-butanediamide



20 Following a procedure analogous to the preparation of  
Examples 12a and 12b, the (S)-benzodiazepine diastereomer  
of Example 13 was isolated. MS (M+H)<sup>+</sup> = 479.3; MS (M+Na)<sup>+</sup>  
= 501.2.

#### Example 17b

(2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-(4-fluorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide



5


Following a procedure analogous to the preparation of Examples 12a and 12b, the (R)-benzodiazepine diastereomer of Example 13 was isolated. MS (M+H)<sup>+</sup> = 479.3; MS (M+Na)<sup>+</sup> = 501.2.

10

#### Example 18

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(pyrid-2-yl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide

15



Following a procedure analogous to the preparation of Example 1 using 1,3-dihydro-1-methyl-3-amino-5-(pyrid-2-yl)-2H-1,4-benzodiazepine-2-one, prepared by methods known to one skilled in the art, in place of (30) in Step 1, the title compound was prepared. MS (2M+Na)<sup>+</sup> = 945.6.


20

#### Example 19

.38618

PCT/US99/0815

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(N-morpholino)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide



5

Following a procedure analogous to the preparation of Example 1 using 1,3-dihydro-1-methyl-3-amino-5-(N-morpholino)-2H-1,4-benzodiazepine-2-one, prepared by methods known to one skilled in the art, in place of (30) in Step 1, the title compound was prepared. Mp 145-150°C.


10

MS (M+H)<sup>+</sup> = 470.

15

#### Example 20

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(N-morpholino)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide



20


Following a procedure analogous to the preparation of Example 1 using (dimethylamino)acetic acid, prepared by methods known to one skilled in the art, in place of (30) in Step 1, the title compound was prepared.



**Exempl 21**

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(N-methyl-N-phenylamino)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide


5



Following a procedure analogous to the preparation of Example 1 using 1,3-dihydro-1-methyl-3-amino-5-(N-methyl-N-phenylamino)-2H-1,4-benzodiazepine-2-one, prepared by  
 10 methods known to one skilled in the art, in place of (30) in Step 1, the title compound was prepared. melting point 266.6-267.8°C; MS (M+H)<sup>+</sup> = 490.

**Example 22**


15 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(N-piperidinyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide



20 Following a procedure analogous to the preparation of Example 1 using 1,3-dihydro-1-methyl-3-amino-5-(N-piperidinyl)-2H-1,4-benzodiazepine-2-one, prepared by methods known to one skilled in the art, in place of (30) in Step 1, the title compound was prepared. melting point  
 25 231-238°C; MS (M+H)<sup>+</sup> = 468.

**Example 23**

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(N-homopiperidinyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide




5

Following a procedure analogous to the preparation of Example 1 using 1,3-dihydro-1-methyl-3-amino-5-(N-homopiperidinyl)-2H-1,4-benzodiazepine-2-one, prepared by methods known to one skilled in the art, in place of (30) in Step 1, the title compound was prepared. melting point 232-237°C; MS (M+H)<sup>+</sup> = 482.

#### Example 24

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(3-methoxyphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide



Following a procedure analogous to the preparation of Example 1 using 1,3-dihydro-1-methyl-3-amino-5-(3-methoxyphenyl)-2H-1,4-benzodiazepine-2-one, prepared by methods known to one skilled in the art, in place of (30) in Step 1, the title compound was prepared. melting point 225-230°C; MS (M+H)<sup>+</sup> = 491.

Step 1: Preparation of 24';

This was prepared using the standard HATU coupling conditions with **97** (1.03 g, 3.5 mmol) to give **24'** (0.48 g, 25%) as a colorless oil (mixture of diastereomers):  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  7.63 (t, 1 H), 7.33-7.38 (m, 2 H), 7.24-7.30 (m, 2 H), 7.05 (d, 1 H), 7.00 (dd, 1 H), 5.65-5.8 (m, 1 H), 5.57 (d, 1 H), 4.92-5.20 (m, 2 H), 3.80 (s, 3 H), 3.77 (s, 3 H), 3.48 (s, 3 H), 2.27-2.67 (m, 3 H), 1.75-1.80 (m, 1 H), 1.60-1.63 (m, 1 H), 1.48 (s, 9 H), 1.46 (s, 9 H), 1.22-1.26 (m, 1 H), 0.88-1.00 (m, 3 H), 0.80 (d, 3 H).

10

Step 2: Preparation of **24''** ;

This was prepared using the standard *t*-Bu ester deprotection with TFA. The acid intermediate **24''** is a mixture of diastereomers:  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  9.75 (bs, 1 H), 7.75-7.85 (m, 1 H), 7.00-7.51 (m, 7 H), 5.70-5.80 (m, 2 H), 5.05-5.15 (m, 2 H), 3.80 (s, 3 H), 3.53, 3.50 (s, 3 H), 5.74-2.93 (m, 1 H), 2.25-2.53 (m, 1 H), 1.53-1.80 (m, 1 H), 1.25-1.30 (m, 1 H), .93 (d, 3 H), 0.89 (d, 3 H); API MS  $m/z$  = 492 [ $\text{C}_{28}\text{H}_{33}\text{N}_3\text{O}_5+\text{H}$ ] $^+$ .

20

Step 3: Preparation of Example **24**;

This was prepared using the HATU mediated amide formation.


mp 225-230°C;  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  7.55-7.70 (m, 2 H), 7.27-7.33 (m, 3 H), 7.24 (s, 1 H), 7.0-7.07 (m, 2 H), 5.70-5.77 (m, 1 H), 5.45 (d, 1 H), 4.93-5.10 (m, 2 H), 3.78, 3.76 (s, 3 H), 3.48 (s, 3 H), 2.75-2.85 (m, 1 H), 2.27-2.50 (m, 2 H), 1.53-1.71 (m, 2 H), 1.10-1.25 (m, 1 H), 0.80-1.00 (m, 6 H); IR (KBr) 3423, 2956, 1655, 1601, 1326  $\text{cm}^{-1}$ ; API MS  $m/z$  = 491 [ $\text{C}_{28}\text{H}_{34}\text{N}_4\text{O}_4+\text{H}$ ] $^+$ ; Anal. Calcd. For

30

C<sub>28</sub>H<sub>34</sub>N<sub>4</sub>O<sub>4</sub>·0.5 H<sub>2</sub>O: C, 67.3; H, 7.06; N, 11.21. Found:  
C, 67.52; H, 6.85; N, 11.21.

**Example 25**

- 5 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(pyrid-4-yl)-2H-  
1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-  
butanediamide




- 10 Following a procedure analogous to the preparation of  
Example 1 using methods known to one skilled in the art,  
the title compound was prepared. MS (M+H)<sup>+</sup> = 462.

15

**Example 26**

- (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-7-methoxy-  
2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-  
butanediamide

20




- Following a procedure analogous to the preparation of  
Example 1 using reagents known to one skilled in the art,  
the title compound was prepared. m.pt. 244-255 °C; MS  
25 (M+H)<sup>+</sup> = 491.

**Example 27**

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(pyrid-3-yl)-2H-  
1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-  
butanediamide

5




Following a procedure analogous to the preparation of  
Example 1 using reagents known to one skilled in the art,  
the title compound was prepared. m.pt. 251-253°C; MS

10 (M+H)<sup>+</sup> = 462.**Example 28a**

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-  
1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-  
(cyclopropylmethyl)-butanediamide

15




A procedure analogous to the preparation of Example 5  
was followed. The appropriate succinate ester was prepared  
20 using the method of Becket et al. (Synlett, 1993, pp. 137-  
138), using cyclopropylmethyl iodide for reagent R<sup>3</sup>-X in  
Scheme 1. This succinate half-ester, a mixture of syn- and  
anti-isomers, was coupled to the (S) benzodiazepinone-(30)  
using the procedures of Example 1, Step 1. After Steps 2  
25 and 3, the mixture of butanediamide isomers was separated

using silica gel chromatography to provide Example 28a. MS  
(M+H)<sup>+</sup> = 475.

**Example 28b**

- 5 (2R,3R) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-  
1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-  
(cyclopropylmethyl)-butanediamide




- 10 From the product of Step 3 of Example 28a, the other  
succinate isomer, Example 28b, was isolated. Mp 146-148 °C;  
MS (M+H)<sup>+</sup> = 475.

15

**Example 29**

- (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(3-fluorophenyl)-  
2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-  
butanediamide




20

Following a procedure analogous to the preparation of  
Example 1 using 1,3-dihydro-1-methyl-3-amino-5-(3-  
fluorophenyl)-2H-1,4-benzodiazepine-2-one, prepared by  
methods known to one skilled in the art, in place of (30)

in Step 1, the title compound was prepared. MS (M+H)<sup>+</sup> = 479.3. ; (M+Na)<sup>+</sup> = 501.3.

**Example 29a**


- 5 (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(3-fluorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide



- 10 Following a procedure analogous to the preparation of Examples 12a and 12b, the (S)-benzodiazepine diastereomer of Example 29 was isolated. MS (M+H)<sup>+</sup> = 479.3.

**Example 29b**


- 15 (2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-(3-fluorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide



- 20 Following a procedure analogous to the preparation of Examples 12a and 12b, the (R)-benzodiazepine diastereomer of Example 29 was isolated. MS (M+H)<sup>+</sup> = 479.3.

**Example 30**

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-(3-buten-1-yl)-butanediamide




5

A procedure analogous to the preparation of Example 5 was followed. The appropriate succinate ester was prepared using the method of Becket et al. (Synlett, 1993, pp. 137-138), using 3-butenyl bromide for  $R^3-X$  in Scheme 1. The anti-succinate half-ester was coupled to (S)-isomer benzodiazepine (30) using the procedures of Step 1. After Steps 2 and 3, the title compound was isolated. MS  $(M+H)^+ = 475.3$ ;  $(M+Na)^+ = 497.3$ .

15

#### Example 31

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-(cyclopentylethyl)-butanediamide



20


A procedure analogous to the preparation of Example 5 was followed. The appropriate succinate ester was prepared using the method of Becket et al. (Synlett, 1993, pp. 137-138), using 2-(cyclopentyl)ethyl iodide for  $R^3-X$  in Scheme 1. The anti-succinate half-ester was coupled to (S)-isomer benzodiazepine (30) using the procedures of Step 1. After



Steps 2 and 3, the title compound was isolated. MS (M+H)<sup>+</sup> = 475; (M+Na)<sup>+</sup> = 497.2.

#### Example 32a

5 (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-(3-buten-1-yl)-butanediamide




10 A procedure analogous to the preparation of Example 5 was followed. The succinate ester of Example 30 was employed. The anti-succinate half-ester was coupled to (S)-1,3-dihydro-1-methyl-3-amino-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepine-2-one using  
15 the procedures of Step 1. The required (S)-3-aminobenzodiazepine was prepared by HBr deprotection of the corresponding benzyloxycarbonyl derivative, which was separated from the (R)-isomer using chiral chromatography. The racemic Cbz-aminobenzodiazepine was prepared using the  
20 method of Sherrill and Sugg (J. Org. Chem. 1995, 60, 730-734). After Steps 2 and 3, the title compound was isolated. The stereochemical assignment of Examples 32a and 32b was performed following the H-NMR method of Examples 12a and 12b. MS (M+H)<sup>+</sup> = 543.2; (M+Na)<sup>+</sup> = 565.3.

25

#### Example 32b

(2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-(3-buten-1-yl)-butanediamide

30




A procedure analogous to the preparation of Example 32a was followed using (R)-1,3-dihydro-1-methyl-3-amino-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepine-2-one in

5 Step 1. MS (M+H)<sup>+</sup> = 543.2; (M+Na)<sup>+</sup> = 565.3.

### Example 33

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide

10




Following a procedure analogous to the preparation of Example 1 using 1,3-dihydro-1-methyl-3-amino-5-(4-

15 fluoromethylphenyl)-2H-1,4-benzodiazepine-2-one, prepared by methods known to one skilled in the art, in place of (30) in Step 1, the title compound was prepared. MS (M+H)<sup>+</sup> = 529.2.

20

### Example 33a

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide




Following a procedure analogous to the preparation of Examples 12a and 12b, the (S)-benzodiazepine diastereomer of Example 33 was isolated. MS (M+H)<sup>+</sup> = 529.2.

5

#### Example 33b

(2R,3S) N1-[(3R)-1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide

10




Following a procedure analogous to the preparation of Examples 12a and 12b, the (S)-benzodiazepine diastereomer of Example 33 was isolated. MS (M+H)<sup>+</sup> = 529.2; (M+Na)<sup>+</sup> =

15 551.2.

#### Example 34

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3,3-diallyl-butanediamide

20




A procedure analogous to the preparation of Example 5 was followed. The diallyl succinate ester was prepared using the method of Curtin et al. (Biorg. Med. Chem. Lett., 8, 1998, 1443-8). The succinate half-ester was coupled to the (S) isomer of benzodiazepinone (30) using the procedures of Step 1. After Steps 2 and 3, the title compound was isolated. MS (M+Na)<sup>+</sup> = 523.2.

10

**Example 35a**

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-n-butyl-butanediamide


15



Following the reduction procedure of Example 7, the product of Example 32a was converted to the title compound. MS (M+H)<sup>+</sup> = 545.2; (M+Na)<sup>+</sup> = 567.3.

**Example 36**

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-propyl-butanediamide




Following the reduction procedure of Example 7, the product of Example 33a was converted to the title compound.

MS (M+H)<sup>+</sup> = 531.2; (M+Na)<sup>+</sup> = 553.2.

#### Example 37

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-chlorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-(3-buten-1-yl)-butanediamide




A procedure analogous to the preparation of Example 5<sup>1</sup> was followed. The succinate ester of Example 30 was employed and was coupled to (S)-2,3-dihydro-1-methyl-3-amino-5-(4-chlorophenyl)-1H-1,4-benzodiazepine-2-one using the procedures of Step 1. The required (S)-3-aminobenzodiazepine was prepared by HBr deprotection of the corresponding benzyloxycarbonyl derivative, which was separated from the (R)-isomer using chiral chromatography.

The racemic Cbz-aminobenzodiazepine was prepared using the

method of Sherrill and Sugg (J. Org. Chem. 1995, 60, 730-734). After Steps 2 and 3, the title compound was isolated. The stereochemical assignment was performed following the H-NMR method of Example 12a. MS (M+H)<sup>+</sup> =  
5 509.2.

**Example 38**


(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-chlorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-n-butyl-butanediamide  
10



The product of Example 37 was taken up in ethanol  
15 (0.25M) together with 10% by weight Wilkinson's catalyst (chlorotris(triphenylphosphine)rhodium(I)). The mixture was shaken under 50 psi hydrogen overnight. The reaction mixture was filtered through a layer of celite and the solvent was evaporated. The residue was purified by flash  
20 chromatography on silica gel using 4% methanol in CH<sub>2</sub>Cl<sub>2</sub> to provide the title compound. MS (M+H)<sup>+</sup> = 511.2; (M+Na)<sup>+</sup> = 533.2.

**Example 39**


25 (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-N4-[benzyl]-butanediamide



Following a procedure analogous to the preparation of  
 Example 5 using benzylamine in place of ammonia in Step 3,  
 the title compound was prepared. MS (M+H)<sup>+</sup> = 551.2; (M+Na)<sup>+</sup>  
 5 = 565.3.

#### Example 40


(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-methyl-2H-1,4-  
 benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-  
 10 butanediamide



Following a procedure analogous to the preparation of  
 15 Example 1 using 1,3-dihydro-1-methyl-3-amino-5-methyl-2H-  
 1,4-benzodiazepine-2-one in place of (30) in Step 1, the  
 title compound was prepared. MS (M+H)<sup>+</sup> = 399.1.

#### Example 41

20 (2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-n-butyl-2H-1,4-  
 benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-  
 butanediamide




Following a procedure analogous to the preparation of Example 1 using 1,3-dihydro-1-methyl-3-amino-5-(n-butyl)-2H-1,4-benzodiazepine-2-one in place of (30) in Step 1, the title compound was prepared. The benzodiazepine diastereomers, Example 41a (3S) isomer and 41b (3R) isomer, were separated by silica gel chromatography. MS (M+H)<sup>+</sup> = 441.2.

10

**Example 42**

(2R,3S) N1-[1,3-dihydro-1-methyl-2-oxo-5-(2-methylpropyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide



15

Following a procedure analogous to the preparation of Example 1 using 1,3-dihydro-1-methyl-3-amino-5-(i-butyl)-2H-1,4-benzodiazepine-2-one in place of (30) in Step 1, the title compound was prepared. MS (M+H)<sup>+</sup> = 441.2.


20

**Example 43a**

(2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-(4-chlorophenyl)-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide

25






Following a procedure analogous to the preparation of Example 1 using 1,3-dihydro-1-methyl-3-amino-5-(4-chlorophenyl)-2H-1,4-benzodiazepine-2-one in place of (30) in Step 1, the mixture of benzodiazepine diastereomers of the title compound was prepared. The isomers were separated using reverse phase chromatography (C-18 column, isocratic eluent composed of 65% 90:10 water:acetonitrile and 35% 10:90 water:acetonitrile). The first eluting isomer was assigned the (S)-benzodiazepine configuration using the H-NMR method described for Example 12a. The required aminobenzodiazepine was prepared using the method of Sherrill and Sugg (J. Org. Chem. 1995, 60, 730-734). MS (M+H)<sup>+</sup> = 495.

#### Example 44a

(2R,3S) N1-[(3S)-1,3-dihydro-1-ethyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide



Following a procedure analogous to the preparation of Example 1 using 1,3-dihydro-1-ethyl-3-amino-5-phenyl-2H-


1,4-benzodiazepine-2-one in place of (30) in Step 1, the title compound was prepared. The required aminobenzodiazepine was prepared using the method of Sherrill and Sugg (J. Org. Chem. 1995, 60, 730-734). The benzodiazepine diastereomers of the product were separated using reverse phase HPLC as described for Example 43a. The first eluting isomer was assigned the (S)-benzodiazepine configuration using the H-NMR method described for Example 12a. MS (M+H)<sup>+</sup> = 475.

10

**Example 45a**

(2R,3S) N1-[(3S)-1,3-dihydro-1-propyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide

15



Following a procedure analogous to the preparation of Example 1 using 1,3-dihydro-1-(n-propyl)-3-amino-5-phenyl-2H-1,4-benzodiazepine-2-one in place of (30) in Step 1, the title compound was prepared as a mixture of benzodiazepine diastereomers, which were separated using reverse phase HPLC as described for Example 43a. The first eluting isomer was assigned the (S)-benzodiazepine configuration using the H-NMR method described for Example 12a. The required aminobenzodiazepine was prepared using the method of Sherrill and Sugg (J. Org. Chem. 1995, 60, 730-734). MS (M+H)<sup>+</sup> = 489.

30

**Example 45b**

(2R,3S) N1-[(3R)-1,3-dihydro-1-propyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide

5


Following the procedure of Example 45a, the second eluting isomer was isolated and assigned the (R)-benzodiazepine configuration using the H-NMR method described for Example 12b. MS (M+H)<sup>+</sup> = 489.

10

**Example 46**

(2R,3S) N1-[1,3-dihydro-1-(isopropyl)-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide

15



Following a procedure analogous to the preparation of Example 1 using 1,3-dihydro-1-(i-propyl)-3-amino-5-phenyl-2H-1,4-benzodiazepine-2-one in place of (30) in Step 1, the title compound was prepared. The required aminobenzodiazepine was prepared using the method of Sherrill and Sugg (J. Org. Chem. 1995, 60, 730-734). MS (M+H)<sup>+</sup> = 489.


25

**Example 47**

(2R,3S) N1-[6,7-dihydro-5-methyl-6-oxo-5H-dibenz[b,d]azepin-7-yl]-2-(2-methylpropyl)-3-allyl-butanediamide

30


Step 1: Preparation of Compound 47';



To a solution of amine **31**, (for preparation see Audia, J.E., PCT patent application WO 99/32453) (0.42 g, 1.76 mmol), succinate **9**, (0.52 g, 1.94 mmol), and DMF (50 mL) at 0 °C was added HATU (0.87 g, 2.29 mmol), and finally DIPEA (1.23 mL, 7.05 mmol). The solution was allowed to warm to room temperature while stirring for 18 h. The DMF was removed under reduced pressure and the residue was dissolved in ethyl acetate (200 mL), washed with 0.1 N HCl (2 x 100 mL), 5% aqueous NaHCO<sub>3</sub> (2 x 100 mL), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, and concentrated to yield a brown oil. The crude material was further purified via silica gel column chromatography (1% MeOH, 99% CH<sub>2</sub>Cl<sub>2</sub>) to yield **47'** (0.61g, 70.5%) as a pale yellow solid.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 0.87-1.42 (m, 9 H), 1.48 (d, 9 H), 2.21-2.79 (m, 4 H), 3.38 (s, 3 H), 4.92-5.11 (m, 3 H), 5.41 (d, 1 H), 5.72 (m, 1 H), 7.29-7.65 (m, 8 H). CI MS *m/z* = 491 [C<sub>30</sub>H<sub>38</sub>N<sub>2</sub>O<sub>4</sub>+H]<sup>+</sup>.

Step 2: Preparation of Compound **47''**;




t-Butyl ester **47'** (0.61 g, 1.24 mmol) was dissolved in TFA (15 mL) and stirred for 4 h at room temperature under

N<sub>2</sub>. The TFA was removed under reduced pressure, the residue was dissolved in ethyl acetate (50 mL), washed with a saturated solution of aqueous NaHCO<sub>3</sub> (2 x 50 mL), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, and concentrated to yield 47" (0.38 g, 71%) as a pale yellow foam.

<sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 0.8-1.82 (m, 9 H), 2.25-2.55 (m, 4 H), 3.28 (s, 3 H), 4.91-5.20 (m, 3H), 5.48 (s, 1 H), 5.7-5.9 (m, 1 H), 7.35-7.76 (m, 8 H).

10

Step 3: Preparation of Example 47;




Ammonia gas was bubbled through as a solution of 47" (0.38g, 0.88 mmol), DMF (15 mL), HATU (0.4 g, 1.1 mmol), and DIPEA (4.3 mmol, 0.76 mL) at 0°C for 10 min. The solution was allowed to warm to room temperature while stirring for 18 h. The DMF was removed under reduced pressure and the resulting residue was dissolved in ethyl acetate (200 mL), washed with 0.1 N HCl (2 x 100 mL), brine (2 x 100 mL), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, and concentrated to yield a light brown oil. The crude material was further purified via silica gel chromatography (2% MeOH, 98% CH<sub>2</sub>Cl<sub>2</sub>) to yield the title compound, Example 47, as an off-white solid (0.21 g, 55%): mp 137-145 °C.

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.82-1.08 (m, 6 H), 1.22-1.79 (m, 3 H) 2.13-2.82 (m, 4 H) 3.35 (s, 3 H) 4.94-5.12 (m, 3 H), 5.36 (d, 1 H), 5.52 (d, 1 H), 5.67-5.81 (m, 1 H), 7.30-7.63 (m, 8 H); IR (KBr) 3334, 2955, 1654, 1498, 1386 cm<sup>-1</sup>; CI MS m/z = 434 [C<sub>26</sub>H<sub>31</sub>N<sub>3</sub>O<sub>3</sub> + H]<sup>+</sup>; HPLC 95.4% tr = 17.65 min.

Anal. Calcd. for  $C_{26}H_{31}N_3O_3 \cdot 0.25 H_2O$ : C, 72.29; H, 7.25; N, 9.59, Found: C, 71.22; H, 7.13; N, 9.32.


**Example 48**

- 5 (2R,3S) N1-[1,3,4,5-tetrahydro-1,5-dimethyl-2,4-dioxo-2H-1,5-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide



10

Step 1: Preparation of Compound 48';



15

To a solution of tert-butyl succinate ester (9) ( 1.0 eq) in DMF ( 0.25 M) at zero degrees was added HATU ( 1.1 eq), then Hunig's base (4.0 eq). The mixture was stirred at zero degrees for 10 mins. A solution of 3 amino-  
 20 1,3,4,5-tetrahydro-1,5-dimethyl-2H-1,5-benzodiazepin-2,4-dione (32) in DMF ( 0.8 M) (1.0 eq) was added to this solution. The reaction mixture was stirred overnight at RT and then transferred to a separatory funnel containing water. 30% N-hexane in ethyl acetate was added which gave  
 25 a clear organic layer. The aqueous solution was extracted twice with 30% n-hexane in ethyl acetate. The combined organic layers were washed with water and brine, dried over magnesium sulfate, and concentrated in vacuo. The residue


was purified by chromatography on flash grade silica gel using 30% ethyl acetate in n-hexane. Compound **48'** was isolated as an amorphous white solid (92%).  $R_f = 0.15$  (7 : 3 n-hexane:ethyl acetate).

5

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  7.34 (s, 4H); 6.97-6.94 (d, 1H); 5.80-5.60 (m, 1H); 5.15 - 4.95 (m, 3H); 3.45 (s, 6H); 2.65-2.20 (m, 4H); 1.80-1.50 (m, 2H); 1.18-1.00 (m, 1H); 0.95-0.92 (d, 3H); 0.87-0.84 (d, 3H); MS (M+H)<sup>+</sup> 472.2.


10

Step 2: Preparation of Compound **48''**;



Following a procedure analogous to the procedure of Step 2 in Example 1, compound **48''** was prepared.

Step 3: Preparation of Example 48;




20

Following a procedure analogous to the procedure of Step 3 in Example 1, the title compound, Example 48, was prepared.

- $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  7.36 (s, 4H); 7.10-7.00 (d, 1H); 6.44 (s, 1H); 5.85-5.75 (m, 1H); 5.40 (m, 1H); 5.19-5.00 (m, 3H); 3.50-3.45 (d, 6H); 2.70-2.33 (m, 4H); 1.60-1.40 (m, 2H); 1.30-1.20 (m, 1H); 0.90-0.85 (q, 6H).
- 5 MS  $(\text{M}+\text{H})^+ = 415.4$ ,  $(\text{M}+\text{Na})^+ = 437.4$ .

**Example 49**

- (2R,3S) N1-[1,3,4,5-tetrahydro-1-methyl-2-oxo-2H-benzoazepin-3-yl]-2-(2-methylpropyl)-3-allyl-butanediamide
- 10




- Following a procedure analogous to the preparation of
- 15 Example 1 using 3-amino-1,3,4,5-tetrahydro-1-methyl-2H-benzoazepine-2-one, prepared by methods known to one skilled in the art, in place of (30) in Step 1, the title compound was prepared. MS  $(\text{M}+\text{H})^+ = 408.3$ .

20

**Example 50**

- (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]-2-methyl-3-propyl-butanediamide



- 25 Using the product of Example 9a as starting material, and following the reduction procedure of Example 7, the title compound was prepared. MS  $(\text{M}+\text{Na})^+ = 443.2$ .







Table 2 demonstrates representative compounds envisaged within the scope of the present invention. Each formulae at the start of Table 2 are intended to be paired with each entry in the table which follows.

5 For example the compound (2R,3S) N1-[6,7-dihydro-5-methyl-6-oxo-5H-dibenz[b,d]azepin-7-yl]-2-(2-methylpropyl)-3-(allyl)-butanediamide is represented by Example #500-B-j, which comprises the core B, succinate j, and entry #500.



10 For example the compound (2R,3S) N1-[(3S)-1,3-dihydro-1-methyl-2-oxo-5-phenyl-7-chloro-2H-1,4-benzodiazepin-3-yl]-2-(2-methylpropyl)-3-(butyl)-butanediamide, is represented by Example #502-D-ab, which comprises the core D, succinate ab, and entry #502.

15











Table 2



20




25




P

Q












R




S


5

wherein R<sup>3</sup> and R<sup>5</sup> are:abcdefghi


15




i




k




l




m




n




o




p




q




r




s




t




u









v









w






x













**Y****Z****aa****ab****ac****ad**


5

**ae****af****ag****ah****ai****aj**


10

**ak****al****am**


anaoapaqarasatauavawaxay




az




ba




bb



bc




bd




be


5




bf




bg




bh



bi












bj



bk

10

**b1****bm****bn****bo****bp****bq****br****bs****bt**

5

10

**Table 2( cont.)**

Ex#	core	R3/R5	R13	Z
500	A - S	a - bt	H	methyl
501	A - S	a - bt	F	methyl
502	A - S	a - bt	Cl	methyl
503	A - S	a - bt	OH	methyl
504	A - S	a - bt	-CH <sub>3</sub>	methyl
505	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	methyl
506	A - S	a - bt	-OCH <sub>3</sub>	methyl
507	A - S	a - bt	-CF <sub>3</sub>	methyl
508	A - S	a - bt	H	ethyl
509	A - S	a - bt	F	ethyl
510	A - S	a - bt	Cl	ethyl
511	A - S	a - bt	OH	ethyl



512	A - S	a - bt	-CH <sub>3</sub>	ethyl
513	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	ethyl
514	A - S	a - bt	-OCH <sub>3</sub>	ethyl
515	A - S	a - bt	-CF <sub>3</sub>	ethyl
516	A - S	a - bt	H	i-propyl
517	A - S	a - bt	F	i-propyl
518	A - S	a - bt	Cl	i-propyl
519	A - S	a - bt	OH	i-propyl
520	A - S	a - bt	-CH <sub>3</sub>	i-propyl
521	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	i-propyl
522	A - S	a - bt	-OCH <sub>3</sub>	i-propyl
523	A - S	a - bt	-CF <sub>3</sub>	i-propyl
524	A - S	a - bt	H	n-propyl
525	A - S	a - bt	F	n-propyl
526	A - S	a - bt	Cl	n-propyl
527	A - S	a - bt	OH	n-propyl
528	A - S	a - bt	-CH <sub>3</sub>	n-propyl
529	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	n-propyl
530	A - S	a - bt	-OCH <sub>3</sub>	n-propyl
531	A - S	a - bt	-CF <sub>3</sub>	n-propyl
532	A - S	a - bt	H	n-butyl
533	A - S	a - bt	F	n-butyl
534	A - S	a - bt	Cl	n-butyl
535	A - S	a - bt	OH	n-butyl
536	A - S	a - bt	-CH <sub>3</sub>	n-butyl
537	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	n-butyl
538	A - S	a - bt	-OCH <sub>3</sub>	n-butyl
539	A - S	a - bt	-CF <sub>3</sub>	n-butyl
540	A - S	a - bt	H	i-butyl
541	A - S	a - bt	F	i-butyl
542	A - S	a - bt	Cl	i-butyl
543	A - S	a - bt	OH	i-butyl
544	A - S	a - bt	-CH <sub>3</sub>	i-butyl
545	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	i-butyl
546	A - S	a - bt	-OCH <sub>3</sub>	i-butyl
547	A - S	a - bt	-CF <sub>3</sub>	i-butyl
548	A - S	a - bt	H	s-butyl
549	A - S	a - bt	F	s-butyl
550	A - S	a - bt	Cl	s-butyl
551	A - S	a - bt	OH	s-butyl
552	A - S	a - bt	-CH <sub>3</sub>	s-butyl
553	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	s-butyl
554	A - S	a - bt	-OCH <sub>3</sub>	s-butyl
555	A - S	a - bt	-CF <sub>3</sub>	s-butyl
556	A - S	a - bt	H	t-butyl
557	A - S	a - bt	F	t-butyl
558	A - S	a - bt	Cl	t-butyl
559	A - S	a - bt	OH	t-butyl
560	A - S	a - bt	-CH <sub>3</sub>	t-butyl
561	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	t-butyl
562	A - S	a - bt	-OCH <sub>3</sub>	t-butyl
563	A - S	a - bt	-CF <sub>3</sub>	t-butyl
564	A - S	a - bt	H	allyl
565	A - S	a - bt	F	allyl

566	A - S	a - bt	Cl	allyl
567	A - S	a - bt	OH	allyl
568	A - S	a - bt	-CH <sub>3</sub>	allyl
569	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	allyl
570	A - S	a - bt	-OCH <sub>3</sub>	allyl
571	A - S	a - bt	-CF <sub>3</sub>	allyl
572	A - S	a - bt	H	cyclopropyl
573	A - S	a - bt	F	cyclopropyl
574	A - S	a - bt	Cl	cyclopropyl
575	A - S	a - bt	OH	cyclopropyl
576	A - S	a - bt	-CH <sub>3</sub>	cyclopropyl
577	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	cyclopropyl
578	A - S	a - bt	-OCH <sub>3</sub>	cyclopropyl
579	A - S	a - bt	-CF <sub>3</sub>	cyclopropyl
580	A - S	a - bt	-CF <sub>3</sub>	cyclopropyl
581	A - S	a - bt	H	cyclopropyl-CH <sub>2</sub> -
582	A - S	a - bt	F	cyclopropyl-CH <sub>2</sub> -
583	A - S	a - bt	Cl	cyclopropyl-CH <sub>2</sub> -
584	A - S	a - bt	OH	cyclopropyl-CH <sub>2</sub> -
585	A - S	a - bt	-CH <sub>3</sub>	cyclopropyl-CH <sub>2</sub> -
586	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	cyclopropyl-CH <sub>2</sub> -
587	A - S	a - bt	-OCH <sub>3</sub>	cyclopropyl-CH <sub>2</sub> -
588	A - S	a - bt	-CF <sub>3</sub>	cyclopropyl-CH <sub>2</sub> -
589	A - S	a - bt	H	cyclobutyl
590	A - S	a - bt	F	cyclobutyl
591	A - S	a - bt	Cl	cyclobutyl
592	A - S	a - bt	OH	cyclobutyl
593	A - S	a - bt	-CH <sub>3</sub>	cyclobutyl
594	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	cyclobutyl
595	A - S	a - bt	-OCH <sub>3</sub>	cyclobutyl
596	A - S	a - bt	-CF <sub>3</sub>	cyclobutyl
597	A - S	a - bt	H	cyclobutyl-CH <sub>2</sub> -
598	A - S	a - bt	F	cyclobutyl-CH <sub>2</sub> -
599	A - S	a - bt	Cl	cyclobutyl-CH <sub>2</sub> -
600	A - S	a - bt	OH	cyclobutyl-CH <sub>2</sub> -
601	A - S	a - bt	-CH <sub>3</sub>	cyclobutyl-CH <sub>2</sub> -
602	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	cyclobutyl-CH <sub>2</sub> -
603	A - S	a - bt	-OCH <sub>3</sub>	cyclobutyl-CH <sub>2</sub> -
604	A - S	a - bt	-CF <sub>3</sub>	cyclobutyl-CH <sub>2</sub> -
605	A - S	a - bt	H	cyclopentyl
606	A - S	a - bt	F	cyclopentyl
607	A - S	a - bt	Cl	cyclopentyl
608	A - S	a - bt	OH	cyclopentyl
609	A - S	a - bt	-CH <sub>3</sub>	cyclopentyl
610	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	cyclopentyl
611	A - S	a - bt	-OCH <sub>3</sub>	cyclopentyl
612	A - S	a - bt	-CF <sub>3</sub>	cyclopentyl
613	A - S	a - bt	H	cyclopentyl-CH <sub>2</sub> -
614	A - S	a - bt	F	cyclopentyl-CH <sub>2</sub> -
615	A - S	a - bt	Cl	cyclopentyl-CH <sub>2</sub> -
616	A - S	a - bt	OH	cyclopentyl-CH <sub>2</sub> -
617	A - S	a - bt	-CH <sub>3</sub>	cyclopentyl-CH <sub>2</sub> -

618	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	cyclopentyl-CH <sub>2</sub> -
619	A - S	a - bt	-OCH <sub>3</sub>	cyclopentyl-CH <sub>2</sub> -
620	A - S	a - bt	-CF <sub>3</sub>	cyclopentyl-CH <sub>2</sub> -
621	A - S	a - bt	H	cyclohexyl
622	A - S	a - bt	F	cyclohexyl
623	A - S	a - bt	Cl	cyclohexyl
624	A - S	a - bt	OH	cyclohexyl
625	A - S	a - bt	-CH <sub>3</sub>	cyclohexyl
626	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	cyclohexyl
627	A - S	a - bt	-OCH <sub>3</sub>	cyclohexyl
628	A - S	a - bt	-CF <sub>3</sub>	cyclohexyl
629	A - S	a - bt	H	cyclohexyl-CH <sub>2</sub> -
630	A - S	a - bt	F	cyclohexyl-CH <sub>2</sub> -
631	A - S	a - bt	Cl	cyclohexyl-CH <sub>2</sub> -
632	A - S	a - bt	OH	cyclohexyl-CH <sub>2</sub> -
633	A - S	a - bt	-CH <sub>3</sub>	cyclohexyl-CH <sub>2</sub> -
634	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	cyclohexyl-CH <sub>2</sub> -
635	A - S	a - bt	-OCH <sub>3</sub>	cyclohexyl-CH <sub>2</sub> -
636	A - S	a - bt	-CF <sub>3</sub>	cyclohexyl-CH <sub>2</sub> -
637	A - S	a - bt	H	phenyl
638	A - S	a - bt	F	phenyl
639	A - S	a - bt	Cl	phenyl
640	A - S	a - bt	OH	phenyl
641	A - S	a - bt	-CH <sub>3</sub>	phenyl
642	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	phenyl
643	A - S	a - bt	-OCH <sub>3</sub>	phenyl
644	A - S	a - bt	-CF <sub>3</sub>	phenyl
645	A - S	a - bt	H	2-F-phenyl
646	A - S	a - bt	F	2-F-phenyl
647	A - S	a - bt	Cl	2-F-phenyl
648	A - S	a - bt	OH	2-F-phenyl
649	A - S	a - bt	-CH <sub>3</sub>	2-F-phenyl
650	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl
651	A - S	a - bt	-OCH <sub>3</sub>	2-F-phenyl
652	A - S	a - bt	-CF <sub>3</sub>	2-F-phenyl
653	A - S	a - bt	H	3-F-phenyl
654	A - S	a - bt	F	3-F-phenyl
655	A - S	a - bt	Cl	3-F-phenyl
656	A - S	a - bt	OH	3-F-phenyl
657	A - S	a - bt	-CH <sub>3</sub>	3-F-phenyl
658	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	3-F-phenyl
659	A - S	a - bt	-OCH <sub>3</sub>	3-F-phenyl
660	A - S	a - bt	-CF <sub>3</sub>	3-F-phenyl
661	A - S	a - bt	H	4-F-phenyl
662	A - S	a - bt	F	4-F-phenyl
663	A - S	a - bt	Cl	4-F-phenyl
664	A - S	a - bt	OH	4-F-phenyl
665	A - S	a - bt	-CH <sub>3</sub>	4-F-phenyl
666	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	4-F-phenyl
667	A - S	a - bt	-OCH <sub>3</sub>	4-F-phenyl
668	A - S	a - bt	-CF <sub>3</sub>	4-F-phenyl
669	A - S	a - bt	H	3-Cl-phenyl
670	A - S	a - bt	F	3-Cl-phenyl
671	A - S	a - bt	Cl	3-Cl-phenyl

672	A - S	a - bt	OH	3-Cl-phenyl
673	A - S	a - bt	-CH <sub>3</sub>	3-Cl-phenyl
674	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	3-Cl-phenyl
675	A - S	a - bt	-OCH <sub>3</sub>	3-Cl-phenyl
676	A - S	a - bt	-CF <sub>3</sub>	3-Cl-phenyl
677	A - S	a - bt	H	4-Cl-phenyl
678	A - S	a - bt	F	4-Cl-phenyl
679	A - S	a - bt	Cl	4-Cl-phenyl
680	A - S	a - bt	OH	4-Cl-phenyl
681	A - S	a - bt	-CH <sub>3</sub>	4-Cl-phenyl
682	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	4-Cl-phenyl
683	A - S	a - bt	-OCH <sub>3</sub>	4-Cl-phenyl
684	A - S	a - bt	-CF <sub>3</sub>	4-Cl-phenyl
685	A - S	a - bt	H	3-Me-phenyl
686	A - S	a - bt	F	3-Me-phenyl
687	A - S	a - bt	Cl	3-Me-phenyl
688	A - S	a - bt	OH	3-Me-phenyl
689	A - S	a - bt	-CH <sub>3</sub>	3-Me-phenyl
690	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	3-Me-phenyl
691	A - S	a - bt	-OCH <sub>3</sub>	3-Me-phenyl
692	A - S	a - bt	-CF <sub>3</sub>	3-Me-phenyl
693	A - S	a - bt	H	4-Me-phenyl
694	A - S	a - bt	F	4-Me-phenyl
695	A - S	a - bt	Cl	4-Me-phenyl
696	A - S	a - bt	OH	4-Me-phenyl
697	A - S	a - bt	-CH <sub>3</sub>	4-Me-phenyl
698	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	4-Me-phenyl
699	A - S	a - bt	-OCH <sub>3</sub>	4-Me-phenyl
700	A - S	a - bt	-CF <sub>3</sub>	4-Me-phenyl
701	A - S	a - bt	H	3-MeO-phenyl
702	A - S	a - bt	F	3-MeO-phenyl
703	A - S	a - bt	Cl	3-MeO-phenyl
704	A - S	a - bt	OH	3-MeO-phenyl
705	A - S	a - bt	-CH <sub>3</sub>	3-MeO-phenyl
706	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	3-MeO-phenyl
707	A - S	a - bt	-OCH <sub>3</sub>	3-MeO-phenyl
708	A - S	a - bt	-CF <sub>3</sub>	3-MeO-phenyl
709	A - S	a - bt	H	4-MeO-phenyl
710	A - S	a - bt	F	4-MeO-phenyl
711	A - S	a - bt	Cl	4-MeO-phenyl
712	A - S	a - bt	OH	4-MeO-phenyl
713	A - S	a - bt	-CH <sub>3</sub>	4-MeO-phenyl
714	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	4-MeO-phenyl
715	A - S	a - bt	-OCH <sub>3</sub>	4-MeO-phenyl
716	A - S	a - bt	-CF <sub>3</sub>	4-MeO-phenyl
717	A - S	a - bt	H	3-F <sub>3</sub> C-phenyl
718	A - S	a - bt	F	3-F <sub>3</sub> C-phenyl
719	A - S	a - bt	Cl	3-F <sub>3</sub> C-phenyl
720	A - S	a - bt	OH	3-F <sub>3</sub> C-phenyl
721	A - S	a - bt	-CH <sub>3</sub>	3-F <sub>3</sub> C-phenyl
722	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	3-F <sub>3</sub> C-phenyl
723	A - S	a - bt	-OCH <sub>3</sub>	3-F <sub>3</sub> C-phenyl
724	A - S	a - bt	-CF <sub>3</sub>	3-F <sub>3</sub> C-phenyl

725	A - S	a - bt	H	4-F <sub>3</sub> C-phenyl
726	A - S	a - bt	F	4-F <sub>3</sub> C-phenyl
727	A - S	a - bt	Cl	4-F <sub>3</sub> C-phenyl
728	A - S	a - bt	OH	4-F <sub>3</sub> C-phenyl
729	A - S	a - bt	-CH <sub>3</sub>	4-F <sub>3</sub> C-phenyl
730	A - S	a - bt	-CH <sub>2</sub> CH <sub>3</sub>	4-F <sub>3</sub> C-phenyl
731	A - S	a - bt	-OCH <sub>3</sub>	4-F <sub>3</sub> C-phenyl
732	A - S	a - bt	-CF <sub>3</sub>	4-F <sub>3</sub> C-phenyl

UTILITY

A $\beta$  production has been implicated in the pathology of Alzheimer's Disease (AD). The compounds of the present invention have utility for the prevention and treatment of AD by inhibiting A $\beta$  production. Methods of treatment target formation of A $\beta$  production through the enzymes involved in the proteolytic processing of  $\beta$  amyloid precursor protein. Compounds that inhibit  $\beta$  or  $\gamma$ secretase activity, either directly or indirectly, control the production of A $\beta$ . Such inhibition of  $\beta$  or  $\gamma$ secretases reduces production of A $\beta$ , and is expected to reduce or prevent the neurological disorders associated with A $\beta$  protein, such as Alzheimer's Disease.

Cellular screening methods for inhibitors of A $\beta$  production, testing methods for the *in vivo* suppression of A $\beta$  production, and assays for the detection of secretase activity are known in the art and have been disclosed in numerous publications, including PCT publication number WO 98/22493, EPO publication number 0652009, US patent 5703129 and US patent 5593846; all hereby incorporated by reference.

The compounds of the present invention have utility for the prevention and treatment of disorders involving A $\beta$  production, such as cerebrovascular disorders.

Compounds of the present invention have been shown to inhibit A $\beta$  production, as determined by the secretase inhibition assay described below.

Compounds of the present invention have been shown to inhibit A $\beta$  production, utilizing the C-terminus  $\beta$  amyloid precursor protein accumulation assay described below.

Compounds of Formula (I) are expected to possess  $\gamma$ -secretase inhibitory activity. The  $\gamma$ -secretase inhibitory activity of the compounds of the present invention is demonstrated using assays for such activity, for example, using the assay described below. Compounds of the present invention have been shown to inhibit the activity of  $\gamma$ -

secretase, as determined by the A $\beta$  immunoprecipitation assay.

Compounds provided by this invention should also be useful as standards and reagents in determining the ability of a potential pharmaceutical to inhibit A $\beta$  production. These would be provided in commercial kits comprising a compound of this invention.

As used herein "ug" denotes microgram, "mg" denotes milligram, "g" denotes gram, "uL" denotes microliter, "mL" denotes milliliter, "L" denotes liter, "nM" denotes nanomolar, "uM" denotes micromolar, "mM" denotes millimolar, "M" denotes molar, "nm" denotes nanometer, "SDS" denotes sodium dodecyl sulfate, and "DMSO" denotes dimethyl sulfoxide, and "EDTA" denotes ethylenediaminetetraacetate.

A compound is considered to be active if it has an IC<sub>50</sub> or K<sub>i</sub> value of less than about 100 uM for the inhibition of A $\beta$  production.

#### 20 $\beta$ amyloid precursor protein accumulation assay

A novel assay to evaluate the accumulation of A $\beta$  protein was developed to detect potential inhibitors of secretase. The assay uses the N 9 cell line, characterized for expression of exogenous APP by immunoblotting and immunoprecipitation.

The effect of test compounds on the accumulation of A $\beta$  in the conditioned medium is tested by immunoprecipitation. Briefly, N 9 cells are grown to confluency in 6-well plates and washed twice with 1 x Hank's buffered salt solution. The cells are starved in methionine/cysteine deficient media for 30 min, followed by replacement with fresh deficient media containing 150uCi S35 Translabel (Amersham). Test compounds dissolved in DMSO (final concentration 1%) are added together with the addition of radiolabel. The cells are incubated for 4 h at 37°C in a tissue culture incubator.

At the end of the incubation period, the conditioned medium is harvested and pre-cleared by the addition of 5 uL

normal mouse serum and 50ul of protein A Sepharose (Pharmacia), mixed by end-over-end rotation for 30 minutes at 4°C, followed by a brief centrifugation in a microfuge. The supernatant is then harvested and transferred to fresh  
5 tubes containing 5ug of a monoclonal antibody (clone 1101.1; directed against an internal peptide sequence in A $\beta$ ) and 50  $\mu$ l protein A Sepharose. After incubation overnight at 4°C, the samples are washed three times with high salt washing buffer (50mM Tris, pH 7.5, 500mM NaCl,  
10 5mM EDTA, 0.5% Nonidet P-40), three times with low salt wash buffer (50mM Tris, pH 7.5, 150mM NaCl, 5mM EDTA, 0.5% Nonidet P-40), and three times with 10mM Tris, pH 7.5. The pellet after the last wash is resuspended in SDS sample buffer (Laemmli, 1970) and boiled for 3 minutes. The  
15 supernatant is then fractionated on either 10-20% Tris/Tricine SDS gels or on 16.5% Tris/Tricine SDS gels. The gels are dried and exposed to X-ray film or analyzed by phosphorimaging. The resulting image is analyzed for the presence of A $\beta$  polypeptides. The steady-state level of A $\beta$   
20 in the presence of a test compound is compared to wells treated with DMSO (1%) alone. A typical test compound blocks A $\beta$  accumulation in the conditioned medium, and is therefore considered active, with an IC<sub>50</sub> less than 100  $\mu$ M.

25 C-Terminus  $\beta$  Amyloid Precursor Protein Accumulation Assay

The effect of test compounds on the accumulation of C-terminal fragments is determined by immunoprecipitation of APP and fragments thereof from cell lysates. N 9 cells are metabolically labeled as above in the presence or absence  
30 of test compounds. At the end of the incubation period, the conditioned medium are harvested and cells lysed in RIPA buffer (10 mM Tris, pH 8.0 containing 1% Triton X-100, 1% deoxycholate, 0.1% SDS, 150mM NaCl, 0.125% NaN<sub>3</sub>). Again, lysates are precleared with 5ul normal rabbit serum  
35 / 50ul protein A Sepharose, followed by the addition of BC-1 antiserum (15ul;) and 50ul protein A Sepharose for 16 hours at 4°C. The immunoprecipitates are washed as above, bound proteins eluted by boiling in SDS sample buffer and



fractionated by Tris/Tricine SDS-PAGE. After exposure to X-ray film or phosphorimager, the resulting images are analyzed for the presence of C-terminal APP fragments. The steady-state level of C-terminal APP fragments is compared to wells treated with DMSO (1%) alone. A typical test compound stimulates C-terminal fragment accumulation in the cell lysates, and is therefore considered active, with an  $IC_{50}$  less than 100  $\mu M$ .

#### 10 A $\beta$ Immunoprecipitation Assay

This immunoprecipitation assay is specific for  $\gamma$  secretase (i.e., proteolytic activity required to generate the C-terminal end of A $\beta$  either by direct cleavage or generating a C-terminal extended species which is subsequently further proteolyzed). N 9 cells are pulse labeled in the presence of a reported  $\gamma$  secretase inhibitor (MDL 28170) for 1 h, followed by washing to remove radiolabel and MDL 28170. The media is replaced and test compounds are added. The cells are chased for increasing periods of times and A  $\beta$  is isolated from the conditioned medium and C-terminal fragments from cell lysates (see above). The test compounds are characterized whether a stabilization of C-terminal fragments is observed and whether A $\beta$  is generated from these accumulated precursor. A typical test compound prevents the generation of A $\beta$  out of accumulated C-terminal fragments and is considered active with an  $IC_{50}$  less than 100  $\mu M$ .

#### Dosage and Formulation

30 The compounds of the present invention can be administered orally using any pharmaceutically acceptable dosage form known in the art for such administration. The active ingredient can be supplied in solid dosage forms such as dry powders, granules, tablets or capsules, or in liquid dosage forms, such as syrups or aqueous suspensions. The active ingredient can be administered alone, but is generally administered with a pharmaceutical carrier. A valuable treatise with respect to pharmaceutical dosage

forms is Remington's Pharmaceutical Sciences, Mack Publishing.

The compounds of the present invention can be administered in such oral dosage forms as tablets, capsules  
5 (each of which includes sustained release or timed release formulations), pills, powders, granules, elixirs, tinctures, suspensions, syrups, and emulsions. Likewise, they may also be administered in intravenous (bolus or infusion), intraperitoneal, subcutaneous, or intramuscular  
10 form, all using dosage forms well known to those of ordinary skill in the pharmaceutical arts. An effective but non-toxic amount of the compound desired can be employed to prevent or treat neurological disorders related to  $\beta$ -amyloid production or accumulation, such as  
15 Alzheimer's disease and Down's Syndrome.

The compounds of this invention can be administered by any means that produces contact of the active agent with the agent's site of action in the body of a host, such as a human or a mammal. They can be administered by any  
20 conventional means available for use in conjunction with pharmaceuticals, either as individual therapeutic agents or in a combination of therapeutic agents. They can be administered alone, but generally administered with a pharmaceutical carrier selected on the basis of the chosen  
25 route of administration and standard pharmaceutical practice.

The dosage regimen for the compounds of the present invention will, of course, vary depending upon known factors, such as the pharmacodynamic characteristics of the  
30 particular agent and its mode and route of administration; the species, age, sex, health, medical condition, and weight of the recipient; the nature and extent of the symptoms; the kind of concurrent treatment; the frequency of treatment; the route of administration, the renal and  
35 hepatic function of the patient, and the effect desired. An ordinarily skilled physician or veterinarian can readily determine and prescribe the effective amount of the drug

required to prevent, counter, or arrest the progress of the condition.

Advantageously, compounds of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three, or four times daily.

The compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

In the methods of the present invention, the compounds herein described in detail can form the active ingredient, and are typically administered in admixture with suitable pharmaceutical diluents, excipients, or carriers (collectively referred to herein as carrier materials) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic, pharmaceutically acceptable, inert carrier such as lactose, starch, sucrose, glucose, methyl cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, mannitol, sorbitol and the like; for oral administration in liquid form, the oral drug components can be combined with any oral, non-toxic, pharmaceutically acceptable inert carrier such as ethanol, glycerol, water, and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents, and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or  $\beta$ -lactose, corn sweeteners, natural and synthetic gums such as acacia,

tragacanth, or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes, and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride, and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum, and the like.

The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine, or phosphatidylcholines.

Compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxyethylaspartamidephenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacylates, and crosslinked or amphipathic block copolymers of hydrogels.

Gelatin capsules may contain the active ingredient and powdered carriers, such as lactose, starch, cellulose derivatives, magnesium stearate, stearic acid, and the like. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous release of medication over a period of hours. Compressed tablets can be sugar coated or film coated to mask any unpleasant taste and protect the tablet from the atmosphere, or enteric coated for selective disintegration in the gastrointestinal tract.

Liquid dosage forms for oral administration can contain coloring and flavoring to increase patient acceptance. In general, water, a suitable oil, saline, aqueous dextrose (glucose), and related sugar solutions and glycols such as

5 propylene glycol or polyethylene glycols are suitable carriers for parenteral solutions. Solutions for parenteral administration preferably contain a water soluble salt of the active ingredient, suitable stabilizing agents, and if necessary, buffer substances. Antioxidizing

10 agents such as sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or combined, are suitable stabilizing agents. Also used are citric acid and its salts and sodium EDTA. In addition, parenteral solutions can contain preservatives, such as benzalkonium chloride,


15 methyl- or propyl-paraben, and chlorobutanol.

Suitable pharmaceutical carriers are described in Remington's Pharmaceutical Sciences, Mack Publishing Company, a standard reference text in this field.

CLAIMS

What is claimed is:

1. A compound of Formula (I):



- or a pharmaceutically acceptable salt or prodrug thereof, wherein:

Q is  $-NR^1R^2$ ;

- $R^1$ , at each occurrence, is independently selected from:
- H;
  - $C_1$ - $C_6$  alkyl substituted with 0-3  $R^{1a}$ ;
  - $C_3$ - $C_{10}$  carbocycle substituted with 0-3  $R^{1b}$ ;
  - $C_6$ - $C_{10}$  aryl substituted with 0-3  $R^{1b}$ ; and
  - 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3  $R^{1b}$ ;
- $R^{1a}$ , at each occurrence, is independently selected from H,  $C_1$ - $C_6$  alkyl Cl, F, Br, I, =O, CN,  $NO_2$ ,  $NR^{15}R^{16}$ ,  $CF_3$ ;
- $C_3$ - $C_{10}$  carbocycle substituted with 0-3  $R^{1b}$ ;
  - $C_6$ - $C_{10}$  aryl substituted with 0-3  $R^{1b}$ ; and
  - 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3  $R^{1b}$ ;

R<sup>1b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>6</sub> haloalkyl, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy;

5 R<sup>2</sup> is independently selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>10</sub> carbocycle, C<sub>6</sub>-C<sub>10</sub> aryl, and 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur;

10 R<sup>3</sup> is -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-R<sup>4</sup>,  
 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-S-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>,  
 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-O-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>,  
 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-N(R<sup>7b</sup>)-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>,  
 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-S(=O)-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>,  
 15 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-S(=O)<sub>2</sub>-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>,  
 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-C(=O)-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>,  
 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-N(R<sup>7b</sup>)C(=O)-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>,  
 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-C(=O)N(R<sup>7b</sup>)-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>,  
 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-N(R<sup>7b</sup>)S(=O)<sub>2</sub>-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>, or  
 20 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-S(=O)<sub>2</sub>N(R<sup>7b</sup>)-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>;

n is 0, 1, 2, or 3;

m is 0, 1, 2, or 3;

25

R<sup>3a</sup> is H, OH, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>2</sub>-C<sub>4</sub> alkenyl or C<sub>2</sub>-C<sub>4</sub> alkenyloxy;

R<sup>4</sup> is H, OH, OR<sup>14a</sup>,

30

C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 0-3 R<sup>4a</sup>,  
 C<sub>2</sub>-C<sub>6</sub> alkenyl substituted with 0-3 R<sup>4a</sup>,  
 C<sub>2</sub>-C<sub>6</sub> alkynyl substituted with 0-3 R<sup>4a</sup>,  
 C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,  
 C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>4b</sup>, or

35

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R<sup>4b</sup>;

- R<sup>4a</sup>, at each occurrence, is independently selected from is  
H, F, Cl, Br, I, CF<sub>3</sub>,  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,  
5 C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>4b</sup>, or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>4b</sup>;  
10
- R<sup>4b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>,  
S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>,  
C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl,  
15 C<sub>1</sub>-C<sub>4</sub> haloalkoxy, and C<sub>1</sub>-C<sub>4</sub> halothioalkyl-S-;
- R<sup>5</sup> is H, OR<sup>14</sup>;  
C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 0-3 R<sup>5b</sup>;  
C<sub>1</sub>-C<sub>6</sub> alkoxy substituted with 0-3 R<sup>5b</sup>;  
20 C<sub>2</sub>-C<sub>6</sub> alkenyl substituted with 0-3 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>6</sub> alkynyl substituted with 0-3 R<sup>5b</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>5c</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>5c</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
25 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>5c</sup>;
- R<sup>5a</sup> is H, OH, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>2</sub>-C<sub>4</sub> alkenyl, or  
30 C<sub>2</sub>-C<sub>4</sub> alkenyloxy;
- R<sup>5b</sup>, at each occurrence, is independently selected from:  
H, C<sub>1</sub>-C<sub>6</sub> alkyl, CF<sub>3</sub>, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>,  
NR<sup>15</sup>R<sup>16</sup>;  
35 C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>5c</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>5c</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and



sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R<sup>5c</sup>;

5 R<sup>5c</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl, C<sub>1</sub>-C<sub>4</sub> haloalkoxy, and C<sub>1</sub>-C<sub>4</sub> halothioalkyl-S-;

10 R<sup>6</sup> is H;

C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 0-3 R<sup>6a</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>6b</sup>; or  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>6b</sup>;

15 R<sup>6a</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, aryl or CF<sub>3</sub>;

20 R<sup>6b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy;

R<sup>7</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, CF<sub>3</sub>, phenyl and C<sub>1</sub>-C<sub>4</sub> alkyl;

25

R<sup>7a</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, CF<sub>3</sub>, and C<sub>1</sub>-C<sub>4</sub> alkyl;

R<sup>7b</sup> is independently selected from H and C<sub>1</sub>-C<sub>4</sub> alkyl;

30

Ring B is a 7 membered lactam,

wherein the lactam is saturated, partially saturated or unsaturated;

35 wherein each additional lactam carbon is substituted with 0-2 R<sup>11</sup>; and,

optionally, the lactam contains a heteroatom selected from -O-, -S-, -S(=O)-, -S(=O)<sub>2</sub>-, -N=, -NH-, and -N(R<sup>10</sup>)-;

additionally, two R<sup>11</sup> substituents on adjacent atoms may be combined to form a benzo fused radical; wherein said benzo fused radical is substituted with 0-4 R<sup>13</sup>;

5 additionally, two R<sup>11</sup> substituents on adjacent atoms may be combined to form a 5 to 6 membered heteroaryl fused radical, wherein said 5 to 6 membered heteroaryl fused radical comprises 1 or 2 heteroatoms selected from N,  
10 O, and S; wherein said 5 to 6 membered heteroaryl fused radical is substituted with 0-3 R<sup>13</sup>;

additionally, two R<sup>11</sup> substituents on the same or adjacent carbon atoms may be combined to form a C<sub>3</sub>-C<sub>6</sub>  
15 carbocycle substituted with 0-3 R<sup>13</sup>;

R<sup>10</sup> is H, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>,  
S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, S(=O)<sub>2</sub>R<sup>17</sup>;  
C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 0-3 R<sup>10a</sup>;  
20 C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>10b</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>10b</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
25 is substituted with 0-3 R<sup>10b</sup>;

R<sup>10a</sup>, at each occurrence, is independently selected from H,  
C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>,  
CF<sub>3</sub>, or aryl substituted with 0-4 R<sup>10b</sup>;

30 R<sup>10b</sup>, at each occurrence, is independently selected from H,  
OH, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, CN, NO<sub>2</sub>,  
NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub>  
alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl, C<sub>1</sub>-C<sub>4</sub> haloalkoxy,  
35 and C<sub>1</sub>-C<sub>4</sub> haloalkyl-S-;

R<sup>11</sup>, at each occurrence, is independently selected from

- H, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>18</sup>R<sup>19</sup>,  
C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, CF<sub>3</sub>;  
C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 0-3 R<sup>11a</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>11b</sup>;  
5 C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>11b</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>11b</sup>;  
10  
R<sup>11a</sup>, at each occurrence, is independently selected from  
H, C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>,  
NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>;  
phenyl substituted with 0-3 R<sup>11b</sup>;  
15 C<sub>3</sub>-C<sub>6</sub> cycloalkyl substituted with 0-3 R<sup>11b</sup>; and  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>11b</sup>;  
20  
R<sup>11b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>,  
S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>,  
C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl,  
25 C<sub>1</sub>-C<sub>4</sub> haloalkoxy, and C<sub>1</sub>-C<sub>4</sub> halothioalkyl-S-;  
  
Z is H;  
C<sub>1</sub>-C<sub>8</sub> alkyl substituted with 1-3 R<sup>12</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 1-3 R<sup>12</sup>;  
30 C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 1-3 R<sup>12</sup>;  
C<sub>1</sub>-C<sub>8</sub> alkyl substituted with 0-3 R<sup>12a</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>12a</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>12a</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>;  
35 C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-4 R<sup>12b</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R<sup>12b</sup>;

5 R<sup>12</sup>, at each occurrence, is independently selected from C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>12b</sup>; C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-4 R<sup>12b</sup>; or 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle  
10 is substituted with 0-3 R<sup>12b</sup>;

R<sup>12a</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, -C(=O)NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>,  
15 C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl, C<sub>1</sub>-C<sub>4</sub> haloalkoxy, or C<sub>1</sub>-C<sub>4</sub> halothioalkyl-S-;

R<sup>12b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>,  
20 C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl, C<sub>1</sub>-C<sub>4</sub> haloalkoxy, and C<sub>1</sub>-C<sub>4</sub> halothioalkyl-S-;

R<sup>13</sup>, at each occurrence, is independently selected from H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;  
25

R<sup>14</sup> is H, phenyl, benzyl, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkoxyalkyl, or C<sub>3</sub>-C<sub>6</sub> cycloalkyl;  
30

R<sup>14a</sup> is H, phenyl, benzyl, or C<sub>1</sub>-C<sub>4</sub> alkyl;

R<sup>15</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-;  
35

R<sup>16</sup>, at each occurrence, is independently selected from H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, phenethyl,

(C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-;

R<sup>17</sup> is H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkoxyalkyl,

5 aryl substituted by 0-4 R<sup>17a</sup>, or  
-CH<sub>2</sub>-aryl substituted by 0-4 R<sup>17a</sup>;

R<sup>17a</sup> is H, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
propoxy, butoxy, -OH, F, Cl, Br, I, CF<sub>3</sub>, OCF<sub>3</sub>, SCH<sub>3</sub>,  
S(O)CH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, -NH<sub>2</sub>, -N(CH<sub>3</sub>)<sub>2</sub>, or C<sub>1</sub>-C<sub>4</sub> haloalkyl;

10

R<sup>18</sup>, at each occurrence, is independently selected from  
H, C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl, benzyl, phenethyl,  
(C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-; and

15 R<sup>19</sup>, at each occurrence, is independently selected from  
H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl, benzyl, phenethyl,  
(C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-;

provided, when R<sup>13</sup> is H,

20 then Z is H;

C<sub>4</sub>-C<sub>8</sub> alkyl substituted with 1-3 R<sup>12</sup>;

C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 1-3 R<sup>12</sup>;

C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 1-3 R<sup>12</sup>;

C<sub>1</sub>-C<sub>8</sub> alkyl substituted with 0-3 R<sup>12a</sup>;

25 C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>12a</sup>; or

C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>12a</sup>; and

provided, when ring B is a 1,3,4,5-tetrahydro-1-(Z)-5-  
(R<sup>10</sup>)-6,6,7,7-tetra(R<sup>11</sup>)-2,4-dioxo-2H-1,5-diazepin-3-yl

30 core, and R<sup>13</sup> is H; then

R<sup>10</sup> is H, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>,


S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, S(=O)<sub>2</sub>R<sup>17</sup>; or

C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 0-3 R<sup>10a</sup>;

35

R<sup>10a</sup>, at each occurrence, is independently selected from  
H, C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>,  
NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>.

2. A compound, according to Claim 1, of Formula (Ia):



5

(Ia)

or a pharmaceutically acceptable salt or prodrug thereof,  
wherein:

10 Z is H;

C<sub>1</sub>-C<sub>8</sub> alkyl substituted with 0-3 R<sup>12a</sup>;

C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>12a</sup>; or

C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>12a</sup>.

15 3. A compound according to Claim 2 of Formula (Ia)  
wherein:

R<sup>3</sup> is -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-R<sup>4</sup>,

-(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-S-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>,

20 -(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-O-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>, or

-(CR<sup>7</sup>R<sup>7a</sup>)<sub>n</sub>-N(R<sup>7b</sup>)-(CR<sup>7</sup>R<sup>7a</sup>)<sub>m</sub>-R<sup>4</sup>;

n is 0, 1, or 2;

25 m is 0, 1, or 2;

R<sup>3a</sup> is H, OH, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, butoxy, allyl, or 3-buten-1-yl;

30 R<sup>4</sup> is H, OH, OR<sup>14a</sup>,

C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 0-3 R<sup>4a</sup>,

C<sub>2</sub>-C<sub>6</sub> alkenyl substituted with 0-3 R<sup>4a</sup>,

C<sub>2</sub>-C<sub>6</sub> alkynyl substituted with 0-3 R<sup>4a</sup>,

C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,

35 C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>4b</sup>, or

- 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R<sup>4b</sup>;
- 5 R<sup>4a</sup>, at each occurrence, is independently selected from is H, F, Cl, Br, I, CF<sub>3</sub>, C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>4b</sup>, C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>4b</sup>, or
- 10 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R<sup>4b</sup>;
- 15 R<sup>4b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy;
- 20 R<sup>5</sup> is H, OR<sup>14</sup>;  
C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 0-3 R<sup>5b</sup>;  
C<sub>1</sub>-C<sub>6</sub> alkoxy substituted with 0-3 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>6</sub> alkenyl substituted with 0-3 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>6</sub> alkynyl substituted with 0-3 R<sup>5b</sup>;
- 25 C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>5c</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>5c</sup>; or  
5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle
- 30 is substituted with 0-3 R<sup>5c</sup>;
- R<sup>5a</sup> is H or C<sub>1</sub>-C<sub>4</sub> alkyl;
- R<sup>5b</sup>, at each occurrence, is independently selected from:
- 35 H, C<sub>1</sub>-C<sub>6</sub> alkyl, CF<sub>3</sub>, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>5c</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>5c</sup>; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R<sup>5c</sup>;

5

R<sup>5c</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy;

10

R<sup>6</sup> is H, methyl, or ethyl;

R<sup>7</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, CF<sub>3</sub>, phenyl and C<sub>1</sub>-C<sub>4</sub> alkyl;

15

R<sup>7a</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, CF<sub>3</sub>, and C<sub>1</sub>-C<sub>4</sub> alkyl;

R<sup>7b</sup> is independently selected from H, methyl, ethyl, propyl, and butyl;

20

Ring B is a 7 membered lactam, wherein the lactam is saturated, partially saturated or unsaturated; wherein each additional lactam carbon is substituted with 0-2 R<sup>11</sup>; and, optionally, the lactam contains a heteroatom selected from, -O-, -S-, -S(=O)-, -S(=O)<sub>2</sub>-, -N=, -NH-, and -N(R<sup>10</sup>)-;

30

additionally, two R<sup>11</sup> substituents on adjacent atoms may be combined to form a benzo fused radical; wherein said benzo fused radical is substituted with 0-3 R<sup>13</sup>;

35 additionally, two R<sup>11</sup> substituents on adjacent atoms may be combined to form a 5 to 6 membered heteroaryl fused radical, wherein said 5 to 6 membered heteroaryl fused radical comprises 1 or 2 heteroatoms selected from N,



O, and S; wherein said 5 to 6 membered heteroaryl  
fused radical is substituted with 0-3 R<sup>13</sup>;

5 additionally, two R<sup>11</sup> substituents on the same or adjacent  
carbon atoms may be combined to form a C<sub>3</sub>-C<sub>6</sub>  
carbocycle substituted with 0-3 R<sup>13</sup>;

R<sup>10</sup> is H, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>,  
S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, S(=O)<sub>2</sub>R<sup>17</sup>;  
10 C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 0-2 R<sup>10a</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-4 R<sup>10b</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>10b</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
15 sulphur, wherein said 5 to 10 membered heterocycle  
is substituted with 0-3 R<sup>10b</sup>;

R<sup>10a</sup>, at each occurrence, is independently selected from H,  
C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>,  
20 CF<sub>3</sub>, or phenyl substituted with 0-4 R<sup>10b</sup>;

R<sup>10b</sup>, at each occurrence, is independently selected from H,  
OH, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, CN, NO<sub>2</sub>,  
NR<sup>15</sup>R<sup>16</sup>, or CF<sub>3</sub>;

25 R<sup>11</sup>, at each occurrence, is independently selected from  
H, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>18</sup>R<sup>19</sup>,  
C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, CF<sub>3</sub>;  
C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 0-3 R<sup>11a</sup>;  
30 C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>11b</sup>;  
C<sub>3</sub>-C<sub>10</sub> carbocycle substituted with 0-3 R<sup>11b</sup>; or  
5 to 10 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 10 membered heterocycle  
35 is substituted with 0-3 R<sup>11b</sup>;

- R<sup>11a</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, or phenyl substituted with 0-3 R<sup>11b</sup>;
- 5 R<sup>11b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy;
- 10 Z is H;  
C<sub>1</sub>-C<sub>6</sub> alkyl substituted with 0-3 R<sup>12a</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>12a</sup>; or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>12a</sup>;
- 15 R<sup>12a</sup>, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkyl, and C<sub>1</sub>-C<sub>4</sub> haloalkoxy;
- 20 R<sup>13</sup>, at each occurrence, is independently selected from H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;
- R<sup>14</sup> is H, phenyl, benzyl, C<sub>1</sub>-C<sub>6</sub> alkyl, or C<sub>2</sub>-C<sub>6</sub> alkoxyalkyl;
- 25 R<sup>14a</sup> is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;
- R<sup>15</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-;
- 30 R<sup>16</sup>, at each occurrence, is independently selected from H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-;
- 35 R<sup>17</sup> is H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkoxyalkyl, aryl substituted by 0-4 R<sup>17a</sup>, or -CH<sub>2</sub>-aryl substituted by 0-4 R<sup>17a</sup>;

R<sup>17a</sup> is H, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, butoxy, -OH, F, Cl, Br, I, CF<sub>3</sub>, OCF<sub>3</sub>, SCH<sub>3</sub>, S(O)CH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, -NH<sub>2</sub>, -N(CH<sub>3</sub>)<sub>2</sub>, or C<sub>1</sub>-C<sub>4</sub> haloalkyl;

5

R<sup>18</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-; and

10 R<sup>19</sup>, at each occurrence, is independently selected from H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>6</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>6</sub> alkyl)-S(=O)<sub>2</sub>-.

15 4. A compound according to Claim 3 of Formula (Ia) wherein:

R<sup>3</sup> is -(CHR<sup>7</sup>)<sub>n</sub>-R<sup>4</sup>,

20 n is 0 or 1;

R<sup>3a</sup> is H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, butoxy, allyl, or 3-buten-1-yl;

25 R<sup>4</sup> is H, OH, OR<sup>14a</sup>,C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-2 R<sup>4a</sup>,C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-2 R<sup>4a</sup>,C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>4a</sup>,C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,30 C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>4b</sup>, or

5 to 6 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R<sup>4b</sup>;

35

R<sup>4a</sup>, at each occurrence, is independently selected from is H, F, Cl, Br, I, CF<sub>3</sub>,

C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,

phenyl substituted with 0-3 R<sup>4b</sup>, or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
5 substituted with 0-3 R<sup>4b</sup>;

R<sup>4b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>,  
S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkoxy, C<sub>1</sub>-C<sub>2</sub>  
10 haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

R<sup>5</sup> is H, OR<sup>14</sup>;  
C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-3 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>5b</sup>;  
15 C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>5b</sup>;

R<sup>5a</sup> is H, methyl, ethyl, propyl, or butyl;

R<sup>5b</sup>, at each occurrence, is independently selected from:  
20 H, methyl, ethyl, propyl, butyl, CF<sub>3</sub>, OR<sup>14</sup>, Cl, F, Br,  
I, =O;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>5c</sup>;  
phenyl substituted with 0-3 R<sup>5c</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
25 heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>5c</sup>;

R<sup>5c</sup>, at each occurrence, is independently selected from H,  
30 OH, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>,  
S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkoxy, C<sub>1</sub>-C<sub>2</sub>  
haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

R<sup>6</sup> is H;  
35

R<sup>7</sup>, at each occurrence, is independently selected from H,  
F, CF<sub>3</sub>, methyl, and ethyl;

Ring B is a 7 membered lactam,  
wherein the lactam is saturated, partially saturated  
or unsaturated;  
wherein each additional lactam carbon is substituted  
5 with 0-2 R<sup>11</sup>; and,  
optionally, the lactam contains a heteroatom selected  
from -N=, -NH-, and -N(R<sup>10</sup>)-;

10 additionally, two R<sup>11</sup> substituents on adjacent atoms may be  
combined to form a benzo fused radical; wherein said  
benzo fused radical is substituted with 0-2 R<sup>13</sup>;

15 additionally, two R<sup>11</sup> substituents on adjacent atoms may be  
combined to form a 5 to 6 membered heteroaryl fused  
radical, wherein said 5 to 6 membered heteroaryl fused  
radical comprises 1 or 2 heteroatoms selected from N,  
O, and S; wherein said 5 to 6 membered heteroaryl  
fused radical is substituted with 0-2 R<sup>13</sup>;

20 additionally, two R<sup>11</sup> substituents on the same or adjacent  
carbon atoms may be combined to form a C<sub>3</sub>-C<sub>6</sub>  
carbocycle substituted with 0-2 R<sup>13</sup>;

R<sup>10</sup> is H, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>;  
25 C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with 0-1 R<sup>10a</sup>;  
phenyl substituted with 0-4 R<sup>10b</sup>;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>10b</sup>; or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
30 sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>10b</sup>;

R<sup>10a</sup>, at each occurrence, is independently selected from H,  
C<sub>1</sub>-C<sub>4</sub> alkyl, OR<sup>14</sup>, Cl, F, Br, I, =O, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>,  
35 CF<sub>3</sub>, or phenyl substituted with 0-4 R<sup>10b</sup>;

- $R^{10b}$ , at each occurrence, is independently selected from H, OH, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkoxy, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, or CF<sub>3</sub>;
- 5  $R^{11}$ , at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, =O, NR<sup>18</sup>R<sup>19</sup>, C(=O)R<sup>17</sup>, C(=O)OR<sup>17</sup>, CF<sub>3</sub>;  
C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with 0-3 R<sup>11a</sup>;  
C<sub>6</sub>-C<sub>10</sub> aryl substituted with 0-3 R<sup>11b</sup>;
- 10 C<sub>3</sub>-6 carbocycle substituted with 0-3 R<sup>11b</sup>; or  
5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R<sup>11b</sup>;
- 15  $R^{11a}$ , at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>4</sub> alkyl, OR<sup>14</sup>, F, =O, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, or phenyl substituted with 0-3 R<sup>11b</sup>;
- 20  $R^{11b}$ , at each occurrence, is independently selected from H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;
- Z is H;
- 25 C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-3 R<sup>12a</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3 R<sup>12a</sup>; or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3 R<sup>12a</sup>;
- $R^{12a}$ , at each occurrence, is independently selected from  
30 H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>3</sub> alkoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;
- $R^{13}$ , at each occurrence, is independently selected from  
35 H, OH, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, Cl, F, Br, I, CN, NO<sub>2</sub>, NR<sup>15</sup>R<sup>16</sup>, and CF<sub>3</sub>;
- R<sup>14</sup> is H, phenyl, benzyl, C<sub>1</sub>-C<sub>4</sub> alkyl, or C<sub>2</sub>-C<sub>4</sub> alkoxyalkyl;

R<sup>15</sup>, at each occurrence, is independently selected from H, C<sub>1</sub>-C<sub>4</sub> alkyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>4</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>4</sub> alkyl)-S(=O)<sub>2</sub>-;

5

R<sup>16</sup>, at each occurrence, is independently selected from H, OH, C<sub>1</sub>-C<sub>4</sub> alkyl, benzyl, phenethyl, (C<sub>1</sub>-C<sub>4</sub> alkyl)-C(=O)-, and (C<sub>1</sub>-C<sub>4</sub> alkyl)-S(=O)<sub>2</sub>-;

10 R<sup>17</sup> is H, methyl, ethyl, propyl, butyl, methoxymethyl, ethoxymethyl, methoxyethyl, ethoxyethyl, phenyl substituted by 0-3 R<sup>17a</sup>, or -CH<sub>2</sub>-phenyl substituted by 0-3 R<sup>17a</sup>;


15 R<sup>17a</sup> is H, methyl, methoxy, -OH, F, Cl, CF<sub>3</sub>, or OCF<sub>3</sub>;

R<sup>18</sup>, at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl;

20

R<sup>19</sup>, at each occurrence, is independently selected from H, methyl, and ethyl.

25 5. A compound of Claim 4 of Formula (Ib):




30


or a pharmaceutically acceptable salt or prodrug thereof wherein:

Ring B is selected from:

35



6. A compound according to Claim 5 of Formula (Ic):



(Ic)

or a pharmaceutically acceptable salt or prodrug thereof  
wherein

R<sup>3</sup> is R<sup>4</sup>,

R<sup>4</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>4a</sup>,  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>4a</sup>, or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>4a</sup>;

R<sup>4a</sup>, at each occurrence, is independently selected from  
H, F, CF<sub>3</sub>,



C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3 R<sup>4b</sup>,  
phenyl substituted with 0-3 R<sup>4b</sup>, or  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
5 sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>4b</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
10 imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>4b</sup>, at each occurrence, is independently selected from H,  
OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
15 ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

R<sup>5</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>5b</sup>;  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>5b</sup>;

20


R<sup>5b</sup>, at each occurrence, is independently selected from:  
H, methyl, ethyl, propyl, butyl, CF<sub>3</sub>, OR<sup>14</sup>, =O;  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-2 R<sup>5c</sup>;  
phenyl substituted with 0-3 R<sup>5c</sup>; or  
25 5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3 R<sup>5c</sup>; wherein said 5 to 6  
membered heterocycle is selected from pyridinyl,  
30 pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R<sup>5c</sup>, at each occurrence, is independently selected from H,  
35 OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>,  
S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;

- $R^{11}$ , at each occurrence, is independently selected from  
H, =O,  $NR^{18}R^{19}$ ,  $CF_3$ ;  
C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with 0-1  $R^{11a}$ ;  
phenyl substituted with 0-3  $R^{11b}$ ;  
5 C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3  $R^{11b}$ ; and  
5 to 6 membered heterocycle containing 1 to 4  
heteroatoms selected from nitrogen, oxygen, and  
sulphur, wherein said 5 to 6 membered heterocycle is  
substituted with 0-3  $R^{11b}$ ; wherein said 5 to 6  
10 membered heterocycle is selected from pyridinyl,  
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,  
pyrrolyl, piperazinyl, piperidinyl, pyrazolyl,  
imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;
- 15  $R^{11a}$ , at each occurrence, is independently selected from H,  
C<sub>1</sub>-C<sub>4</sub> alkyl,  $OR^{14}$ , F, Cl, =O,  $NR^{15}R^{16}$ ,  $CF_3$ , or phenyl  
substituted with 0-3  $R^{11b}$ ;
- $R^{11b}$ , at each occurrence, is independently selected from H,  
20 OH, Cl, F,  $NR^{15}R^{16}$ ,  $CF_3$ , methyl, ethyl, propyl, butyl,  
methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub>  
haloalkoxy;
- Z is H;  
25 C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-3  $R^{12a}$ ;  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-3  $R^{12a}$ ; or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-3  $R^{12a}$ ;
- $R^{12a}$ , at each occurrence, is independently selected from  
30 H, OH, Cl, F,  $NR^{15}R^{16}$ ,  $CF_3$ , acetyl,  $SCH_3$ ,  $S(=O)CH_3$ ,  
 $S(=O)_2CH_3$ , methyl, ethyl, propyl, butyl, methoxy,  
ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;
- $R^{13}$ , at each occurrence, is independently selected from  
35 H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy,  
Cl, F, Br, CN,  $NR^{15}R^{16}$ , and  $CF_3$ ;
- $R^{14}$  is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;

- $R^{15}$ , at each occurrence, is independently selected from H, methyl, ethyl, propyl, and butyl;
- 5  $R^{16}$ , at each occurrence, is independently selected from H, OH, methyl, ethyl, propyl, butyl, benzyl, phenethyl, methyl-C(=O)-, ethyl-C(=O)-, methyl-S(=O)<sub>2</sub>-, and ethyl-S(=O)<sub>2</sub>-;
- 10  $R^{18}$ , at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl; and
- $R^{19}$ , at each occurrence, is independently selected from  
15 H, methyl, and ethyl.

7. A compound according to Claim 5 of Formula (Id):



20

(Id)

or a pharmaceutically acceptable salt or prodrug thereof  
wherein:

25  $R^3$  is  $R^4$ ,

$R^4$  is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1  $R^{4a}$ ,  
C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1  $R^{4a}$ , or  
C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1  $R^{4a}$ ;

30

$R^{4a}$ , at each occurrence, is independently selected from  
H, F, CF<sub>3</sub>,  
C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-3  $R^{4b}$ ,  
phenyl substituted with 0-3  $R^{4b}$ , or

- 5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R<sup>4b</sup>; wherein said 5 to 6 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;
- 10 R<sup>4b</sup>, at each occurrence, is independently selected from H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;
- 15 R<sup>5</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl substituted with 0-1 R<sup>5b</sup>; C<sub>2</sub>-C<sub>4</sub> alkenyl substituted with 0-1 R<sup>5b</sup>; C<sub>2</sub>-C<sub>4</sub> alkynyl substituted with 0-1 R<sup>5b</sup>;
- R<sup>5b</sup>, at each occurrence, is independently selected from:
- 20 H, methyl, ethyl, propyl, butyl, CF<sub>3</sub>, OR<sup>14</sup>, =O; C<sub>3</sub>-C<sub>6</sub> carbocycle substituted with 0-2 R<sup>5c</sup>; phenyl substituted with 0-3 R<sup>5c</sup>; or 5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R<sup>5c</sup>; wherein said 5 to 6 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;
- 25 30
- R<sup>5c</sup>, at each occurrence, is independently selected from H, OH, Cl, F, NR<sup>15</sup>R<sup>16</sup>, CF<sub>3</sub>, acetyl, SCH<sub>3</sub>, S(=O)CH<sub>3</sub>, S(=O)<sub>2</sub>CH<sub>3</sub>, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C<sub>1</sub>-C<sub>2</sub> haloalkyl, and C<sub>1</sub>-C<sub>2</sub> haloalkoxy;
- 35
- R<sup>11</sup>, at each occurrence, is independently selected from H, =O, NR<sup>18</sup>R<sup>19</sup>, CF<sub>3</sub>;